




## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification:</b> C07D 263/32, A61K 31/421, A61K 31/422, A61K 31/426, A61K 31/427, C07D 277/24, C07D 413/12, C07D 417/12	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 00/08002</b> <b>(43) International Publication Date:</b> 17 February 2000 (17.02.2000)
<b>(21) International Application Number:</b> PCT/EP99/05666 <b>(22) International Filing Date:</b> 05 August 1999 (05.08.1999) <b>(30) Priority Data:</b> 9817118.4 07 August 1998 (07.08.1998) GB <b>(60) Parent Application or Grant</b> GLAXO GROUP LIMITED [/]; (). COLLINS, Jon, Loren [/]; (). DEZUBE, Milana [/]; (). OPLINGER, Jeffrey, Alan [/]; (). WILLSON, Timothy, Mark [/]; (). COLLINS, Jon, Loren [/]; (). DEZUBE, Milana [/]; (). OPLINGER, Jeffrey, Alan [/]; (). WILLSON, Timothy, Mark [/]; (). LEAROYD, Stephanie, Anne ; ().	<b>Published</b>	
<b>(54) Title:</b> SUBSTITUTED OXAZOLES AND THIAZOLES DERIVATIVES AS hPPAR GAMMA AND hPPAR ALPHA ACTIVATORS <b>(54) Titre:</b> OXAZOLES SUBSTITUES ET DERIVES DE THIAZOLES COMME ACTIVATEURS DE ALPHA hPPAR ET DE GAMMA hPPAR  <b>(57) Abstract</b> <p>The present invention discloses compounds of formula (I), and tautomeric forms, pharmaceutically acceptable salts, or solvates thereof. Preferably, the compounds of the invention are dual activators of hPPAR'gamma' and hPPAR'alpha'.</p> <b>(57) Abrégé</b> <p>L'invention concerne des composés représentés par la formule (I), et des formes tautomères, des sels pharmaceutiquement acceptables ou des solvates de ceux-ci. Les composés de l'invention sont de préférence des activateurs doubles de hPPAR'gamma' et hPPAR'alpha'.</p>		

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification <sup>7</sup> : C07D 263/32, 277/24, A61K 31/421, 31/426, C07D 413/12, 417/12, A61K 31/422, 31/427</p>	A1	<p>(11) International Publication Number: <b>WO 00/08002</b></p> <p>(43) International Publication Date: 17 February 2000 (17.02.00)</p>
<p>(21) International Application Number: PCT/EP99/05666</p> <p>(22) International Filing Date: 5 August 1999 (05.08.99)</p> <p>(30) Priority Data: 9817118.4 7 August 1998 (07.08.98) GB</p> <p>(71) Applicant (for all designated States except US): GLAXO GROUP LIMITED [GB/GB]; Glaxo Wellcome House, Berkeley Avenue, Greenford, Middlesex UB6 0NN (GB).</p> <p>(72) Inventors; and (75) Inventors/Applicants (for US only): COLLINS, Jon, Loren [US/US]; Glaxo Wellcome Inc., Five Moore Drive, Research Triangle Park, NC 27709 (US). DEZUBE, Milana [US/US]; Glaxo Wellcome Inc., Five Moore Drive, Research Triangle Park, NC 27709 (US). OPLINGER, Jeffrey, Alan [US/US]; Glaxo Wellcome Inc., Five Moore Drive, Research Triangle Park, NC 27709 (US). WILLSON, Timothy, Mark [GB/US]; Glaxo Wellcome Inc., Five Moore Drive, Research Triangle Park, NC 27709 (US).</p> <p>(74) Agent: LEAROYD, Stephanie, Anne; Glaxo Wellcome plc, Glaxo Wellcome House, Berkeley Avenue, Greenford, Middlesex UB6 0NN (GB).</p>		<p>(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SI, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p><b>Published</b> With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</p>
<p>(54) Title: SUBSTITUTED OXAZOLES AND THIAZOLES DERIVATIVES AS hPPAR GAMMA AND hPPAR ALPHA ACTIVATORS</p> <div style="text-align: center;">  <p>(I)</p> </div> <p>(57) Abstract</p> <p>The present invention discloses compounds of formula (I), and tautomeric forms, pharmaceutically acceptable salts, or solvates thereof. Preferably, the compounds of the invention are dual activators of hPPAR<math>\gamma</math> and hPPAR<math>\alpha</math>.</p>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

**Description**

5

10

15

20

25

30

35

40

45

50

55

## SUBSTITUTED OXAZOLES AND THIAZOLES DERIVATIVES AS hPPAR GAMMA AND hPPAR ALPHA ACTIVATORS

The present invention relates to certain novel compounds, to processes for their preparation, to pharmaceutical compositions containing them and to their use in medicine. More particularly, it relates to compounds which exhibit activation, including dual agonist activity, to peroxisome proliferator-activated receptors gamma (PPAR $\gamma$ ) and alpha (PPAR $\alpha$ ) thereby enabling them to modulate the blood glucose and lipid levels in mammals.

Treatment of type 2 diabetes mellitus usually begins with a combination of diet and exercise, with progression to oral hypoglycaemics (e.g. sulfonylureas) and in more severe cases, insulin. In the last decade a class of compounds known as thiazolidinediones (e.g. U.S. Pat Nos. 5,089,514, 4,342,771, 4,367,234, 4,340,605, 5,306,726) have emerged as effective antidiabetic agents that enhance the insulin sensitivity of target tissues (skeletal muscle, liver, adipose) in animal models of type 2 diabetes mellitus and also reduce lipid and insulin levels in these animal models.

It has been reported that thiazolidinediones are potent and selective activators of PPAR $\gamma$  and bind directly to the PPAR $\gamma$  receptor (J. M. Lehmann et al., *J. Biol. Chem.* 12953-12956, 270 (1995)), providing evidence that PPAR $\gamma$  is a possible target for the therapeutic actions of the thiazolidinediones.

PCT patent publication WO 97/31907 discloses certain novel compounds that bind to and activate PPAR $\gamma$ . These compounds are indicated to be useful for the treatment of type 2 diabetes mellitus and other diseases.

Activators of the nuclear receptor PPAR $\gamma$ , for example troglitazone, have been shown in the clinic to enhance insulin-action, reduce serum glucose and have small but significant effects on reducing serum triglyceride levels in patients with Type 2 diabetes. See, for example, D. E. Kelly et al., *Curr. Opin. Endocrinol. Diabetes*, 90-96, 5 (2), (1998); M. D. Johnson et al., *Ann. Pharmacother.*, 337-348, 32 (3), (1997); and M. Leutenegger et al., *Curr. Ther. Res.*, 403-416, 58 (7), (1997).

5 The mechanism for this triglyceride lowering effect appears to be  
predominantly increased clearance of very low density lipoproteins (VLDL)  
10 through induction of lipoprotein lipase (LPL) gene expression. See, for  
example, B. Staels et al., *Arterioscler. Thromb., Vasc. Biol.*, 1756-1764, 17  
5 (9), (1997).


Fibrates are a class of drugs which may lower serum triglycerides 20-  
15 50%, lower LDLc 10-15%, shift the LDL particle size from the more  
atherogenic small dense to normal dense LDL, and increase HDLc 10-15%.  
Experimental evidence indicates that the effects of fibrates on serum lipids are  
10 mediated through activation of PPAR $\alpha$ . See, for example, B. Staels et al.,  
20 *Curr. Pharm. Des.*, 1-14, 3 (1), (1997). Activation of PPAR $\alpha$  results in  
transcription of enzymes that increase fatty acid catabolism and decrease de-  
novo fatty acid synthesis in the liver resulting in decreased triglyceride  
25 synthesis and VLDL production/secretion. In addition, PPAR $\alpha$  activation  
15 decreases production of apoC-III. Reduction in apoC-III, an inhibitor of LPL  
activity, increases clearance of VLDL. See, for example, J. Auwerx et al.,  
*Atherosclerosis*, (Shannon, Ire.), S29-S37, 124 (Suppl), (1996).

30 In addition, a dual agonist of PPAR $\alpha$  and PPAR $\gamma$  could be effective in  
reducing the dyslipidemia and hyperinsulinemia associated with impaired  
20 glucose tolerance (IGT) or metabolic syndrome and could be effective in  
patients with mixed hyperlipidemia. See, for example, U.S. Pat Nos.  
35 5,478,852.

PCT patent publication WO 98/05331 (Paterniti et al.) relates to  
40 methods for treating diabetes and cardiovascular disease using a PPAR $\gamma$   
25 agonist in combination with a PPAR $\alpha$  agonist, or a compound that activates  
both PPAR $\gamma$  and PPAR $\alpha$ .

#### 45 Brief description

Briefly, in one aspect, the present invention provides compounds of  
30 formula (I), and tautomeric forms, pharmaceutically acceptable salts and  
solvates thereof,



(I)

wherein;

R<sup>1</sup> is hydrogen or C<sub>1-3</sub>alkyl;

R<sup>2</sup> is hydrogen, or C<sub>1-6</sub>alkyl optionally substituted by one or more halogens;

R<sup>3</sup> is C<sub>1-6</sub>alkyl, C<sub>4-7</sub>cycloalkyl or cycloalkenyl, -OC<sub>1-6</sub>alkyl, -NR'R' (where each R' is independently hydrogen or C<sub>1-3</sub>alkyl), a 5 or 6 membered heterocyclic group containing at least one oxygen, nitrogen, or sulfur ring atom (optionally substituted by one or more halogen, C<sub>1-6</sub>alkyl optionally substituted by one or more halogens, -OC<sub>1-6</sub>alkyl optionally substituted by one or more halogens, -CN, or -NO<sub>2</sub>), or phenyl (optionally substituted by one or more halogen, C<sub>1-6</sub>alkyl optionally substituted by one or more halogens, -OC<sub>1-6</sub>alkyl optionally substituted by one or more halogens, -CN, or -NO<sub>2</sub>);

R<sup>4</sup> is a 5 or 6 membered heterocyclic group containing at least one oxygen, nitrogen, or sulfur ring atom (optionally substituted by one or more halogen, C<sub>1-6</sub>alkyl optionally substituted by one or more halogens, -OC<sub>1-6</sub>alkyl optionally substituted by one or more halogens, -CN, or -NO<sub>2</sub>), or phenyl (optionally substituted by one or more halogen, C<sub>1-6</sub>alkyl optionally substituted by one or more halogens, -OC<sub>1-6</sub>alkyl optionally substituted by one or more halogens, -NR'R' (as defined above), -CN, or -NO<sub>2</sub>);

R<sup>5</sup> is hydrogen, halogen, or C<sub>1-3</sub>alkyl optionally substituted by one or more halogens;

R<sup>6</sup> is hydrogen or C<sub>1-3</sub>alkyl;

X is O or S; and

n is 1, 2, or 3.

5 Preferably, the compounds of this invention activate both the hPPAR $\gamma$  and hPPAR $\alpha$  receptors.

10 In another aspect, the present invention provides pharmaceutical compositions comprising a compound of the invention. As used herein, "a  
5 compound of the invention" means a compound of formula (I) or a tautomeric form, pharmaceutically acceptable salt, or solvates thereof.

15 The invention further provides a compound of the invention for use in therapy, and in particular, in human medicine.

20 In another aspect, the present invention provides a method for treatment or prevention of a hPPAR $\gamma$  and/or hPPAR $\alpha$  mediated disease, risk factor, or condition, comprising administration of a therapeutically effective amount of a compound of this invention.

25 According to another aspect, the present invention provides the use of a compound of the invention for the manufacture of a medicament for the  
15 treatment or prevention of a hPPAR $\gamma$  and/or hPPAR $\alpha$  mediated disease.

30 hPPAR $\gamma$  and/or hPPAR $\alpha$  mediated diseases, risk factors, or conditions include hyperglycemia, dyslipidemia, type II diabetes mellitus including associated diabetic dyslipidemia, type I diabetes, hypertriglyceridemia, syndrome X, insulin resistance, heart failure, hyperlipidemia,  
20 hypercholesteremia, hypertension, cardiovascular disease, including atherosclerosis, regulation of appetite and food intake in subjects suffering  
35 from disorders such as obesity, anorexia bulimia, and anorexia nervosa. In particular, the compounds of the present invention are useful in the treatment or prevention of hyperglycaemia, dyslipidemia, and type II diabetes mellitus  
40 25 including associated diabetic dyslipidemia.

#### Detailed Description

45 Preferably, R<sup>1</sup> is hydrogen or methyl. Most preferably, R<sup>1</sup> is hydrogen.

50 Preferably, R<sup>2</sup> is C<sub>1-8</sub>alkyl optionally substituted by one or more  
30 halogens. Preferably, said halogen is fluorine. Most preferably, R<sup>2</sup> is straight-chain.

55 Preferably, R<sup>3</sup> is pyridine, pyrazine, thiophene, furan, thiazole, or phenyl (any of which may be optionally substituted by one or more halogen,



C<sub>1-6</sub>alkyl optionally substituted by one or more halogens, -OC<sub>1-6</sub>alkyl optionally substituted by one or more halogens, -CN, or -NO<sub>2</sub>), or C<sub>4-7</sub>cycloalkyl. Most preferably, R<sup>3</sup> is phenyl (optionally substituted by one or more halogen, C<sub>1-6</sub>alkyl optionally substituted by one or more halogens, -OC<sub>1-6</sub>alkyl optionally substituted by one or more halogens, -CN, or -NO<sub>2</sub>).


Preferably R<sup>4</sup> is phenyl (optionally substituted by one or more halogen, C<sub>1-6</sub>alkyl optionally substituted by one or more halogens, or -OC<sub>1-6</sub>alkyl optionally substituted by one or more halogens). Preferably, said halogen is fluorine. Most preferably R<sup>4</sup> is phenyl either unsubstituted or substituted with 1, 2, or 3 fluorine atoms.

Preferably, R<sup>5</sup> is hydrogen, halogen, or C<sub>1-3</sub>alkyl optionally substituted by one or more halogens. Most preferably R<sup>5</sup> is hydrogen.

Preferably R<sup>6</sup> is methyl or ethyl.

Preferably n is 2.

Preferably, the carbon atom bonded to CO<sub>2</sub>R<sup>1</sup> is in the S configuration. In other words, preferably, the absolute configuration around that carbon is:



Suitable compounds of the present invention include:

- (2S)-2-[[[(Z)-1-methyl-3-oxo-3-phenyl-1-propenyl]amino]-3-{4-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]phenyl}propanoic acid
- (2S)-3-{4-[2-[2-(4-fluorophenyl)-5-methyl-1,3-oxazol-4-yl]ethoxy]phenyl}-2-[[[(Z)-1-methyl-3-oxo-3-[4-(trifluoromethyl)phenyl]-1-propenyl]amino]propanoic acid
- (2S)-3-{4-[2-[2-(4-isopropoxyphenyl)-5-methyl-1,3-oxazol-4-yl]ethoxy]phenyl}-2-[[[(Z)-1-methyl-3-oxo-3-[4-(trifluoromethyl)phenyl]-1-propenyl]amino]propanoic acid
- (2S)-3-{4-[2-[2-(4-methoxyphenyl)-5-methyl-1,3-oxazol-4-yl]ethoxy]phenyl}-2-[[[(Z)-1-methyl-3-oxo-3-[4-(trifluoromethyl)phenyl]-1-propenyl]amino]propanoic acid

(2S)-2-(((Z)-1-ethyl-3-oxo-3-phenyl-1-propenyl)amino)-3-[4-(2-(5-methyl-2-[4-(trifluoromethyl)phenyl]-1,3-oxazol-4-yl)ethoxy)phenyl]propanoic acid  
(2S)-2-(((Z)-1-ethyl-3-(4-fluorophenyl)-3-oxo-1-propenyl)amino)-3-[4-(2-[2-(4-methoxyphenyl)-5-methyl-1,3-oxazol-4-yl]ethoxy)phenyl]propanoic acid  
(2S)-2-(((Z)-1-methyl-3-oxo-3-phenyl-1-propenyl)amino)-3-[4-(2-(5-methyl-2-[4-(trifluoromethyl)phenyl]-1,3-oxazol-4-yl)ethoxy)phenyl]propanoic acid  
(2S)-2-(((Z)-1-ethyl-3-oxo-3-phenyl-1-propenyl)amino)-3-[4-(2-[2-(4-fluorophenyl)-5-methyl-1,3-oxazol-4-yl]ethoxy)phenyl]propanoic acid  
(2S)-2-(((Z)-1-ethyl-3-(4-fluorophenyl)-3-oxo-1-propenyl)amino)-3-[4-(2-[2-(4-fluorophenyl)-5-methyl-1,3-oxazol-4-yl]ethoxy)phenyl]propanoic acid  
(2S)-2-(((Z)-1-methyl-3-oxo-3-[4-(trifluoromethyl)phenyl]-1-propenyl)amino)-3-[4-(2-(5-methyl-2-phenyl-1,3-thiazol-4-yl)ethoxy)phenyl]propanoic acid  
(2S)-2-(((Z)-3-(4-fluorophenyl)-1-methyl-3-oxo-1-propenyl)amino)-3-[4-(2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy)phenyl]propanoic acid  
(2S)-2-(((Z)-1-methyl-3-oxo-3-(2,3,4-trifluorophenyl)-1-propenyl)amino)-3-[4-(2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy)phenyl]propanoic acid  
(2S)-2-(((Z)-1-methyl-3-(4-nitrophenyl)-3-oxo-1-propenyl)amino)-3-[4-(2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy)phenyl]propanoic acid  
(2S)-2-(((Z)-1-methyl-3-oxo-3-[4-(trifluoromethyl)phenyl]-1-propenyl)amino)-3-[4-(2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy)phenyl]propanoic acid  
(2S)-2-(((Z)-1-ethyl-3-(4-fluorophenyl)-3-oxo-1-propenyl)amino)-3-[4-(2-[2-(4-isopropoxyphenyl)-5-methyl-1,3-oxazol-4-yl]ethoxy)phenyl]propanoic acid  
(2S)-2-(((Z)-1-methyl-3-oxo-3-(2,4,5-trifluorophenyl)-1-propenyl)amino)-3-[4-(2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy)phenyl]propanoic acid  
(2S)-2-(((Z)-1-ethyl-3-oxo-3-phenyl-1-propenyl)amino)-3-[4-(2-[2-(4-isopropoxyphenyl)-5-methyl-1,3-oxazol-4-yl]ethoxy)phenyl]propanoic acid  
(2S)-2-(((Z)-1-ethyl-3-oxo-3-[4-(trifluoromethyl)phenyl]-1-propenyl)amino)-3-[4-(2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy)phenyl]propanoic acid  
(2S)-2-(((Z)-1-methyl-3-(4-methylphenyl)-3-oxo-1-propenyl)amino)-3-[4-(2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy)phenyl]propanoic acid  
(2S)-2-(((Z)-1-ethyl-3-oxo-3-phenyl-1-propenyl)amino)-3-[4-(2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy)phenyl]propanoic acid

(2S)-3-(4-{2-[2-(4-fluorophenyl)-5-methyl-1,3-thiazol-4-yl]ethoxy}phenyl)-2-  
{[(Z)-3-oxo-3-phenyl-1-(trifluoromethyl)-1-propenyl]amino}propanoic acid  
(2S)-2-([(Z)-1-ethyl-3-oxo-3-[4-(trifluoromethyl)phenyl]-1-propenyl]amino)-3-  
(4-{2-[2-(4-fluorophenyl)-5-methyl-1,3-thiazol-4-yl]ethoxy}phenyl)propanoic  
acid  
(2S)-2-([(Z)-1-butyl-3-oxo-3-phenyl-1-propenyl]amino)-3-(4-{2-(5-methyl-2-  
phenyl-1,3-oxazol-4-yl)ethoxy}phenyl)propanoic acid  
(2S)-2-([(Z)-3-(4-chlorophenyl)-1-methyl-3-oxo-1-propenyl]amino)-3-(4-{2-(5-  
methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy}phenyl)propanoic acid  
(2S)-2-([(Z)-1-methyl-3-(3-nitrophenyl)-3-oxo-1-propenyl]amino)-3-(4-{2-(5-  
methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy}phenyl)propanoic acid  
(2S)-2-([(Z)-3-[2-fluoro-3-(trifluoromethyl)phenyl]-1-methyl-3-oxo-1-  
propenyl]amino)-3-(4-{2-(5-methyl-2-phenyl-1,3-oxazol-4-  
yl)ethoxy}phenyl)propanoic acid  
(2S)-2-([(Z)-3-(4-isopropoxyphenyl)-1-methyl-3-oxo-1-propenyl]amino)-3-(4-{2-  
(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy}phenyl)propanoic acid  
(2S)-2-([(Z)-3-(2-chlorophenyl)-1-methyl-3-oxo-1-propenyl]amino)-3-(4-{2-(5-  
methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy}phenyl)propanoic acid  
(2S)-2-([(Z)-3-(2-furyl)-1-methyl-3-oxo-1-propenyl]amino)-3-(4-{2-(5-methyl-2-  
phenyl-1,3-oxazol-4-yl)ethoxy}phenyl)propanoic acid  
(2S)-2-([(Z)-1-methyl-3-oxo-3-(2-pyrazinyl)-1-propenyl]amino)-3-(4-{2-(5-  
methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy}phenyl)propanoic acid  
(2S)-2-([(Z)-3-(2,4-difluorophenyl)-1-methyl-3-oxo-1-propenyl]amino)-3-(4-{2-  
(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy}phenyl)propanoic acid  
(2S)-2-([(Z)-1-methyl-3-oxo-3-(1,3-thiazol-2-yl)-1-propenyl]amino)-3-(4-{2-(5-  
methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy}phenyl)propanoic acid  
(2S)-2-([(Z)-1-methyl-3-oxo-3-(3-thienyl)-1-propenyl]amino)-3-(4-{2-(5-methyl-  
2-phenyl-1,3-oxazol-4-yl)ethoxy}phenyl)propanoic acid  
(2S)-2-([(Z)-1-methyl-3-oxo-3-(2-pyridinyl)-1-propenyl]amino)-3-(4-{2-(5-  
methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy}phenyl)propanoic acid  
(2S)-2-([(Z)-1-ethyl-3-(4-fluorophenyl)-3-oxo-1-propenyl]amino)-3-(4-{2-(5-  
methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy}phenyl)propanoic acid

- (2S)-2-([(Z)-1-methyl-3-oxo-3-phenyl-1-propenyl]amino)-3-{4-[2-(5-methyl-2-phenyl-1,3-thiazol-4-yl)ethoxy]phenyl}propanoic acid
- (2S)-2-([(Z)-3-(2-fluorophenyl)-1-methyl-3-oxo-1-propenyl]amino)-3-{4-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]phenyl}propanoic acid
- (2S)-2-([(Z)-3-(2,3-difluorophenyl)-1-methyl-3-oxo-1-propenyl]amino)-3-{4-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]phenyl}propanoic acid
- (2S)-2-([(Z)-3-(2-hydroxyphenyl)-1-methyl-3-oxo-1-propenyl]amino)-3-{4-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]phenyl}propanoic acid
- (2S)-3-{4-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]phenyl}-2-([(Z)-3-oxo-3-phenyl-1-propyl-1-propenyl]amino)propanoic acid
- (2S)-2-([(Z)-3-(4-methoxyphenyl)-1-methyl-3-oxo-1-propenyl]amino)-3-{4-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]phenyl}propanoic acid
- (2S)-3-{4-[2-[2-(4-methoxyphenyl)-5-methyl-1,3-oxazol-4-yl]ethoxy]phenyl}-2-([(Z)-1-methyl-3-oxo-3-phenyl-1-propenyl]amino)propanoic acid
- (2S)-2-([(Z)-3-cyclohexyl-1-methyl-3-oxo-1-propenyl]amino)-3-{4-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]phenyl}propanoic acid
- (2S)-3-{4-[2-[2-(4-isopropoxyphenyl)-5-methyl-1,3-oxazol-4-yl]ethoxy]phenyl}-2-([(Z)-1-methyl-3-oxo-3-phenyl-1-propenyl]amino)propanoic acid
- (2S)-2-([(Z)-1-heptyl-3-oxo-3-phenyl-1-propenyl]amino)-3-{4-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]phenyl}propanoic acid
- (2S)-2-([(Z)-1-methyl-3-(3-methylphenyl)-3-oxo-1-propenyl]amino)-3-{4-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]phenyl}propanoic acid
- (2S)-2-([(Z)-1-ethyl-3-oxo-3-phenyl-1-propenyl]amino)-3-{4-[2-[2-(4-methoxyphenyl)-5-methyl-1,3-oxazol-4-yl]ethoxy]phenyl}propanoic acid
- (2S)-3-{4-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]phenyl}-2-([(Z)-3-oxo-3-phenyl-1-(trifluoromethyl)-1-propenyl]amino)propanoic acid
- (2S)-3-{4-[2-(5-methyl-2-phenyl-1,3-thiazol-4-yl)ethoxy]phenyl}-2-([(Z)-3-oxo-3-phenyl-1-(trifluoromethyl)-1-propenyl]amino)propanoic acid
- (2S)-2-([(Z)-1-ethyl-3-oxo-3-{4-(trifluoromethyl)phenyl}-1-propenyl]amino)-3-{4-[2-[2-(4-fluorophenyl)-5-methyl-1,3-oxazol-4-yl]ethoxy]phenyl}propanoic acid

(2S)-2-(((Z)-1-ethyl-3-oxo-3-[4-(trifluoromethyl)phenyl]-1-propenyl)amino)-3-(4-[2-[2-(4-isopropoxyphenyl)-5-methyl-1,3-oxazol-4-yl]ethoxy]phenyl)propanoic acid

(2S)-2-(((Z)-1-ethyl-3-oxo-3-[4-(trifluoromethyl)phenyl]-1-propenyl)amino)-3-(4-[2-[2-(4-methoxyphenyl)-5-methyl-1,3-oxazol-4-yl]ethoxy]phenyl)propanoic acid

(2S)-3-(4-[2-[2-(4-fluorophenyl)-5-methyl-1,3-oxazol-4-yl]ethoxy]phenyl)-2-(((Z)-3-oxo-3-phenyl-1-(trifluoromethyl)-1-propenyl)amino)propanoic acid

(2S)-2-(((Z)-1-methyl-3-oxo-3-phenyl-1-propenyl)amino)-3-(4-[(5-methyl-2-phenyl-1,3-oxazol-4-yl)methoxy]phenyl)propanoic acid

(2S)-3-(4-[2-(5-ethyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]phenyl)-2-(((Z)-1-methyl-3-oxo-3-phenyl-1-propenyl)amino)propanoic acid

(2S)-2-(((Z)-1-methyl-3-oxo-3-phenyl-1-propenyl)amino)-3-(4-[2-(2-phenyl-5-propyl-1,3-oxazol-4-yl)ethoxy]phenyl)propanoic acid

(2S)-2-(((Z)-1-methyl-3-oxo-3-phenyl-1-propenyl)amino)-3-(4-[3-(5-methyl-2-phenyl-1,3-oxazol-4-yl)propoxy]phenyl)propanoic acid

(2S)-3-(4-[2-(5-ethyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]phenyl)-2-(((Z)-3-oxo-3-phenyl-1-(trifluoromethyl)-1-propenyl)amino)propanoic acid

(2S)-3-(4-[3-(5-methyl-2-phenyl-1,3-oxazol-4-yl)propoxy]phenyl)-2-(((Z)-3-oxo-3-phenyl-1-(trifluoromethyl)-1-propenyl)amino)propanoic acid

(2S)-3-(4-[2-(5-ethyl-2-phenyl-1,3-thiazol-4-yl)ethoxy]phenyl)-2-(((Z)-3-oxo-3-phenyl-1-(trifluoromethyl)-1-propenyl)amino)propanoic acid

(2S)-3-(4-[2-[5-ethyl-2-(4-fluorophenyl)-1,3-oxazol-4-yl]ethoxy]phenyl)-2-(((Z)-3-oxo-3-phenyl-1-(trifluoromethyl)-1-propenyl)amino)propanoic acid

(2S)-3-(4-[3-[5-ethyl-2-(4-fluorophenyl)-1,3-oxazol-4-yl]propoxy]phenyl)-2-(((Z)-3-oxo-3-phenyl-1-(trifluoromethyl)-1-propenyl)amino)propanoic acid

(2S)-3-(4-[2-[5-ethyl-2-(4-fluorophenyl)-1,3-thiazol-4-yl]ethoxy]phenyl)-2-(((Z)-3-oxo-3-phenyl-1-(trifluoromethyl)-1-propenyl)amino)propanoic acid

(2S)-3-(4-[3-[5-ethyl-2-(4-fluorophenyl)-1,3-oxazol-4-yl]propoxy]phenyl)-2-(((Z)-1-ethyl-3-oxo-3-[4-(trifluoromethyl)phenyl]-1-propenyl)amino)propanoic acid

(2S)-2-(((Z)-1-ethyl-3-oxo-3-[4-(trifluoromethyl)phenyl]-1-propenyl)amino)-3-(4-[3-(5-ethyl-2-phenyl-1,3-oxazol-4-yl)propoxy]phenyl)propanoic acid

(2S)-3-(4-{3-[5-ethyl-2-(4-fluorophenyl)-1,3-oxazol-4-yl]propoxy}phenyl)-2-  
(((Z)-1-methyl-3-oxo-3-[4-(trifluoromethyl)phenyl]-1-propenyl)amino)propanoic  
acid

(2S)-3-(4-{3-[5-ethyl-2-phenyl-1,3-oxazol-4-yl]propoxy}phenyl)-2-(((Z)-1-  
methyl-3-oxo-3-[4-(trifluoromethyl)phenyl]-1-propenyl)amino)propanoic acid

(2S)-3-(4-{3-[5-ethyl-2-phenyl-1,3-oxazol-4-yl]propoxy}phenyl)-2-(((Z)-3-oxo-3-  
phenyl-1-(trifluoromethyl)-1-propenyl)amino)propanoic acid

(2S)-3-(4-{[(5-ethyl-2-phenyl-1,3-thiazol-4-yl)methoxy]phenyl}-2-(((Z)-3-oxo-3-  
phenyl-1-(trifluoromethyl)-1-propenyl)amino)propanoic acid

(2S)-3-(4-{[(5-ethyl-2-phenyl-1,3-thiazol-4-yl)methoxy]phenyl}-2-(((Z)-1-methyl-  
3-oxo-3-[4-(trifluoromethyl)phenyl]-1-propenyl)amino)propanoic acid

(2S)-3-(4-{[(5-ethyl-2-phenyl-1,3-oxazol-4-yl)methoxy]phenyl}-2-(((Z)-3-oxo-3-  
phenyl-1-(trifluoromethyl)-1-propenyl)amino)propanoic acid

(2S)-3-(4-{[5-ethyl-2-(4-fluorophenyl)-1,3-oxazol-4-yl]methoxy}phenyl)-2-(((Z)-  
3-oxo-3-phenyl-1-(trifluoromethyl)-1-propenyl)amino)propanoic acid

(2S)-2-(((Z)-1-ethyl-3-oxo-3-[4-(trifluoromethyl)phenyl]-1-propenyl)amino)-3-  
{4-[(5-ethyl-2-phenyl-1,3-oxazol-4-yl)methoxy]phenyl}propanoic acid

(2S)-3-(4-{[5-ethyl-2-(4-fluorophenyl)-1,3-oxazol-4-yl]methoxy}phenyl)-2-(((Z)-  
1-ethyl-3-oxo-3-[4-(trifluoromethyl)phenyl]-1-propenyl)amino)propanoic acid

(2S)-3-(4-{[(5-ethyl-2-phenyl-1,3-oxazol-4-yl)methoxy]phenyl}-2-(((Z)-1-methyl-  
3-oxo-3-[4-(trifluoromethyl)phenyl]-1-propenyl)amino)propanoic acid

(2S)-3-(4-{[5-ethyl-2-(4-fluorophenyl)-1,3-oxazol-4-yl]methoxy}phenyl)-2-(((Z)-  
1-methyl-3-oxo-3-[4-(trifluoromethyl)phenyl]-1-propenyl)amino)propanoic acid

(2S)-2-(((Z)-1-ethyl-3-oxo-3-[4-(trifluoromethyl)phenyl]-1-propenyl)amino)-3-  
{4-[(5-ethyl-2-phenyl-1,3-thiazol-4-yl)methoxy]phenyl}propanoic acid

(2S)-3-(4-{2-[5-ethyl-2-(4-fluorophenyl)-1,3-oxazol-4-yl]ethoxy}phenyl)-2-(((Z)-  
1-ethyl-3-oxo-3-[4-(trifluoromethyl)phenyl]-1-propenyl)amino)propanoic acid

(2S)-3-(4-{2-[5-ethyl-2-(4-fluorophenyl)-1,3-thiazol-4-yl]ethoxy}phenyl)-2-(((Z)-  
1-ethyl-3-oxo-3-[4-(trifluoromethyl)phenyl]-1-propenyl)amino)propanoic acid

(2S)-2-(((Z)-1-ethyl-3-oxo-3-[4-(trifluoromethyl)phenyl]-1-propenyl)amino)-3-  
{4-[2-(5-ethyl-2-phenyl-1,3-thiazol-4-yl)ethoxy]phenyl}propanoic acid,

and pharmaceutically acceptable salts and solvates thereof.

Preferred compounds of the invention include

5 (2S)-2-(((Z)-1-methyl-3-oxo-3-phenyl-1-propenyl)amino)-3-{4-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]phenyl}propanoic acid

10 (2S)-2-(((Z)-1-ethyl-3-oxo-3-phenyl-1-propenyl)amino)-3-{4-[2-(5-methyl-2-{4-(trifluoromethyl)phenyl}-1,3-oxazol-4-yl)ethoxy]phenyl}propanoic acid

5 (2S)-2-(((Z)-1-methyl-3-oxo-3-(2,3,4-trifluorophenyl)-1-propenyl)amino)-3-{4-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]phenyl}propanoic acid

15 (2S)-2-(((Z)-1-methyl-3-(4-nitrophenyl)-3-oxo-1-propenyl)amino)-3-{4-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]phenyl}propanoic acid

(2S)-2-(((Z)-1-methyl-3-oxo-3-{4-(trifluoromethyl)phenyl}-1-propenyl)amino)-3-{4-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]phenyl}propanoic acid

20 (2S)-2-(((Z)-1-methyl-3-oxo-3-(2,4,5-trifluorophenyl)-1-propenyl)amino)-3-{4-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]phenyl}propanoic acid

(2S)-2-(((Z)-1-ethyl-3-oxo-3-phenyl-1-propenyl)amino)-3-{4-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]phenyl}propanoic acid

25 (2S)-3-{4-[2-[2-(4-fluorophenyl)-5-methyl-1,3-thiazol-4-yl]ethoxy]phenyl}-2-(((Z)-3-oxo-3-phenyl-1-(trifluoromethyl)-1-propenyl)amino)propanoic acid

(2S)-2-(((Z)-1-ethyl-3-oxo-3-{4-(trifluoromethyl)phenyl}-1-propenyl)amino)-3-{4-[2-[2-(4-fluorophenyl)-5-methyl-1,3-thiazol-4-yl]ethoxy]phenyl}propanoic acid

20 (2S)-2-(((Z)-1-butyl-3-oxo-3-phenyl-1-propenyl)amino)-3-{4-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]phenyl}propanoic acid

35 (2S)-2-(((Z)-3-(4-chlorophenyl)-1-methyl-3-oxo-1-propenyl)amino)-3-{4-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]phenyl}propanoic acid

(2S)-3-{4-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]phenyl}-2-(((Z)-3-oxo-3-phenyl-1-propyl-1-propenyl)amino)propanoic acid

40 (2S)-3-{4-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]phenyl}-2-(((Z)-3-oxo-3-phenyl-1-(trifluoromethyl)-1-propenyl)amino)propanoic acid

(2S)-2-(((Z)-1-ethyl-3-oxo-3-{4-(trifluoromethyl)phenyl}-1-propenyl)amino)-3-{4-[2-[2-(4-fluorophenyl)-5-methyl-1,3-oxazol-4-yl]ethoxy]phenyl}propanoic acid

30 acid

(2S)-2-(((Z)-1-ethyl-3-oxo-3-{4-(trifluoromethyl)phenyl}-1-propenyl)amino)-3-{4-[2-[2-(4-methoxyphenyl)-5-methyl-1,3-oxazol-4-yl]ethoxy]phenyl}propanoic acid

5

(2S)-3-(4-{2-[2-(4-fluorophenyl)-5-methyl-1,3-oxazol-4-yl]ethoxy}phenyl)-2-  
{[(Z)-3-oxo-3-phenyl-1-(trifluoromethyl)-1-propenyl]amino}propanoic acid

10

(2S)-2-([(Z)-1-methyl-3-oxo-3-phenyl-1-propenyl]amino)-3-{4-[(5-methyl-2-  
phenyl-1,3-oxazol-4-yl)methoxy]phenyl}propanoic acid

5

(2S)-3-(4-[2-(5-ethyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]phenyl)-2-([(Z)-1-methyl-  
3-oxo-3-phenyl-1-propenyl]amino)propanoic acid

15

(2S)-2-([(Z)-1-methyl-3-oxo-3-phenyl-1-propenyl]amino)-3-{4-[3-(5-methyl-2-  
phenyl-1,3-oxazol-4-yl)propoxy]phenyl}propanoic acid

10

(2S)-3-(4-[2-(5-ethyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]phenyl)-2-([(Z)-3-oxo-3-  
phenyl-1-(trifluoromethyl)-1-propenyl]amino)propanoic acid

20

(2S)-3-(4-[2-(5-ethyl-2-phenyl-1,3-thiazol-4-yl)ethoxy]phenyl)-2-([(Z)-3-oxo-3-  
phenyl-1-(trifluoromethyl)-1-propenyl]amino)propanoic acid

(2S)-3-(4-{2-[5-ethyl-2-(4-fluorophenyl)-1,3-oxazol-4-yl]ethoxy}phenyl)-2-([(Z)-  
3-oxo-3-phenyl-1-(trifluoromethyl)-1-propenyl]amino)propanoic acid

25

15

(2S)-3-(4-{3-[5-ethyl-2-(4-fluorophenyl)-1,3-oxazol-4-yl]propoxy}phenyl)-2-  
{[(Z)-3-oxo-3-phenyl-1-(trifluoromethyl)-1-propenyl]amino}propanoic acid

(2S)-3-(4-{2-[5-ethyl-2-(4-fluorophenyl)-1,3-thiazol-4-yl]ethoxy}phenyl)-2-([(Z)-  
3-oxo-3-phenyl-1-(trifluoromethyl)-1-propenyl]amino)propanoic acid

30

(2S)-3-(4-{3-[5-ethyl-2-(4-fluorophenyl)-1,3-oxazol-4-yl]propoxy}phenyl)-2-

20

([(Z)-1-ethyl-3-oxo-3-[4-(trifluoromethyl)phenyl]-1-propenyl]amino)propanoic  
acid

35

(2S)-2-([(Z)-1-ethyl-3-oxo-3-[4-(trifluoromethyl)phenyl]-1-propenyl]amino)-3-  
{4-[3-(5-ethyl-2-phenyl-1,3-oxazol-4-yl)propoxy]phenyl}propanoic acid

(2S)-3-(4-{3-[5-ethyl-2-(4-fluorophenyl)-1,3-oxazol-4-yl]propoxy}phenyl)-2-

40

25

([(Z)-1-methyl-3-oxo-3-[4-(trifluoromethyl)phenyl]-1-propenyl]amino)propanoic  
acid

(2S)-3-(4-[3-(5-ethyl-2-phenyl-1,3-oxazol-4-yl)propoxy]phenyl)-2-([(Z)-1-  
methyl-3-oxo-3-[4-(trifluoromethyl)phenyl]-1-propenyl]amino)propanoic acid

45

(2S)-3-(4-[(5-ethyl-2-(4-fluorophenyl)-1,3-oxazol-4-yl)methoxy]phenyl)-2-([(Z)-  
3-oxo-3-phenyl-1-(trifluoromethyl)-1-propenyl]amino)propanoic acid

30

(2S)-3-(4-[(5-ethyl-2-phenyl-1,3-oxazol-4-yl)methoxy]phenyl)-2-([(Z)-1-methyl-  
3-oxo-3-[4-(trifluoromethyl)phenyl]-1-propenyl]amino)propanoic acid

50

55



(2S)-3-(4-[[5-ethyl-2-(4-fluorophenyl)-1,3-oxazol-4-yl]methoxy]phenyl)-2-(((Z)-1-methyl-3-oxo-3-[4-(trifluoromethyl)phenyl]-1-propenyl)amino)propanoic acid  
(2S)-3-(4-{2-[5-ethyl-2-(4-fluorophenyl)-1,3-thiazol-4-yl]ethoxy}phenyl)-2-(((Z)-1-ethyl-3-oxo-3-[4-(trifluoromethyl)phenyl]-1-propenyl)amino)propanoic acid  
and pharmaceutically acceptable salts and solvates thereof.

Many of the compounds of formula (I) are dual activators of hPPAR $\gamma$  and hPPAR $\alpha$ . As used herein, by "activating compound", or "activator", or the like, is meant those compounds which achieve at least 50% activation of human PPAR $\gamma$  ("hPPAR $\gamma$ ") or hPPAR $\alpha$  (relative to the appropriate indicated positive control) in the transfection assay described below at concentrations of  $10^{-7}$  M or less. As used herein, a "dual activator" is a compound that is an activator of both PPAR $\gamma$  and PPAR $\alpha$ .

The ratio  $\frac{EC_{50} \text{ hPPAR}\alpha}{EC_{50} \text{ hPPAR}\gamma}$

defines the relative activity of hPPAR $\alpha$  to hPPAR $\gamma$ . Preferably, the dual activators of this invention have a relative hPPAR $\alpha$  to hPPAR $\gamma$  activity of from 0.02 to 50.

Those skilled in the art will recognize that stereocenters exist in compounds of formula (I). Accordingly, the present invention includes all possible stereoisomers and geometric isomers of formula (I) and includes not only racemic compounds but also the optically active isomers as well. When a compound of formula (I) is desired as a single enantiomer, it may be obtained either by resolution of the final product or by stereospecific synthesis from either isomerically pure starting material or any convenient intermediate. Resolution of the final product, an intermediate or a starting material may be effected by any suitable method known in the art. See, for example, Stereochemistry of Carbon Compounds by E. L. Eliel (McGraw Hill, 1962) and Tables of Resolving Agents by S. H. Wilen. Additionally, in situations where tautomers of the compounds of formula (I) are possible, the present invention is intended to include all tautomeric forms of the compounds.

It will also be appreciated by those skilled in the art that the compounds of the present invention may also be utilized in the form of a pharmaceutically

5 acceptable salt or solvate thereof. The physiologically acceptable salts of the compounds of formula (I) include conventional salts formed from pharmaceutically acceptable inorganic or organic acids or bases as well as quaternary ammonium acid addition salts. More specific examples of suitable acid salts include hydrochloric, hydrobromic, sulfuric, phosphoric, nitric, perchloric, fumaric, acetic, propionic, succinic, glycolic, formic, lactic, maleic, tartaric, citric, pantoic, malonic, hydroxymaleic, phenylacetic, glutamic, benzoic, salicylic, fumaric, toluenesulfonic, methanesulfonic, naphthalene-2-sulfonic, benzenesulfonic hydroxynaphthoic, hydroiodic, malic, stearic, tannic and the like. Other acids such as oxalic, while not in themselves pharmaceutically acceptable, may be useful in the preparation of salts useful as intermediates in obtaining the compounds of the invention and their pharmaceutically acceptable salts. More specific examples of suitable basic salts include sodium, lithium, potassium, magnesium, aluminium, calcium, zinc, N,N'-dibenzylethylenediamine, chlorprocaine, choline, diethanolamine, ethylenediamine, N-methylglucamine and procaine salts. References hereinafter to a compound according to the invention include both compounds of formula (I) and their pharmaceutically acceptable salts and solvates.

The terms C<sub>1-3</sub>alkyl, C<sub>4-6</sub>cycloalkyl, C<sub>1-6</sub>alkylene, C<sub>2-6</sub>alkenyl and the like, as used herein, indicate groups that may contain the indicated range of carbon atoms, for example 1 to 3 carbon atoms. Unless otherwise indicated, such groups can be straight chained or branched.

The term 5- or 6-membered heterocyclic group as used herein includes 5- or 6-membered substituted or unsubstituted heterocycloalkyl and heteroaryl groups, e.g. substituted or unsubstituted imidazolidinyl, piperidyl, piperazinyl, pyrrolidinyl, morpholinyl, pyridyl, pyridazinyl, pyrimidinyl, pyrazinyl, pyrrolyl, pyrazolyl, imidazolyl, pyranyl, furyl, thienyl, oxazolyl, isoxazolyl, oxadiazolyl, thiazolyl, thiadiazolyl, triazolyl or tetrazolyl. Particularly preferred heterocycles are pyridine, pyrazine, thiophene, furan, and thiazole.

It will be appreciated by those skilled in the art that reference herein to treatment extends to prophylaxis as well as the treatment of established diseases or symptoms. Moreover, it will be appreciated that the amount of a compound of the invention required for use in treatment will vary with the

5 nature of the condition being treated and the age and the condition of the  
patient and will be ultimately at the discretion of the attendant physician or  
veterinarian. In general, however, doses employed for adult human treatment  
10 will typically be in the range of 0.02 - 5000 mg per day, preferably 1-1500 mg  
5 per day. The desired dose may conveniently be presented in a single dose or  
as divided doses administered at appropriate intervals, for example as two,  
three, four or more sub-doses per day.

15 While it is possible that compounds of the present invention may be  
therapeutically administered as the raw chemical, it is preferable to present  
10 the active ingredient as a pharmaceutical formulation. Accordingly, the  
present invention further provides for a pharmaceutical formulation comprising  
20 a compound of formula (I) or a pharmaceutically acceptable salt or solvate  
thereof together with one or more pharmaceutically acceptable carriers  
therefor and, optionally, other therapeutic and/or prophylactic ingredients.

25 15 Formulations of the present invention include those especially  
formulated for oral, buccal, parenteral, transdermal, inhalation, intranasal,  
transmucosal, implant, or rectal administration, however, oral administration is  
30 preferred. For buccal administration, the formulation may take the form of  
tablets or lozenges formulated in conventional manner. Tablets and capsules  
20 for oral administration may contain conventional excipients such as binding  
agents, (for example, syrup, acacia, gelatin, sorbitol, tragacanth, mucilage of  
35 starch or polyvinylpyrrolidone), fillers (for example, lactose, sugar,  
microcrystalline cellulose, maize-starch, calcium phosphate or sorbitol),  
lubricants (for example, magnesium stearate, stearic acid, talc, polyethylene  
40 25 glycol or silica), disintegrants (for example, potato starch or sodium starch  
glycollate) or wetting agents, such as sodium lauryl sulfate. The tablets may  
be coated according to methods well-known in the art.

45 Alternatively, the compounds of the present invention may be  
incorporated into oral liquid preparations such as aqueous or oily  
30 suspensions, solutions, emulsions, syrups or elixirs, for example. Moreover,  
formulations containing these compounds may be presented as a dry product  
for constitution with water or other suitable vehicle before use. Such liquid  
50 preparations may contain conventional additives such as suspending agents

5 such as sorbitol syrup, methyl cellulose, glucose/sugar syrup, gelatin,  
hydroxyethylcellulose, carboxymethyl cellulose, aluminum stearate gel or  
hydrogenated edible fats; emulsifying agents such as lecithin, sorbitan mono-  
10 oleate or acacia; non-aqueous vehicles (which may include edible oils) such  
5 as almond oil, fractionated coconut oil, oily esters, propylene glycol or ethyl  
alcohol; and preservatives such as methyl or propyl p-hydroxybenzoates or  
sorbic acid. Such preparations may also be formulated as suppositories, e.g.,  
15 containing conventional suppository bases such as cocoa butter or other  
glycerides.


10 Additionally, formulations of the present invention may be formulated  
20 for parenteral administration by injection or continuous infusion. Formulations  
for injection may take such forms as suspensions, solutions, or emulsions in  
oily or aqueous vehicles, and may contain formulatory agents such as  
suspending, stabilizing and/or dispersing agents. Alternatively, the active  
25 ingredient may be in powder form for constitution with a suitable vehicle (e.g.,  
15 sterile, pyrogen-free water) before use.

The formulations according to the invention may also be formulated as  
30 a depot preparation. Such long acting formulations may be administered by  
implantation (for example, subcutaneously or intramuscularly) or by  
20 intramuscular injection. Accordingly, the compounds of the invention may be  
formulated with suitable polymeric or hydrophobic materials (as an emulsion  
35 in an acceptable oil, for example), ion exchange resins or as sparingly soluble  
derivatives as a sparingly soluble salt, for example.

The formulations according to the invention may contain between 0.1 -  
40 25 99% of the active ingredient, conveniently from 30 - 95% for tablets and  
capsules and 3 - 50% for liquid preparations.


There is further provided by the present invention processes for the  
preparation of compounds of the invention Unless otherwise indicated, R<sup>1</sup>, R<sup>1</sup>,  
45 R<sup>2</sup>, R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup>, R<sup>6</sup>, n, and X are as defined above.

30 A compound of structural formula (I) may be prepared from the  
condensation of a compound of formula (II)



(II)


with a compound of formula (III), for example, in a polar organic solvent, such




(III)

as methanol, at temperatures from  $-20^{\circ}\text{C}$  to  $150^{\circ}\text{C}$ , such as  $65^{\circ}\text{C}$ , in the presence of a dehydrating agent such as trimethylorthoformate and/or molecular sieves.

Alternatively, a compound of formula (I) where  $\text{R}^5$  is hydrogen may be prepared by reaction of a compound of formula (II) with a compound of



(IVa)




(IVb)

formula (IVa) in an organic solvent such as methanol at temperatures from  $-20^{\circ}\text{C}$  to  $150^{\circ}\text{C}$ , such as  $65^{\circ}\text{C}$ , in the presence of a base, such as diisopropylethylamine.

Compounds of Formula (I) where  $\text{R}^2$  is  $\text{CF}_3$  may be prepared from the reaction of compounds of formula (II) with compounds of formula (IVb) in an organic solvent such as methanol at temperatures from  $-20^{\circ}\text{C}$  to  $150^{\circ}\text{C}$ , such as  $23^{\circ}\text{C}$ , in the presence of a base such as diisopropylamine.

Compounds of formula (II) may be prepared via alkylation of compounds of formula (V), preferably where  $\text{R}^1$  is hydrogen, with compounds of formula (VI), where  $\text{R}^*$  is an activating group such as mesylate followed by



deprotection of the amine under acidic conditions such as 4.0 N HCl in dioxane solution at 20 °C, or trifluoroacetic acid in dichloromethane at 20 °C.


Alternatively, compounds of formula (II) may be prepared from reaction of compounds of formula (V), preferably where R<sup>1</sup> is methyl, with compounds of formula (VI), preferably where R\* is hydrogen, in an organic solvent such as toluene at temperatures from -20 °C to 150 °C such as 20 °C in the presence of triphenylphosphine and diethylazodicarboxylate in a Mitsunobu type procedure. Standard methyl ester hydrolysis conditions such as lithium hydroxide in tetrahydrofuran and water at 20 °C followed by amine deprotection as hereinbefore described (see also Greene, T. W. and Wutz, P. G. M. "Protective Groups in Organic Synthesis" 2<sup>nd</sup> edition, 1991, John Wiley and Sons for a general discussion on protective group installation/removal) provides compounds of structural formula (II).

Compounds of formula (V) are commercially available and may also be prepared by someone skilled in the art from commercially available compounds.

Compounds of formula (VI), for example where R\* is mesylate, may be prepared by reaction of an alcohol of formula (VI) where R\* is hydrogen, with methanesulfonyl chloride in an organic solvent such as tetrahydrofuran at temperatures from -20 °C to 150 °C such as 0 °C in the presence of an amine base such as triethylamine.



Alcohols of formula (VI) where R\* is hydrogen, may be prepared via reduction of the corresponding carboxylic acid or ester of formula (IX) in an organic solvent such as diethyl ether at temperatures of from -20 °C to 100 °C, such as 0 °C, in the presence of a metal hydride reagent such as lithium aluminum hydride. These acids or esters of formula (IX) may be prepared by a two stage alkylation/cyclization process between an amide or thioamide

compound of formula (VII), and a bromoketoester compound of formula (VIII), where  $R^m$  is alkyl. The alkylation/cyclization reaction may be performed in an organic solvent such as toluene or ethanol at temperatures of from  $-20\text{ }^{\circ}\text{C}$  to  $200\text{ }^{\circ}\text{C}$ , such as  $110\text{ }^{\circ}\text{C}$ , in the presence of a base such as triethylamine. For the preparation of (VIII), see: Chem. Pharm. Bull. (1986), 34(7), 2840-51, J. Med. Chem. (1992), 35(14), 2617-26, patent WO 9731907.




Amides or thioamides of formula (VII) are commercially available or may be prepared from readily available compounds by one skilled in the art.

Compounds of formula (III) are commercially available or may be readily prepared by one skilled in the art, for example, by reacting a ketone of formula (X) with an ester of formula (XII), or by reacting a ketone of formula (XIII) with an ester of formula (XI), by reaction in an organic solvent, such as tetrahydrofuran, at temperatures from  $-20\text{ }^{\circ}\text{C}$  to  $150\text{ }^{\circ}\text{C}$ , such as  $20\text{ }^{\circ}\text{C}$ , in the presence of a base, such as sodium hydride, a metal cation scavenger, such as dibenzo-18-crown-6, and catalytic quantities of an alcohol, such as ethanol (See: Popic, V. V. et al. *Synthesis* 1991, 195-197).



Compounds of formula (IVa) may be prepared from the addition of compounds of formula (XV) to compounds of formula (XIV) in an organic solvent such as tetrahydrofuran at temperatures of from  $-100^{\circ}\text{C}$  to  $100^{\circ}\text{C}$  such as  $-78^{\circ}\text{C}$  followed by oxidation of the intermediate propargylic alcohol with an oxidant, such as manganese dioxide, in an organic solvent, such as dichloromethane, at temperatures from  $-20^{\circ}\text{C}$  to  $100^{\circ}\text{C}$ , such as  $20^{\circ}\text{C}$ .



Alternatively, certain yne-one compounds of formula (IVa) may be prepared by dehydration of a compound of formula (III) where  $\text{R}^5$  is hydrogen in an organic solvent, such as dichloromethane, at temperatures from  $-20^{\circ}\text{C}$  to  $100^{\circ}\text{C}$ , such as  $20^{\circ}\text{C}$ , in the presence of triphenylphosphine, bromine, and a base, such as triethylamine. When  $\text{R}^2$  is  $\text{CF}_3$ ,  $\text{R}^5$  is hydrogen, and  $\text{R}^3$  is phenyl, a compound of formula (III) in an organic solvent, such as dichloromethane, at temperatures from  $-20^{\circ}\text{C}$  to  $100^{\circ}\text{C}$ , such as  $20^{\circ}\text{C}$ , in the presence of triphenylphosphine, bromine, and a base, such as triethylamine yields a compound of formula (IVb). Compounds of formula (X), (XI), (XII), (XIII), (XIV), and (XV) are commercially available or may be prepared from readily available materials from one skilled in the art.

Suitable reaction conditions are described below and in the accompanying Examples. See also, for example, Chung et al., *Selective Functionalization of (S)-Tyrosine*, Tetrahedron, 49(26), pp. 5767-5776, (1993), Solar et al., *Selective O-Alkylation of Tyrosine*, Journal of Organic Chemistry, 31, pp 1996-1997 (1966), O. Mitsunobu, *Synthesis*, p 1 (1981), and D.L. Hughes, *Org. React.* Vol. 42, p 335 (1992).

A compound of formula (I) can be converted to another compound of formula (I). A particular interconversion reaction involves conversion of a compound of formula (I) wherein  $\text{R}^1$  represents  $\text{C}_{1-3}$ alkyl, to a compound of formula (I) wherein  $\text{R}^1$  represents hydrogen, suitably employing hydrolytic techniques e.g. an alkali metal hydroxide, in the presence of an ether solvent



5 e.g. tetrahydrofuran and an alcoholic solvent e.g. methanol or the like. It will  
therefore be appreciated by persons skilled in the art that compounds which  
fall within general formula (I), may in some instances, be hereinafter  
10 described in the intermediate section, as they are useful for the preparation of  
5 other compounds of formula (I).

For any of the general processes and schemes described above, it  
may be necessary and/or desirable to protect sensitive or reactive groups.  
15 Protecting groups are employed according to standard methods of organic  
synthesis (T. W. Green and P. G. M. Wuts (1991) Protecting Groups in  
10 Organic Synthesis, John Wiley & Sons). These groups are removed at a  
convenient stage of synthesis using methods known from the art. Thus, for  
20 example, amino groups may be protected by a group selected from aralkyl  
(e.g. benzyl), acyl, or sulfonyl, e.g. allylsulfonyl, tert-butoxycarbonyl,  
phthalimide, or tosyl; subsequent removal of the protecting group being  
25 effected when desired by hydrolysis or hydrogenolysis as appropriate using  
standard conditions. Thus, for example, tert-butoxycarbonyl groups may be  
removed by hydrolysis under acidic conditions. Hydroxyl and carboxyl groups  
30 may be protected using any conventional hydroxyl or carboxyl protecting  
group. Examples of suitable hydroxyl and carboxyl protecting groups include  
20 groups selected from alkyl, e.g. methyl, tert-butyl, or methoxymethyl, aralkyl,  
e.g. benzyl, diphenylmethyl, or triphenylmethyl, heterocyclic groups such as  
35 tetrahydropyranyl, acyl, e.g. acetyl or benzoyl, and silyl groups such as  
trialkylsilyl, e.g. tert-butyldimethylsilyl. The hydroxyl protecting groups may be  
removed by conventional techniques. Thus, for example, alkyl, silyl, acyl, and  
40 25 heterocyclic groups may be removed by hydrolysis under acidic or basic  
conditions. Aralkyl groups such as triphenylmethyl may similarly be removed  
by hydrolysis under acidic conditions. Aralkyl groups such as benzyl may be  
cleaved by hydrogenolysis in the presence of a Noble metal catalyst such as  
45 palladium-on-charcoal. Silyl groups may also conveniently be removed using  
30 a source of fluoride ions such as tetra-n-butylammonium fluoride.

The following examples are set forth to illustrate the synthesis of some  
50 particular compounds of the present invention and to further exemplify  
particular applications of general processes described above. Accordingly,

the following Example section is in no way intended to limit the scope of the invention contemplated herein.

### Examples

As used herein the symbols and conventions used in these processes, schemes and examples are consistent with those used in the contemporary scientific literature, for example, the Journal of the American Chemical Society. Unless otherwise noted, all starting materials were obtained from commercial suppliers and used without further purification. Specifically, the following abbreviations may be used in the examples and throughout the specification: g (grams); mg (milligrams); L (liters); mL (milliliters);  $\mu$ L (microliters); N (normal); mM (millimolar); mmol (millimoles); i. v. (intravenous); Hz (Hertz); MHz (megahertz); mol (moles); RT or rt (room temperature); min (minutes); h (hours); mp. (melting point); TLC (thin layer chromatography); HPLC (high pressure liquid chromatography); ms (mass spectrum); ES+ (electrospray);  $R_f$  (retention fraction);  $t_r$  (retention time); RP (reverse phase); MeOH (methanol); TFA (trifluoroacetic acid); HCl (hydrochloric acid); HCO<sub>2</sub>H (formic acid); THF (tetrahydrofuran); CH<sub>3</sub>CN (acetonitrile); EtOH (ethanol); CDCl<sub>3</sub> (deuterated chloroform); DMSO (dimethylsulfoxide); DMSO-d<sub>6</sub> (dimethylsulfoxide-deuterated); EtOAc (ethyl acetate); DCM or CH<sub>2</sub>Cl<sub>2</sub> (dichloromethane); DMF (dimethylformamide); Et<sub>3</sub>N (triethylamine); MgSO<sub>4</sub> (magnesium sulfate); H<sub>2</sub>O (water); LAH (lithium aluminum hydride); NaH (sodium hydride); Na<sub>2</sub>CO<sub>3</sub> (sodium carbonate); Na<sub>2</sub>SO<sub>4</sub> (sodium sulfate); MnO<sub>2</sub> (manganese dioxide); KCN (potassium cyanide); BH<sub>3</sub>·THF (borane.tetrahydrofuran complex); NaOMe (sodium methoxide); IPA (isopropanol); Pd/C (palladium on carbon); NaOH (sodium hydroxide); LiOH (lithium hydroxide); DIEA (diisopropylethylamine); Et<sub>2</sub>O (diethyl ether; diethyl azodicarboxylate (DEAD); Diisopropyl azodicarboxylate (DIAD); tert-butyloxycarbonyl (BOC); NaHCO<sub>3</sub> (saturated aqueous sodium bicarbonate). Brine refers to a saturated aqueous solution of NaCl. Unless otherwise indicated, all temperatures are expressed in °C (degrees

Centigrade). All reactions conducted at room temperature unless otherwise noted.

The  $^1\text{H}$  NMR spectra were recorded on a Varian VXR-300, a Varian Unity-300, or a Varian Unity-400 instrument. Chemical shifts are expressed in parts per million (ppm,  $\delta$  units). Coupling constants are in units of hertz (Hz). Splitting patterns are designated as s, singlet; d, doublet; t, triplet; q, quartet; m, multiplet; br, broad; hept, heptuplet.

Low-resolution mass spectra (MS) were recorded on a JOEL JMS-AX505HA, JOEL SX-102, Micromass Platform 2 LC/Ms, or a SCIEX-APIiii spectrometers. All mass spectra were taken under electrospray ionization (ES, either in the positive ion mode or negative ion mode) or by fast atom bombardment (FAB) methods. Infrared (IR) spectra were obtained on a Nicolet 510 FT-IR spectrometer using a 1-mm NaCl cell. All reactions were monitored by thin-layer chromatography on 0.25 mm E. Merck silica gel plates (60F-254), visualized with UV light, iodine staining, or 7% ethanolic phosphomolybdic acid or p-anisaldehyde solutions. Flash column chromatography was performed on silica gel (230-400 mesh, Merck).

Analytical purity was assessed on a Hewlett Packard series 1050 or 1100 system equipped with a diode array spectrometer. The stationary phase was either a Dynamax C8 column (25 cm x 4.1 mm), a Dynamax 60A C18 column (25 cm x 4.6 mm), a Vydac C18 column (5m, 4.6 mm X 250 mm), a Supelco C18 column (5m, 4.6 mm X 150 mm), or a Rainin C18 column (5m, 4.6 mm X 250 mm). The flow rate was 1.0 to 1.5 ml/min. ( $t_0$  = 2.8 or 3.0 min.) and the solvent systems were as described below. Enantiomeric purity was assessed using either a Chiralpak AD column (25 cm x 4.6 mm) or a Chiralpak OD column (25cm x 4.6 mm) on either a Hewlett Packard series 1050 HPLC system equipped with a diode array spectrometer or on a Supercritical Fluid (SFC) system using  $\text{CO}_2$  / methanol as the mobile phase.

**Intermediate 1: 2-[5-methyl-2-phenyl-1,3-oxazol-4-yl]acetic acid methyl ester**

5 To a solution of 140 g (0.67 mol) of methyl 4-bromo-3-oxo-pentanoate  
in 300 mL of anhydrous toluene was added 210 g (1.73 mol) of benzamide,  
132 g (0.93 mol) of sodium hydrogen phosphate, and ~ 20 mL of EtOH. The  
10 suspension was stirred with heating to 95 °C for 24 h, then cooled to 0 °C,  
5 filtered, and the solids washed with cyclohexane (2x100 mL) followed by 20 %  
EtOAc in hexanes (200 mL). The combined filtrates were washed with 2x100  
mL 10% KOH solution and brine (100 mL). The organics were dried (MgSO<sub>4</sub>),  
15 concentrated to a thin oil, and purified by chromatography on 1 kg of silica gel  
(230-400 mesh). Gradient elution of the column with 10% to 50% EtOAc in  
10 Hexanes gave 54 grams (35% yield) of the title compound as a pale yellow  
oil; <sup>1</sup>H NMR (CDCl<sub>3</sub>, 400 MHz) δ 8.0-7.98 (m, 2H), 7.43-7.40 (m, 3H), 3.73 (s,  
20 3H), 3.59 (s, 2H), 2.37 (s, 3H); low resolution MS (ES+) *m* / *e* 232.4 (MH<sup>+</sup>);  
TLC R<sub>f</sub> = 0.64 (4/1 hexanes/EtOAc).

**Intermediate 1B: 2-[5-ethyl-2-phenyl-1,3-oxazol-4-yl]acetic acid ethyl ester**

The title compound was prepared (as described above for the preparation of example 1) from ethyl 4-bromo-3-oxo-hexanoate (3.32 g, 14 mmol) and 6.79 g (56 mmol) of benzamide to give 710 mg of Intermediate 1B: TLC  $R_f$  = 0.69 (2/1 hexanes/EtOAc);  $^1\text{H}$  NMR ( $\text{CDCl}_3$ , 300 MHz)  $\delta$  7.98 (m, 2H), 7.41 (m, 3H), 4.18 (q, 2H,  $J$  = 6.9), 3.57 (s, 2H), 2.72 (q, 2H,  $J$  = 7.5), 1.29-1.24 (m, 6H); low resolution MS ( $\text{ES}^+$ )  $m/e$  260.1 ( $\text{MH}^+$ ).

**Intermediate 1C: 2-[2-phenyl-5-propyl-1,3-oxazol-4-yl]acetic acid ethyl ester**

The title compound was prepared (as described above for the preparation of example 1) from ethyl 4-bromo-3-oxo-heptanoate (2.83 g, 11.4 mmol) and 5.51 g (45 mmol) of benzamide to give 820 mg of Intermediate 1C: TLC  $R_f$  = 0.70 (2/1 hexanes/EtOAc);  $^1\text{H}$  NMR ( $\text{CDCl}_3$ , 300 MHz)  $\delta$  7.98 (m, 2H), 7.43-7.4 (m, 3H), 4.18 (q, 2H,  $J$  = 7.2), 3.56 (s, 2H), 2.66 (t, 2H,  $J$  = 7.5), 1.76-1.68 (m, 2H), 1.27 (t, 3H,  $J$  = 7.2), 0.99 (t, 3H,  $J$  = 7.2); low resolution MS ( $\text{ES}^+$ )  $m/e$  274.1 ( $\text{MH}^+$ ).

**Intermediate 1D: 5-methyl-2-phenyl-1,3-oxazole-4-carboxylic acid**  
Commercial sources.

**Intermediate 1E: 5-ethyl-2-phenyl-1,3-oxazole-4-carboxylic acid**

Intermediate 1E was prepared (as described below for the preparation of intermediate 2B) from 4.08 g of benzamide, 5.0 g of ethyl 3-bromo-2-oxo-pentanoate, and 10 mL of toluene. Solids were collected and washed with cold water after cooling of the NaOH solution. Heating these solids with citric acid as described followed by cooling and isolation of the resulting solids gave 410 mg (9% yield) of the title compound;  $^1\text{H}$  NMR ( $\text{CDCl}_3$ , 300 MHz)  $\delta$  8.06-8.03 (m, 2H), 7.49-7.43 (m, 3H), 3.14 (q, 2H,  $J$  = 8.0), 1.34 (t, 3H,  $J$  = 8.0); low resolution MS ( $\text{ES}^+$ )  $m/e$  217.8 ( $\text{MH}^+$ ).

**Intermediate 2: 2-[5-methyl-2-phenyl-1,3-thiazol-4-yl]acetic acid methyl ester**

A solution of 13.13g (95.7 mmol) of thiobenzamide and 5.0 g (23.9 mmol) of methyl 4-bromo-3-oxo-pentanoate in 25 mL of dry toluene was heated at 90 °C for 5 h. The solution was poured into EtOAc/water and the aqueous phase extracted with EtOAc. The organics were washed with brine, dried (MgSO<sub>4</sub>), filtered, and concentrated. The crude oil was purified by silica gel chromatography eluting with hexanes/EtOAc (3/1 to 2/1) to afford 4.14 g (70%) yield) of the title compound as a pale orange-colored oil: <sup>1</sup>H NMR (CDCl<sub>3</sub>, 400 MHz) δ 7.97 (m, 2H), 7.11 (m, 2H), 3.73 (s, 3H), 3.56 (s, 2H), 2.36 (s, 3H).

**Intermediate 2B: 2-[5-ethyl-2-phenyl-1,3-thiazol-4-yl]acetic acid**

A suspension of 8.7 g (63.3 mmol) of thiobenzamide and 10 g (42.2 mmol) of ethyl 4-bromo-3-oxo-hexanoate in 45 mL of toluene was refluxed for 6 hrs with Dean-Stark trap removal of water. Cooled, added methanol, and continued reflux for 1 h. Cooled, partially concentrated (~1/2 volume), then 30 mL of 2.0 M NaOH solution was added. Stirred the mixture at 80 °C for 2 hrs before cooling and extraction of the aqueous with EtOAc. The aqueous phase was treated with 70 mL of 1.0 M citric acid with heating (70 °C) for 45 min. The mixture was extracted with EtOAc, and the extracts were dried over MgSO<sub>4</sub>, filtered and concentrated to yield a yellow colored solid upon standing at ambient temperature (8.4 g, ~80 % yield); <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) δ 7.95-7.87 (m, 2H), 7.48-7.45 (m, 3H), 3.86 (s, 2H), 2.85 (q, 2H, J = 7.5), 1.36 (t, 3H, J = 7.5).

**Intermediate 2C: 5-ethyl-2-phenyl-1,3-thiazole-4-carboxylic acid**

Intermediate 2C was prepared as described above for the preparation of intermediate 2B from 4.61 g of thiobenzamide, 5.0 g of ethyl 3-bromo-2-oxo-pentanoate, and 14 mL of toluene. Solids were collected and washed with cold water after cooling of the NaOH solution. Heating these solids with citric acid as described followed by cooling and isolation of the resulting solids gave

3.74 g (72% yield) of intermediate 2C;  $^1\text{H}$  NMR ( $\text{CDCl}_3$ , 300 MHz)  $\delta$  7.913-7.91(m, 2H), 7.51-7.50 (m, 3H), 3.39 (q, 2H,  $J=7.8$ ), 1.42 (t, 3H,  $J=7.8$ ); low resolution MS ( $\text{ES}^+$ )  $m/e$  233.81 ( $\text{MH}^+$ ).

**Intermediate 3: 2-[2-(4-fluorophenyl)-5-methyl-1,3-oxazol-4-yl]acetic acid methyl ester**

A solution of 667 mg (4.80 mmol) of 4-fluorobenzamide and 1.0 g (4.80 mmol) of methyl 4-bromo-3-oxo-pentanoate in 6 mL of dry toluene was heated at 120 °C for 16 h. The resulting dark slurry was cooled to rt, diluted with 10 mL of EtOAc, and washed with  $\text{NaHCO}_3$  (1 x 10 mL). The organic layer was separated, dried ( $\text{MgSO}_4$ ), and the solvents removed under reduced pressure. Purification of the material by silica gel flash column chromatography using hexane/EtOAc 4 / 1 as eluent to afford 308 mg of the title compound as a clear oil:  $^1\text{H}$  NMR ( $\text{CDCl}_3$ , 400 MHz)  $\delta$  7.97 (m, 2H), 7.11 (m, 2H), 3.73 (s, 3H), 3.56 (s, 2H), 2.36 (s, 3H).

**Intermediate 3B: 2-[5-ethyl-2-(4-fluorophenyl)-1,3-oxazol-4-yl]acetic acid**

The title compound was prepared from 5 g of 4-fluorobenzamide and 5.67 g of ethyl 4-bromo-3-oxo-hexanoate as described in example 2B to give 1.8 grams of intermediate 3B as a solid:  $^1\text{H}$  NMR ( $\text{CDCl}_3$ , 300 MHz)  $\delta$  7.96 (dd, 2H,  $J=8.8, 5.3$ ), 7.11 (t, 2H,  $J=8.6$ ), 3.61 (s, 2H), 2.69 (q, 2H,  $J=7.5$ ), 1.27 (t, 3H,  $J=7.5$ ).

**Intermediate 3C: 5-ethyl-2-(4-fluorophenyl)-1,3-oxazole-4-carboxylic acid**

The title compound was prepared from 6.5 g of 4-fluorobenzamide and 6.08 g of ethyl 3-bromo-2-oxo-pentanoate (as described in example 2B) to give 1.55 g (22%) of intermediate 3C as a solid:  $^1\text{H}$  NMR ( $\text{CDCl}_3$ , 300 MHz)  $\delta$  8.12-8.08 (m, 2H), 7.23-7.17 (m, 2H), 3.19 (q, 2H,  $J=7.5$ ), 1.40 (t, 3H,  $J=7.5$ ).

**Intermediate 4: 2-[2-(4-fluorophenyl)-5-methyl-1,3-thiazol-4-yl]acetic acid ethyl ester**

5 A mixture of 8.75 g (56.4 mmol) of 4-fluorobenzamide and 11.79 g  
(56.4 mmol) of methyl 4-bromo-3-oxo-pentanoate in 45 mL of EtOH was  
10 refluxed for 14 h. The solids formed upon cooling were removed via filtration  
and the solvent removed under reduced pressure. The residue was taken into  
5 EtOAc and washed with water, brine, and dried over MgSO<sub>4</sub>. <sup>1</sup>H NMR of the  
crude Intermediate 4 indicated purity sufficient to proceed into the next stage  
of synthesis. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 400 MHz) δ 7.97 (m, 2H), 7.11 (m, 2H), 3.73 (s,  
15 3H), 3.56 (s, 2H), 2.36 (s, 3H).

10 **Intermediate 4B: 2-[5-ethyl-2-(4-fluorophenyl)-1,3-thiazol-4-yl]acetic acid**

20 The title compound was prepared from 6.8 g of 4-fluorothiobenzamide and  
7.42 g of ethyl 4-bromo-3-oxo-hexanoate as described in example 2B to give  
2.9 grams of intermediate 4B as a solid: <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) δ 7.96  
(dd, 2H, J = 8.8, 5.3), 7.11 (t, 2H, J = 8.6), 3.61 (s, 2H), 2.69 (q, 2H, J = 7.5),  
25 1.27 (t, 3H, J = 7.5).

30 **Intermediate 5: 2-[2-(4-methoxyphenyl)-5-methyl-1,3-oxazol-4-yl]acetic acid methyl ester**

20 A mixture of 725 mg (4.80 mmol) of 4-methoxybenzamide and 1.0 g  
(4.80 mmol) of methyl 4-bromo-3-oxo-pentanoate was heated neat at 120 °C  
35 for 2 h. The resulting dark slurry was cooled to RT, diluted with 2 mL of DCM  
and purified by silica gel flash column chromatography using hexane/EtOAc  
(3/1) as eluent to afford 189 mg of the title compound as a yellow solid: <sup>1</sup>H  
40 25 NMR (CDCl<sub>3</sub>, 400 MHz) δ 7.92 (d, 2H, J = 8.9), 6.93 (d, 2H, J = 8.9), 3.85 (s,  
3H), 3.73 (s, 3H), 3.56 (s, 2H), 2.34 (s, 3H); low resolution MS (FAB) m/e  
285 (MH<sup>+</sup>), 284 (M<sup>+</sup>).

45 **Intermediate 6: 2-[2-(4-trifluoromethylphenyl)-5-methyl-1,3-oxazol-4-yl]acetic acid methyl ester**

30 A mixture of 3.62 g (19.1 mmol) of 4-trifluoromethylbenzamide and 4.0  
50 g (19 mmol) of methyl 4-bromo-3-oxo-pentanoate in 15 mL of toluene and 10



5 mL of dioxane (with 2 mL of EtOH) was refluxed for 22 h. The solution was concentrated under reduced pressure. The residue was taken into EtOAc and washed with water, brine, and dried over MgSO<sub>4</sub>. The crude material was purified by silica gel chromatography (10% - 40% EtOAc in hexanes) to yield 130 mg of a yellow solid: <sup>1</sup>H NMR (CDCl<sub>3</sub>, 400 MHz) δ 8.1 (d, 2H, J = 8.3), 7.65 (d, 2H, J = 8.4), 3.75 (s, 3H), 3.58 (s, 2H), 2.39 (s, 3H); low resolution MS (ES<sup>+</sup>) *m/e* 300.0 (MH<sup>+</sup>).

**Intermediate 7: 2-[2-(4-isopropoxyphenyl)-5-methyl-1,3-oxazol-4-yl]acetic acid ethyl ester**

A mixture of 2.8 g (15.6 mmol) of 4-isopropoxybenzamide and 3.26 g (15.6 mmol) of methyl 4-bromo-3-oxo-pentanoate in 10 mL of toluene and 1 mL of EtOH was refluxed for 24 h. The solvents were removed and the crude residue purified directly via silica gel chromatography (4/1 hexanes/EtOAc) to yield 1.31 g (28%) of Intermediate 6: <sup>1</sup>H NMR (CDCl<sub>3</sub>, 400 MHz) δ 7.92 (d, 2H, J = 8.9), 6.93 (d, 2H, J = 8.9), 3.85 (s, 3H), 3.73 (s, 3H), 3.56 (s, 2H), 2.34 (s, 3H); low resolution MS (FAB) *m/e* 285 (MH<sup>+</sup>).

**Intermediate 8: 2-[5-methyl-phenyl-1,3-oxazol-4-yl]ethanol**

To a solution of 330 mg (1.43 mmol) of intermediate 1 in 5 mL of THF at 0 °C was added dropwise 1.43 mL (1.43 mmol) of 1.0 M LAH in Et<sub>2</sub>O solution. The solution was stirred at 0 °C for 1 hr, quenched by the addition of 0.054 mL H<sub>2</sub>O, 0.054 mL of 1.0 N NaOH solution, and 0.162 mL of H<sub>2</sub>O, and MgSO<sub>4</sub> was added. The mixture was filtered, the filtrate was concentrated, and the crude oil dried under vacuum for several hours to give 280 mg of Intermediate 8 as a pale yellow-colored solid: <sup>1</sup>H NMR (CDCl<sub>3</sub>, 400 MHz) δ 7.97 (m, 2H), 7.42 (m, 3H), 3.92 (t, 2H, J = 5.6), 2.72 (t, 2H, J = 5.6), 2.33 (s, 3H); low resolution MS (ES<sup>+</sup>) *m/e* 204 (MH<sup>+</sup>); TLC R<sub>f</sub> = 0.21 (1/1 hexanes/EtOAc)

**Intermediate 8B: 2-[5-ethyl-2-phenyl-1,3-oxazol-4-yl]ethanol** The title compound was prepared (as described above for the preparation of

Intermediate 8 from intermediate 1B (710 mg, 2.74 mmol) and 125 mg LAH to give 480 mg (81% yield) of intermediate 8B: TLC  $R_f$  = 0.21 (2/1 hexanes/EtOAc);  $^1\text{H}$  NMR ( $\text{CDCl}_3$ , 400 MHz)  $\delta$  8.0-7.97 (m, 2H), 7.46-7.41 (m, 3H), 3.92 (t, 2H,  $J$  = 5.6), 3.33 (broad s, 1H), 2.75-2.66 (m, 4H), 1.27 (t, 3H,  $J$  = 7.6); low resolution MS ( $\text{ES}^+$ )  $m/e$  218.15 ( $\text{MH}^+$ ).

**Intermediate 8C: 2-[2-phenyl-5-propyl-1,3-oxazol-4-yl]ethanol** The title compound was prepared (as described above for the preparation of Intermediate 8) from intermediate 1C (820 mg, 3.0 mmol) and 137 mg LAH to give 360 mg (52% yield) of intermediate 8C: TLC  $R_f$  = 0.29 (2/1 hexanes/EtOAc);

$^1\text{H}$  NMR ( $\text{CDCl}_3$ , 400 MHz)  $\delta$  7.91(m, 2H), 7.39-7.32 (m, 3H), 3.86 (t, 2H,  $J$  = 5.6), 2.68-2.55 (m, 2H), 4.48-2.41 (m, 2H), 1.68-1.53 (m, 2H), 0.91 (t, 3H,  $J$  = 7.6); low resolution MS ( $\text{ES}^+$ )  $m/e$  232.14 ( $\text{MH}^+$ ).

**Intermediate 8D: [5-methyl-2-phenyl-1,3-oxazol-4-yl]methanol** The title compound was prepared via LAH reduction of intermediate 1D. Thus, 670 mg of Intermediate 1D (3.30 mmol) was stirred with 165 mg of LAH at 23 °C for two hours. Silica gel chromatography gave 120 mg (19% yield) of Intermediate 8D: TLC  $R_f$  = 0.28 (2/1 hexanes/EtOAc);  $^1\text{H}$  NMR ( $\text{CDCl}_3$ , 300 MHz)  $\delta$  8.0-7.97 (m, 2H), 7.49-7.41 (m, 3H), 4.59 (s, 2H), 4.18-4.05 (broad s, 1H), 2.38 (s, 3H); low resolution MS ( $\text{ES}^+$ )  $m/e$  190.1 ( $\text{MH}^+$ ).

**Intermediate 8E: 3-[5-methyl-2-phenyl-1,3-oxazol-4-yl]propanol** The title compound was prepared from intermediate 8 via a 1-carbon homologation sequence as follows: The mesylate (3.88 g, 1.38 mmol) of alcohol Intermediate 8 (from  $\text{Et}_3\text{N}$ , methanesulfonyl chloride, THF, 0 °C) was heated to 120 °C with 1.08 g (16.6 mmol) of KCN for 14 hr. Aqueous work-up and silica gel chromatography (EtOAc/hexanes) gave 860 mg (30%) of cyanide intermediate. Hydrolysis to the acid intermediate was accomplished in refluxing EtOH/water for 2.5 hrs with NaOH (650 mg). Acidification of the aqueous phase gave 580 mg (62% yield) of the acid. Reduction with LAH (95

mg, 3.0 mmol) in Et<sub>2</sub>O for 3 hrs followed by silica gel chromatography provided 370 mg (68% yield) of Intermediate 8E: TLC R<sub>f</sub> = 0.14 (2/1 hexanes/EtOAc); <sup>1</sup>H NMR (CDCl<sub>3</sub>, 400 MHz) δ 7.98-7.93 (m, 2H), 7.45-7.37 (m, 3H), 3.75 (t, 2H, J = 8.7), 3.36 (br s, 1H), 2.64 (t, 2H, J = 6.9), 2.33 (s, 3H), 1.94-1.85 (m, 2H); low resolution MS (ES<sup>+</sup>) m/e 218.17 (MH<sup>+</sup>).

**Intermediate 8F: 3-[5-ethyl-2-phenyl-1,3-oxazol-4-yl]propanol**

Intermediate 8F was prepared from intermediate 8B via a one carbon homologation procedure as described for the preparation of intermediate 8E. Thus, 495 mg of alcohol intermediate 8B was converted to intermediate 8F via mesylate (380 mg), cyanide (355 mg), and acid (280 mg) intermediates. Borane reduction (THF, 0 °C, 6.5 mL of 1.0 M BH<sub>3</sub>/THF complex) on 273 mg of acid intermediate gave 215 mg of alcohol intermediate 8F; <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) δ 8.05-8.01 (m, 2H), 7.48-7.44 (m, 3H), 3.79 (t, 2H, J = 5.8), 2.78-2.67 (m, 4H), 1.97-1.91 (m, 2H), 1.31 (t, 3H, J = 7.5); low resolution MS (ES<sup>+</sup>) m/e 231.91 (MH<sup>+</sup>); TLC R<sub>f</sub> = 0.49 (1/2 hexanes/EtOAc).

**Intermediate 8G: [5-ethyl-2-phenyl-1,3-oxazol-4-yl]methanol**

Intermediate 8G was prepared (as described below for intermediate 9B) from 410 mg of intermediate 1E to give 335 mg (87%) of the title compound; <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) δ 7.96-7.940 (m, 2H), 7.39-7.37 (m, 3H), 4.54 (s, 2H), 2.71 (q, 2H, J = 7.5), 2.38 (br s, 1H), 1.25 (t, 3H, J = 7.5); low resolution MS (ES<sup>+</sup>) m/e 203.94 (MH<sup>+</sup>).

**Intermediate 9: 2-[5-methyl-2-phenyl-1,3-thiazol-4-yl]ethanol**

To a 0 °C stirred solution of 6.11 g (24.7 mmol) of ester Intermediate 2 in 100 mL of THF was added 935 mg (25 mmol) of LAH. The reaction was stirred for 20 minutes at 20 °C, cooled to 0 °C, and quenched with water. The solids were filtered, washed with Et<sub>2</sub>O, and the filtrate was dried (MgSO<sub>4</sub>). Silica gel chromatography with EtOAc in hexanes (40% to 50%) gave 2.33 g (38% yield) of Intermediate 9: <sup>1</sup>H NMR (CDCl<sub>3</sub>, 400 MHz) δ 7.85 (m, 2H), 7.42

(m, 3H), 3.99 (m, 3), 2.90 (m, 2H), 2.41 (s, 3H); low resolution MS (ES<sup>+</sup>) *m/e* 220.1 (MH<sup>+</sup>); TLC *R<sub>f</sub>* = 0.43 (1/1 hexanes/EtOAc).

**Intermediate 9B: 2-[5-ethyl-2-phenyl-1,3-thiazol-4-yl]ethanol**

To a THF (15 mL) solution of 4.0 g (16.2 mmol) of intermediate 2B at 0 °C was added dropwise over 10 minutes 57 mL of 1.0 M BH<sub>3</sub>-THF complex in THF. Stirred at 35-40 °C for 3 h before cooling to 0 °C and quenching with 70 mL of MeOH. Concentrated, taken into 50 mL of n-butanol and refluxed for 45 min. The solution was concentrated and the residue collected was purified by silica gel chromatography. Elution with 20-50% EtOAc in hexanes gave 2.4 g (64% yield) of a yellow viscous oil after concentration of the pooled product fractions and drying under vacuum: <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) δ 7.86-7.83 (m, 2H), 7.41-7.34 (m, 3H), 3.96 (t, 2H, *J* = 5.5), 2.89 (t, 2H, *J* = 5.5), 2.77 (q, 2H, *J* = 7.5), 1.27 (t, 3H, *J* = 7.5).

**Intermediate 9C: [5-ethyl-2-phenyl-1,3-thiazol-4-yl]methanol**

Intermediate 9C was prepared (as described above for intermediate 9B) from 3.74 g of intermediate 2C to give 2.45 g (70%) of the title compound; <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) δ 7.93-7.90 (m, 2H), 7.47-7.43 (m, 3H), 4.75 (s, 2H), 3.02 (br s, 1H), 2.89 (q, 2H, *J* = 7.2), 1.35 (t, 3H, *J* = 7.2); low resolution MS (ES<sup>+</sup>) *m/e* 219.88 (MH<sup>+</sup>).

**Intermediate 10: 2-[2-(4-fluorophenyl)-5-methyl-1,3-oxazol-4-yl]ethanol**

To a 0 °C stirred solution of 380 mg (1.44 mmol) of ester Intermediate 3 in 10 mL of THF was added 55 mg (1.44 mmol) of LAH. The reaction was stirred for 20 minutes at 20 °C, cooled to 0 °C, and quenched with saturated aqueous NH<sub>4</sub>Cl solution. The solids were filtered, washed with Et<sub>2</sub>O, and the filtrate was dried (MgSO<sub>4</sub>). Silica gel chromatography with EtOAc/hexanes (1/2 to 2/1) as eluent gave 175 mg (55% yield) of intermediate 10: <sup>1</sup>H NMR (CDCl<sub>3</sub>, 400 MHz) δ 7.96 (m, 2H), 7.12 (m, 2H), 3.92 (d, 2H, *J* = 5.0), 3.21 (s, 1H), 2.71 (t, 2H, *J* = 5.6), 2.32 (s, 3H); low resolution MS (FAB) *m/e* 221 (M<sup>+</sup>); TLC *R<sub>f</sub>* = 0.20 (1/1 hexanes/EtOAc).

**Intermediate 10B: 2-[5-ethyl-2-(4-fluorophenyl)-1,3-oxazol-4-yl]ethanol**

The title compound was prepared as described above for example 9B. From 1.3 g of intermediate 3B was prepared 1.14 g (94%) of intermediate 10B as a yellow-colored, viscous oil; <sup>1</sup>H NMR (CDCl<sub>3</sub>, 400 MHz) δ 7.98-7.94 (m, 2H), 7.10 (t, 2H, J = 8.7), 3.89 (t, 2H, J = 5.6), 2.71 (t, 2H, J = 5.7), 2.66 (t, 2H, J = 7.5), 1.25 (t, 3H, J = 7.5); low resolution MS (ES+) *m/e* 236.17 (MH<sup>+</sup>); TLC R<sub>f</sub> = 0.37 (1/1 hexanes/EtOAc)

**Intermediate 10C: 3-[5-ethyl-2-(4-fluorophenyl)-1,3-oxazol-4-yl]propanol**

Intermediate 10C was prepared from intermediate 10B via a one carbon homologation procedure as described for the preparation of intermediate 8E. Thus, 720 mg of alcohol intermediate 10B was converted to intermediate 10C via mesylate (828 mg), cyanide (495 mg), and acid (440 mg) intermediates. Borane reduction (THF, 0 °C, 6.5 mL of 1.0 M BH<sub>3</sub>/THF complex) instead of LAH reduction on 430 mg of acid intermediate gave 320 mg of alcohol intermediate 10C; <sup>1</sup>H NMR (CDCl<sub>3</sub>, 400 MHz) δ 7.98-7.93 (m, 2H), 7.09 (t, 2H, J = 8.6), 3.72 (t, 2H, J = 5.8), 2.70-2.56 (m, 4H), 1.91-1.84 (m, 2H), 1.24 (t, 3H, J = 7.5); low resolution MS (ES+) *m/e* 250.15 (MH<sup>+</sup>); TLC R<sub>f</sub> = 0.55 (1/2 hexanes/EtOAc).

**Intermediate 10D: [5-ethyl-2-(4-fluorophenyl)-1,3-oxazol-4-yl]methanol**

The title compound was prepared as described above for example 9B. From 1.52 g of intermediate 3C was prepared 735 mg (51%) of intermediate 10D; <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) δ 8.05-8.04 (m, 2H), 7.18-7.16 (m, 2H), 4.64 (s, 2H), 2.82-2.78 (m, 2H), 2.30 (br s, 1H), 1.35-1.31 (m, 3H); low resolution MS (ES+) *m/e* 221.94 (MH<sup>+</sup>); TLC R<sub>f</sub> = 0.60 (1/2 hexanes/EtOAc).

**Intermediate 11: 2-[2-(4-fluorophenyl)-5-methyl-1,3-thiazol-4-yl]ethanol**

The title compound was prepared from crude ester Intermediate 4 and 1.07 g of LAH as described above for the preparation of Intermediate 10 to yield 2.8 g of Intermediate 11 (21% yield over two steps, preparation of Intermediates 4 plus 11); <sup>1</sup>H NMR (CDCl<sub>3</sub>, 400 MHz) δ 7.96 (m, 2H), 7.12 (m,

2H), 3.92 (d, 2H,  $J=5.0$ ), 3.21 (s, 1H), 2.71 (t, 2H,  $J=5.6$ ), 2.32 (s, 3H); low resolution MS (FAB) $m/e$  221 ( $M^+$ ).

**Intermediate 11B: 2-[5-ethyl-2-(4-fluorophenyl)-1,3-thiazol-4-yl]ethanol**

The title compound was prepared as described above for example 9B. From 2.7 g of intermediate 4B was prepared 1.75 g (68%) of intermediate 11B as a colorless, viscous oil;  $^1\text{H}$  NMR ( $\text{CDCl}_3$ , 400 MHz)  $\delta$  7.85-7.80 (m, 2H), 7.07 (t, 2H,  $J=8.7$ ), 3.95 (t, 2H,  $J=5.6$ ), 2.88 (t, 2H,  $J=5.5$ ), 2.77 (q, 2H,  $J=7.5$ ), 1.27 (t, 3H,  $J=7.5$ ); low resolution MS ( $\text{ES}^+$ ) $m/e$  252.17 ( $M^+$ ); TLC  $R_f$  = 0.47 (1/1 hexanes/EtOAc)

**Intermediate 12: 2-[2-(4-methoxyphenyl)-5-methyl-1,3-oxazol-4-yl]ethanol**

To a stirring solution of 185 mg (0.71 mmol) of Intermediate 42 in 5 mL of THF at 0 °C was added 0.71 mL (0.71 mmol, 1.0 equiv) of a 1.0 M solution of  $\text{LiAlH}_4$  in THF. The resulting solution was stirred at RT for 45 min then cooled to 0 °C and quenched by careful addition of 0.027 mL of  $\text{H}_2\text{O}$ , followed by addition of 0.027 mL of 15% NaOH and 0.080 mL of  $\text{H}_2\text{O}$ . The resulting slurry was filtered to remove the solids and the filtrate was concentrated under reduced pressure to afford 164 mg of the title compound as a light yellow oil:  $^1\text{H}$  NMR ( $\text{CDCl}_3$ , 400 MHz)  $\delta$  7.92 (d, 2H,  $J=8.8$ ), 6.94 (d, 2H,  $J=8.8$ ), 3.92 (dt, 2H,  $J=5.7, 11.5$ ), 3.86 (s, 3H), 3.35 (t, 1H,  $J=6.2$ ), 2.71 (t, 2H,  $J=5.7$ ), 2.32 (s, 3H).

**Intermediate 13: 2-[2-(4-trifluoromethylphenyl)-5-methyl-1,3-oxazol-4-yl]ethanol**

The title compound was prepared from 390 mg of ester Intermediate 6 (as described above for the preparation of Intermediate 10) to yield 170 mg of Intermediate 13 : TLC  $R_f$  = 0.24 (1/1 hexanes/EtOAc),  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  8.08 (d, 2H,  $J=8.2$ ), 7.68 (d, 2H,  $J=8.4$ ), 3.93 (q, 2H,  $J=11.4, 5.8$ ), 2.99 (t, 1H,  $J=6.0$ ), 2.74 (t, 2H,  $J=5.8$ ), 1.56 (s, 3H).

**Intermediate 14: 2-[2-(4-isopropoxyphenyl)-5-methyl-1,3-oxazol-4-yl]ethanol**

The title compound was prepared from 1.3 g of ester Intermediate 7 (as described above for the preparation of Intermediate 10) to yield 540 mg of Intermediate 14 : TLC  $R_f$  = 0.21 (1/1 hexanes/EtOAc),  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.87 (d, 2H,  $J$  = 8.8), 6.9 (d, 2H,  $J$  = 9.0), 4.6 (hept, 1H,  $J$  = 6.0), 3.9 (q, 2H,  $J$  = 11.4, 5.7), 3.35 (t, 1H,  $J$  = 6.0), 2.69 (t, 2H,  $J$  = 5.6), 2.3 (s, 3H), 1.35 (d, 6H,  $J$  = 6.0).

**Intermediate 15: 5-[2-(methanesulfonyloxy)ethyl]-4-methyl-2-phenyl-1,3-oxazole**

To a solution of 9.41 g (46.3 mmol) of alcohol Intermediate 8 in  $\text{CH}_2\text{Cl}_2$  (300 mL) at 0 °C was added 3.94 mL (50.9 mmol) of methanesulfonyl chloride followed by 7.75 mL (55.6 mmol) of triethylamine. After 1 hr with gradual warming to 20 °C, the solution was washed with water (3x), brine (1x), and dried over  $\text{MgSO}_4$ . Concentration gave 12.92 g (99%) of Intermediate 15 homogeneous by TLC analysis: ( $R_f$  = 0.24, 1/1 hexanes/EtOAc);  $^1\text{H}$  NMR ( $\text{CDCl}_3$ , 400 MHz)  $\delta$  7.97 (m, 2H), 7.43 (m, 3H), 4.53 (t, 2H,  $J$  = 6.5), 2.95 (s, 3H), 2.95 (t, 2H,  $J$  = 6.5), 2.36 (s, 3H); low resolution MS ( $\text{ES}^+$ )  $m/e$  281.9 ( $\text{MH}^+$ ); TLC  $R_f$  = 0.35 (1/1 hexanes/EtOAc)

**Intermediate 16: 1-(phenyl)-1,3-pentanedione**

The title compound was prepared according to Popio, V.V. et al. *Synthesis* (March 1991), pp 195-197. To a stirred suspension of NaH (1.2 g, 50 mmol) and ethyl propionate (5.73 mL, 50 mmol) in 20 mL THF at 20 °C was added EtOH (2 drops), acetophenone (3.0 g, 25 mmol) in 20 mL of THF, and dibenzo-18-crown-6 (150 mg, 0.4 mmol) in 20 mL of THF. Stirred for 30 min, then at reflux for 1 hr. Cooled (0 °C), then added 25 mL of 10%  $\text{H}_2\text{SO}_4$  solution and the aqueous was extracted with  $\text{Et}_2\text{O}$ . The organics were washed with  $\text{H}_2\text{O}$ , aqueous  $\text{Na}_2\text{CO}_3$ , and brine. The solution was dried over  $\text{Na}_2\text{SO}_4$  and concentrated. Purification by silica gel chromatography (1% to 10% EtOAc in hexanes) gave 4.0 g of Intermediate 16 as a thin oil.  $^1\text{H}$  NMR

indicated an ~10:1 mixture of tautomers favoring the enol form:  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ , enol form)  $\delta$  7.87 (m, 2H), 7.52-7.4 (m, 3H), 6.17 (s, 1H), 2.47 (q, 2H,  $J=7.5$ ), 1.21 (t, 3H,  $J=7.5$ )

Intermediates 17, 18, and 20-33 were prepared analogous to the method described above for the preparation of Intermediate 16.

**Intermediate 17: 1-(phenyl)-1,3-hexanedione**

The title compound was prepared (as described above for Intermediate 16) from 3.0 g of acetophenone and 5.8 g of ethyl butyrate to yield 4.0 grams of Intermediate 17 as an oil: TLC  $R_f = 0.51$  (1/9 EtOAc/Hexanes);  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.88 (m, 2H), 7.48 (m, 3H), 6.17 (s, 1H), 2.4 (t, 2H,  $J=7.6$ ), 1.72 (q, 2H,  $J=7.6$ ), 1.0 (t, 3H,  $J=7.2$ ); low resolution MS (ES+)  $m/e$  213.0 ( $M+\text{Na}^+$ ).

**Intermediate 18: 1-(phenyl)-1,3-heptanedione**

The title compound was prepared (as described above for Intermediate 16) from 2.2 g (18.3 mmol) of acetophenone and 4.51 g (35 mmol) of ethyl valerate to yield 2.6 grams of Intermediate 18: TLC  $R_f = 0.52$  (1/9 EtOAc/Hexanes);  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.87 (m, 2H), 7.48 (m, 3H), 6.17 (s, 1H), 2.42 (t, 2H,  $J=7.6$ ), 1.67 (m, 2H), 1.4 (m, 2H), 0.95 (t, 3H,  $J=7.2$ ); low resolution MS (ES+)  $m/e$  205.1 (MH+), 227.1 ( $M+\text{Na}^+$ ).

**Intermediate 19: 1-(phenyl)-1,3-decanedione**

To a solution of 1 g (6.1 mmol) of benzoylacetone in 6 mL of THF at  $-78^\circ\text{C}$  was added 8.2 mL (12.3 mmol) of a 1.5 M LDA solution in THF. Warmed to  $0^\circ\text{C}$  over 2 h, then 2.6 mL (12.3 mmol) of 1-iodohexane was added. The mixture was stirred at rt for 3 h before quenching with saturated aqueous  $\text{NH}_4\text{Cl}$  solution and dilution with  $\text{Et}_2\text{O}$ . The organics were dried ( $\text{Na}_2\text{SO}_4$ ), filtered, concentrated, and purified by silica gel chromatography (15/1 hexanes/ $\text{Et}_2\text{O}$ ) to yield Intermediate 19: TLC  $R_f = 0.52$  (1/10  $\text{Et}_2\text{O}$ /Hexanes);  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ , enol form)  $\delta$  7.88 (m, 2H), 7.54-



7.43 (m, 3H), 6.17 (s, 1H), 2.42 (t, 2H,  $J=7.6$ ), 1.66 (m, 2H), 1.4-1.25 (m, 8H), 0.85 (m, 3H); low resolution MS (ES+)  $m/e$  269.0 ( $M+Na^+$ ).

**Intermediate 20: 1-(2-Fluorophenyl)-1,3-butanedione**

The title compound was prepared (as described above for Intermediate 16) from 2.0 g of 2'-fluoroacetophenone and 2.86 mL of EtOAc to yield Intermediate 20:  $^1H$  NMR (400 MHz,  $CDCl_3$ , enol form)  $\delta$  7.70 (dt, 1H,  $J=1.79$ , 7.69), 7.45 (m, 1H), 7.24 (m, 1H), 7.10 (m, 1H), 6.28 (s, 1H), 2.20 (s, 3H).

**Intermediate 21: 1-(3-methylphenyl)-1,3-butanedione**

The title compound was prepared (as described above for Intermediate 16) from 2.72 mL of 3'-methylacetophenone and 3.95 mL of EtOAc to yield 2.94 grams of Intermediate 21 as an oil:  $^1H$  NMR (400 MHz,  $CDCl_3$ , enol form)  $\delta$  7.70 (m, 2H), 7.34 (m, 2H), 6.17 (s, 1H), 2.41 (s, 3H), 2.20 (s, 3H); low resolution MS (ES)  $m/e$  177.1 ( $MH^+$ ).

**Intermediate 22: 1-(4-methoxyphenyl)-1,3-butanedione**

The title compound was prepared (as described above for Intermediate 16) from 3.0 g of 4'-methoxyacetophenone and 3.95 mL of EtOAc. Purification by recrystallization from toluene/hexane afforded the title compound:  $^1H$  NMR (400 MHz,  $CDCl_3$ , enol form)  $\delta$  7.70 (d, 2H,  $J=8.20$ ), 7.34 (d, 2H,  $J=8.20$ ), 6.17 (s, 1H), 3.87 (s, 3H), 2.17 (s, 3H); low resolution MS (ES)  $m/e$  193.1 ( $MH^+$ ).

**Intermediate 23: 1-(2-hydroxyphenyl)-1,3-butanedione**

The title compound was purchased from Aldrich Chemical Co.

**Intermediate 24: 1-(4-trifluoromethylphenyl)-1,3-butanedione**

The title compound was prepared (as described above for Intermediate 16) from 51.43 g of 4'-trifluoroacetophenone and 53.4 mL of EtOAc to yield 35.78 g of Intermediate 24 : TLC analysis:  $R_f = 0.70$  (2/1, hexanes/EtOAc),  $^1H$

NMR (400 MHz, CDCl<sub>3</sub>, enol form)  $\delta$  7.96 (d, 2H,  $J$  = 7.0), 7.7 (d, 2H,  $J$  = 7.0), 6.2 (s, 1H), 2.23 (s, 3H).

**Intermediate 25: 1-(4-trifluoromethylphenyl)-1,3-pentanedione**

The title compound was prepared (as described above for Intermediate 16) from 1.5 g of 4'-trifluoromethylacetophenone and 1.54 mL of methyl propionate to yield 0.69 g of intermediate 25: TLC analysis:  $R_f$  = 0.75 (2/1, hexanes/EtOAc), <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>, enol form)  $\delta$  7.92-7.86 (m, 2H), 7.12 (t, 2H,  $J$  = 8.6), 6.1 (s, 1H), 2.45 (q, 2H,  $J$  = 7.5), 1.21 (t, 3H,  $J$  = 7.5).

**Intermediate 26: 1-(4-Fluorophenyl)-1,3-butanedione**

The title compound was prepared (as described above for Intermediate 16) from 2.42 mL of 4'-fluoroacetophenone and 3.95 mL of EtOAc to yield Intermediate 26 as an oil: <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>, enol form)  $\delta$  7.89 (dd, 2H,  $J$  = 8.90, 5.40), 7.13 (dd, 2H,  $J$  = 17.2, 8.60), 6.13 (s, 1H), 2.19 (s, 3H).

**Intermediate 27: 1-(4-methylphenyl)-1,3-butanedione**

The title compound was prepared (as described above for Intermediate 16) from 2.0 g of 4'-methylacetophenone and 2.9 mL of EtOAc to yield 1.68 grams of Intermediate 27: TLC analysis:  $R_f$  = 0.61 (5/1, hexanes/EtOAc), <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>, enol form)  $\delta$  7.72 (d, 2H,  $J$  = 8.2), 7.24 (d, 2H,  $J$  = 7.4), 6.15 (s, 1H), 2.4 (s, 3H), 2.2 (s, 3H).

**Intermediate 28: 1-(4-chlorophenyl)-1,3-butanedione**

The title compound was prepared (as described above for Intermediate 16) from 2.0 g of 4'-chloroacetophenone and 2.53 mL of EtOAc to yield 1.7 grams of Intermediate 28: TLC analysis:  $R_f$  = 0.73 (2/1, hexanes/EtOAc); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>, enol form)  $\delta$  7.82 (d, 2H,  $J$  = 8.0), 7.42 (d, 2H,  $J$  = 8.0), 6.13 (s, 1H), 2.2 (s, 3H).

**Intermediate 29: 1-(4-isopropoxyphenyl)-1,3-butanedione**

The title compound was prepared (as described above for Intermediate 16) from 2.42 g of 4'-isopropoxyacetophenone and 2.65 mL of EtOAc to yield 2.25 grams of Intermediate 29 : TLC analysis:  $R_f = 0.73$  (2/1, hexanes/EtOAc);  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ , enol form)  $\delta$  7.84 (d, 2H,  $J = 8.9$ ), 6.9 (d, 2H,  $J = 9.0$ ), 6.1 (s, 1H), 4.62 (hept, 1H,  $J = 6.0$ ), 2.16 (s, 3H), 1.36 (d, 6H,  $J = 6.0$ ).

**Intermediate 30: 1-(2-chlorophenyl)-1,3-butanedione**

The title compound was prepared (as described above for Intermediate 16) from 2.0 g of 2'-chloroacetophenone and 2.53 mL of EtOAc to yield 0.5 g of Intermediate 30: TLC analysis:  $R_f = 0.76$  (2/1, hexanes/EtOAc);  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ , enol form)  $\delta$  7.6-7.55 (m, 1H), 7.45-7.43 (m, 1H), 7.4-7.32 (m, 1H), 6.05 (s, 1H), 2.18 (s, 3H).

**Intermediate 31: 1-(3-thienyl)-1,3-butanedione**

The title compound was prepared (as described above for Intermediate 16) from 1.5 g of 3-acetylthiophene and 2.32 mL of EtOAc to yield 1.28 grams of Intermediate 31 as a yellow oil: TLC  $R_f = 0.65$  (2/1, hexanes/EtOAc);  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ , enol form major, ~ 85:15)  $\delta$  8.0 (m, 1H), 7.45 (m, 1H), 7.35 (m, 1H), 6.0 (s, 1H), 2.18 (s, 3H).

**Intermediate 32: 1-(4-Fluorophenyl)-1,3-pentanedione**

The title compound was prepared (as described above for Intermediate 16) from 0.525 g of 4'-fluoroacetophenone and 0.73 mL of methyl propionate to yield 120 mg of Intermediate 32 : TLC analysis:  $R_f = 0.77$  (2/1, hexanes/EtOAc);  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ , enol form)  $\delta$  7.98 (d, 2H,  $J = 8.3$ ), 7.7 (d, 2H,  $J = 8.3$ ), 6.12 (s, 1H), 1.55 (s, 3H); low resolution mass spectrum (ES+) 195 (MH+)

**Intermediate 33: 1-(cyclohexyl)-1,3-butanedione**

The title compound was prepared as described above from 1.5 g of cyclohexyl methyl ketone and 2.32 mL of EtOAc to yield 1.06 grams of

Intermediate 33 as a yellow oil: TLC  $R_f$  = 0.65 (3/1, hexanes/EtOAc);  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ , enol form)  $\delta$  5.5 (s, 1H), 2.19-2.12 (m, 1H), 2.08 (s, 3H), 1.85-1.55 (m, 5 H), 1.41-1.15 (m, 5H).

**Intermediate 34: 1-(2-pyrazinyl)-1,3-butanedione**

To a solution of methyl 2-pyridazinecarboxylate (1.5 g, 10.86 mmol) in 3 mL of acetone and 10 mL of THF was slowly added 590 mg (10.86 mmol) of NaOMe and the resulting mixture was refluxed for 4 hr. The mixture was filtered, the filtrate was acidified and extracted with  $\text{CH}_2\text{Cl}_2$ , and the organics were dried over  $\text{MgSO}_4$  to yield 940 mg (53%) homogeneous by TLC:  $R_f$  = 0.5 (2/1, EtOAc/hexanes);  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ , enol form)  $\delta$  9.27 (d, 1H,  $J$  = 1.4), 8.7 (d, 1H,  $J$  = 2.4), 8.6 (t, 1H,  $J$  = 1.8), 2.26 (s, 3H).

**Intermediate 35: 1-(2-pyridinyl)-1,3-butanedione**

Intermediate 36 was prepared as described above for Intermediate 35 from 2.0 g of methyl 2-pyridylcarboxylate and 4.0 mL of acetone to yield 1.08 g of the title compound (45% yield) as a light brown oil: TLC analysis:  $R_f$  = 0.73 (2/1, hexanes/EtOAc);  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ , mixture of enol/non-enol forms ~80/20)  $\delta$  8.66 (m, 1H), 8.05 (m, 1H), 7.85 (m, 1H), 7.46 (m, 1H), 6.8 (s, 1H), 2.24 (s, 3H), (non-enol form singlets observed at 4.28 and 2.35); low resolution mass spectrum (ES+) 164.1 (MH+).

**Intermediate 36: 1-(2,3,4-trifluorophenyl)-2-butyne-1-one**

Propynyllithium (720 mg, 14.05 mmol) was added to a 0 °C stirred solution of 1.5 g (9.37 mmol) of 2,3,4-trifluorobenzaldehyde and the solution was allowed to stir for 1 h with warming to ambient temperature. The solution was diluted with EtOAc and washed with  $\text{H}_2\text{O}$  and brine. The organics were dried over  $\text{MgSO}_4$  and concentrated (yield of intermediate alcohol, 1.88 g). The crude product was stirred in 30 mL of  $\text{CH}_2\text{Cl}_2$  with 4 g of activated  $\text{MnO}_2$  at 20 °C for 2.5 h. The mixture was filtered through a pad of celite/silica gel and concentrated to yield 1.31 g (47% overall yield) of crude product homogeneous by TLC analysis:  $R_f$  = 0.6 (2/1, hexanes/EtOAc);  $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ )  $\delta$  7.9-7.8 (m, 1H), 7.1-6.95 (m 1H), 2.13 (s, 3H).

Intermediates 37-43 were prepared analogous to the method described above for the preparation of Intermediate 36.

**Intermediate 37: 1-(2-fluoro-3-trifluoromethylphenyl)-1,3-pentanedione**

The title compound was prepared (as described above for intermediate 36) from 0.5 g of 2'-fluoro-3'-trifluoromethylbenzaldehyde and 1.03 g of MnO<sub>2</sub> to yield 0.46 g (75% overall yield) of Intermediate 37: <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 8.19 (m, 1H), 7.84 (m, 1H), 7.32 (m, 1H), 2.1 (s, 3H).

**Intermediate 38: 1-(2,4,5-trifluorophenyl)-2-butyne-1-one**

Intermediate 38 was prepared (as described above for intermediate 36) from 1-propynyllithium, 0.5 g of 2',4',5'-trifluorobenzaldehyde, and 0.87 g of MnO<sub>2</sub> to yield 0.39 g (63% overall yield) of the title compound: <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.91-7.82 (m, 1H), 7.05-6.96 (m, 1H), 2.14 (s, 3H). For the alcohol intermediate: <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.54-7.48 (m, 1H), 6.97-6.88 (m, 1H), 5.64 (s, 1H), 1.91 (s, 3H).

**Intermediate 39: 1-(2,4-difluorophenyl)-2-butyne-1-one**

Intermediate 39 was prepared (as described above for intermediate 36) from 1-propynyllithium, 500 mg of 2',4'-difluorobenzaldehyde, and 1.5 g of MnO<sub>2</sub> to yield 400 mg of the title compound homogeneous by TLC (R<sub>f</sub> = 0.63, 2/1 hexanes/EtOAc); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 8.13-8.06 (m, 1H), 6.99-6.94 (m, 1H), 6.91-6.85 (m, 1H), 2.14 (s, 3H).

**Intermediate 40: 1-(2,3-difluorophenyl)-2-butyne-1-one**

Intermediate 40 was prepared (as described above for intermediate 36) from 1-propynyllithium, 500 mg of 2,3-difluorobenzaldehyde, and 1.5 g of MnO<sub>2</sub> to yield 440 mg of the title compound homogeneous by TLC (R<sub>f</sub> = 0.5, 2/1 hexanes/EtOAc); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.82-7.86 (m, 1H), 7.43-7.35 (m, 1H), 7.22-7.15 (m, 1H), 2.15 (s, 3H).

**Intermediate 41: 1-(4-nitrophenyl)-2-butyne-1-one**

Intermediate 41 was prepared (as described above for intermediate 36) from 1-propynyllithium, 1.0 gram of 4'-nitrobenzaldehyde, and 1.0 g of MnO<sub>2</sub> to yield 430 mg of the title compound homogeneous by TLC analysis: R<sub>f</sub> = 0.55 (2/1, hexanes/EtOAc), <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 8.32-8.28 (m, 4H), 2.12 (s, 3H).

**Intermediate 42: 1-(3-nitrophenyl)-2-butyne-1-one**

Intermediate 42 was prepared (as described above for Intermediate 36) from 1-propynyllithium and 0.5 g of 3'-nitrobenzaldehyde and 1.5 g MnO<sub>2</sub> to yield 280 mg of the title compound: TLC analysis: R<sub>f</sub> = 0.50 (2/1, hexanes/EtOAc), <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 8.95 (t, 1H, J = 1.8), 8.47 (t, 1H, J = 1.7), 8.44 (t, 1H, J = 1.7), 7.7 (t, 1H, J = 8.0), 2.23 (s, 3H).

**Intermediate 43: 1-(2-thiazolyl)-2-butyne-1-one**

Intermediate 43 was prepared (as described above for Intermediate 36) from 1-propynyllithium, 2-thiophene carboxaldehyde (108 mg) and 300 mg of MnO<sub>2</sub> to yield 65 mg of the title compound : TLC analysis: R<sub>f</sub> = 0.50 (1/1, hexanes/EtOAc); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 8.1 (d, 1H, J = 2.8), 7.71 (d, 1H, J = 2.9), 2.21 (s, 3H); low resolution mass spectrum (ES+) 152.1 (MH<sup>+</sup>).

**Intermediate 44: 3-Bromo-4,4,4-trifluoro-1-phenylbut-2-en-1-one  
(mixture of E- and Z-isomers, ~ 85:15)**

To a solution of 7.58 g (28.9 mmol) of triphenylphosphine in 60 mL of CH<sub>2</sub>Cl<sub>2</sub> at 0 °C was added 1.49 mL (28.9 mmol) of bromine dropwise. The solution was warmed to ambient temperature and 5.0 g (23.1 mmol) of trifluoroacetyl-acetophenone in 25 mL CH<sub>2</sub>Cl<sub>2</sub> and 6.45 mL (46.3 mmol) of triethylamine were added. The mixture was stirred for two hours until no further change was observed by TLC analysis (3/1 hexanes/EtOAc). The mixture was washed with 1.0 N HCl, H<sub>2</sub>O, and brine. The organics were dried over MgSO<sub>4</sub>, concentrated, and the residue purified by silica gel chromatography (20/1 hexanes/EtOAc) to give 1.6 g of impure material and 1.15 g (25% yield) of Intermediate 44 as an oil: TLC R<sub>f</sub> = 0.74 (3/1,

hexanes/EtOAc);  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.95 – 7.5 (series of multiplets, 5H);  $^{19}\text{F}$  NMR (282 MHz,  $\text{CDCl}_3$ ) singlet,  $\delta$  –100.5 ppm)

**Intermediate 45: (2S)-2-amino-3-{4-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]phenyl}propanoic acid**

Dianion coupling method: To a solution of 5.4 g (19.3 mmol) of N-(Boc)-L-Tyrosine in 5 mL of DMSO and 10 mL of  $\text{H}_2\text{O}$  was added 1.7 g (42.5 mmol) of freshly ground NaOH pellets. The mixture was heated to 55–60 °C and a solution of 6.5 g (23.1 mmol) of mesylate intermediate 15 in 10 mL of DMSO was added dropwise over ~ 5 min. The resulting mixture was stirred vigorously for 8 hr at 55–60 °C, cooled to 20 °C, and poured into 100 mL of  $\text{H}_2\text{O}$  and extracted with  $\text{Et}_2\text{O}$  (2 x 50 mL). The aqueous phase was separated and acidified to pH~2–3 with glacial acetic acid/HCl. The resulting slurry was extracted with EtOAc several times, dried ( $\text{MgSO}_4$ ), and the organics were concentrated to a tan oil. The oil was triturated with hexanes/EtOAc (3/1) to give a cream-colored solid. This solid was recrystallized from hot MeOH/  $\text{H}_2\text{O}$  (2.5/1) to yield 5.5 g (61 % yield) of Boc-protected intermediate as a white solid after drying in a vacuum oven at 60 °C for ~14 hr. To a solution of 43.54 g (93.3 mmol) of the above Boc-protected intermediate in 150 mL of dioxane was added 200 mL of 4.0 N HCl in dioxane solution and the mixture was stirred over 7 hr with the gradual formation of a white solid. The solids were filtered (rinsing with  $\text{Et}_2\text{O}$ ), suspended into 1 L of  $\text{H}_2\text{O}$ , and the pH was adjusted to pH~5 with 15% NaOH. The resulting white solids were collected via filtration, washed with  $\text{H}_2\text{O}$  and  $\text{CH}_3\text{CN}$ , and dried overnight at 70 °C in a vacuum oven to give 32.9 g (96%) of intermediate 45 as a white solid:  $^1\text{H}$  NMR ( $\text{DMSO}-d_6$ , 300 MHz)  $\delta$  7.95 (m, 2H), 7.55 (m, 3H), 7.22 (d, 2H,  $J$  = 8.5), 6.9 (d, 2H,  $J$  = 8.5), 4.21 (t, 2H,  $J$  = 6.4), 3.1 (dd, 1H,  $J$  = 14.3, 4.1), 4.17 (m, 1H), 2.96 (t, 2H,  $J$  = 6.6), 2.83 (dd, 1H,  $J$  = 14.3, 8.4), 2.53 (s, 3H),  $\alpha$ -methine proton assumed under  $\text{H}_2\text{O}$  peak;; low resolution MS ( $\text{ES}^+$ )  $m/e$  367.4 ( $\text{MH}^+$ );

Alternative Mitsunobu coupling method.

To a solution of 4.46 g (21.9 mmol) of alcohol Intermediate 8, 5.88 g (19.9 mmol) of N-Boc-L-tyrosine methyl ester, and 6.26 g (23.9 mmol) of triphenylphosphine in 150 mL of toluene at 45 °C was added dropwise a solution of 3.81 g (21.9 mmol) of DEAD in 50 mL of toluene. The resulting clear solution was stirred at 20 °C for 8 hr, concentrated, and the residue re-dissolved into Et<sub>2</sub>O (200 mL). To this was added 75 mL of 1N NaOH solution. The mixture was rapidly stirred for 30 min until phenol no longer present by TLC. The aqueous phase was extracted with EtOAc and the combined organics were dried (MgSO<sub>4</sub>) and concentrated. The crude tan-colored solid was purified by silica gel chromatography (CH<sub>2</sub>Cl<sub>2</sub>/ Et<sub>2</sub>O, 12/1 to 8/1) to give an impure product. A second column eluting with CH<sub>2</sub>Cl<sub>2</sub>/hexane (4/1), CH<sub>2</sub>Cl<sub>2</sub>, and CH<sub>2</sub>Cl<sub>2</sub>/ Et<sub>2</sub>O (10/1) gave 5.5 g (57% yield ) of a clear yellow oil. To this oil in 150 mL of THF was added dropwise 13.7 mL of 1.0 M LiOH in H<sub>2</sub>O solution. To this was added an additional 75 mL of THF and 5 mL of MeOH to give a clear solution. The solution was stirred at 20 °C and the progress of the reaction was monitored by TLC (CHCl<sub>3</sub>/MeOH, 9/1). The THF was removed at reduced pressure and the aqueous phase extracted with EtOAc/1.0 N HCl. The organics were washed with H<sub>2</sub>O, dried over MgSO<sub>4</sub>, concentrated, and the resulting crude oil was purified by silica gel chromatography (EtOAc/hexanes, 1/1 to 8/1 gradient) to give 3.22 g (61%) of a foam. To this foam in 15 mL of dioxane was added dropwise 15 mL of 4.0 N HCl in dioxane solution at 20 °C. Stirred the mixture for 4 hr producing a white solid precipitate. The solids were filtered washing with Et<sub>2</sub>O, then dissolved into H<sub>2</sub>O and the pH adjusted down to ~pH 5 with the dropwise addition of 1.0 N NaOH solution. The resulting white precipitate was collected by filtration and dried in a vacuum oven at 60 °C overnight to yield 1.72 g of Intermediate 45 as a white solid.

**Intermediate 46: (2S)-2-amino-3-(4-(2-(2-(4-fluorophenyl)-5-methyl-1,3-oxazol-4-yl)ethoxy)phenyl)propanoic acid**

DEAD (1.26 mL, 8.16 mmol) was added dropwise to a solution of 2.14 g (8.16 mmol) of triphenylphosphine in 16 mL of THF at 0 °C. This solution was then added to a mixture of 2.41 g (8.16 mmol) of N-(Boc)-L-Tyrosine



5 methyl ester (see, e.g., A. Kolodziejczyk, et al., J. Org. Chem., 46(9), pp 1944-  
1946 (1981)) and Intermediate 9 in 16 mL of THF at 0 °C. The solution was  
10 stirred at 20 °C for 20 h and concentrated to an oil. The crude material was  
purified by silica gel chromatography (4:1 hexanes:EtOAc) to yield 2.78 grams  
5 (69% yield) of the protected amino acid Intermediate. Deprotection was  
achieved as follows: Ester hydrolysis - To a solution of 2.77 g (5.58 mmol) of  
the above ester in 45 mL of THF and 15 mL of H<sub>2</sub>O at 0 °C was added 8.5 mL  
15 (8.5 mmol) of 1.0 M aqueous LiOH solution. After 2 h at 20 °C, EtOAc was  
added and the mixture was acidified with 0.1 N HCl. The aqueous was  
10 extracted with EtOAc and the organics were dried (MgSO<sub>4</sub>) and concentrated  
to give a white glass-like solid (2.67 g). Boc Deprotection - The solid from the  
20 ester hydrolysis above (2.64 g, 5.47 mmol) was taken into 60 mL CH<sub>2</sub>Cl<sub>2</sub> and  
treated with 30 mL of TFA with stirring for 1 h. The solution was concentrated,  
the residue was dissolved into ~400 mL of H<sub>2</sub>O, and the pH was adjusted to  
25 ~pH 7 with aqueous 1.0 N NaOH solution. The suspension was filtered and  
the collected solids were dried for 48 h at 70 °C under vacuum to yield 1.61 g  
15 (77% yield for deprotection, 53% overall yield) of Intermediate 46 as a fluffy  
beige-colored solid: <sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 300 MHz) δ 7.93 (dd, 2H, J = 8.7,  
30 5.4), 7.3 (t, 2H, J = 8.7), 7.12 (d, 2H, J = 8.0), 6.83 (d, 2H, J = 8.2), 4.15 (t, 2H, J  
= 6.5), 3.02 (dd, 1H, J = 14.2, 4.3), 2.89 (t, 2H, J = 6.5), 2.76 (dd, 1H, J = 14.4,  
20 8.2), 2.34 (s, 3H); low resolution MS (ES+) *m* / *e* 384.9 (MH<sup>+</sup>).

35  
Intermediates 47-51 were prepared analogous to the method described  
above for the preparation of Intermediate 46

40  
25  
**Intermediate 47: (2S)-2-amino-3-(4-[2-(5-methyl-2-phenyl-1,3-thiazol-4-yl)ethoxy]phenyl)propanoic acid**

45  
Intermediate 47 was prepared from 2.41 g of N-(Boc)-L-Tyrosine  
methyl ester and 1.79 g of Intermediate 10 to yield 1.61 g (51% overall yield)  
30 of the title compound as a solid: <sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 400 MHz) δ 7.82 (d, 2H,  
J = 7.5), 7.42 (m, 3H), 7.12 (d, 2H, J = 8.2), 6.82 (d, 2H, J = 8.4), 4.23 (t, 2H, J  
= 6.6), 3.1 (t, 2H, J = 6.6), 2.83 (dd, 1H, J = 14.1, 3.7), 2.53 (s, 3H), second  
50

proton of methylene assumed under H<sub>2</sub>O peak; low resolution MS (ES<sup>+</sup>)*m/e* 383.1 (MH<sup>+</sup>);

**Intermediate 48: (2S)-2-amino-3-[4-{2-[2-(4-fluorophenyl)-5-methyl-1,3-thiazol-4-yl]ethoxy}phenyl]propanoic acid**

Intermediate 48 was prepared from 224 mg of N-(Boc)-L-Tyrosine methyl ester and 180 mg of Intermediate 11 to yield 149 mg (51% overall yield) of the title compound as a solid: <sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 400 MHz) δ 7.82 (dd, 2H, *J* = 8.6, 5.5), 7.23 (t, 2H, *J* = 8.7), 7.08 (d, 2H, *J* = 8.4), 6.8 (d, 2H, *J* = 8.5), 4.56 (br s, 1H), 4.19 (t, 2H, *J* = 6.5), 3.56 (m, 1H), 3.03 (t, 2H, *J* = 6.6), 2.95 (m, 1H), 2.79 (dd, 1H, *J* = 14.5, 7.5), 2.37 (s, 3H); low resolution MS (ES<sup>+</sup>)*m/e* 400.9 (MH<sup>+</sup>).

**Intermediate 49: (2S)-2-amino-3-[4-{2-[2-(4-methoxyphenyl)-5-methyl-1,3-oxazol-4-yl]ethoxy}phenyl]propanoic acid**

Intermediate 49 was prepared from 735 mg of N-(Boc)-L-Tyrosine methyl ester and 580 mg of Intermediate 12 to yield 220 mg (22% overall yield) of the title compound as a solid. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 400 MHz) δ 8.16 (d, 2H, *J* = 8.2), 7.92 (d, 2H, *J* = 8.4), 7.21 (d, 2H, *J* = 8.5), 6.93 (d, 2H, *J* = 8.5), 4.25 (t, 2H, *J* = 6.5), 3.1 (dd, 1H, *J* = 14.4, 4.8), 3.0 (t, 2H, *J* = 6.5), 2.76 (dd, 1H, *J* = 14.4, 7.9), 2.46 (s, 3H); low resolution MS (ES<sup>+</sup>)*m/e* 434.9 (MH<sup>+</sup>).

**Intermediate 50: (2S)-2-amino-3-[4-{2-[5-methyl-2-(4-(trifluoromethyl)phenyl)-1,3-oxazol-4-yl]ethoxy}phenyl]propanoic acid**

Intermediate 50 was prepared from 185 mg of N-(Boc)-L-Tyrosine methyl ester and 170 mg of Intermediate 13 to yield 130 mg (48 % overall yield) of the title compound as a solid: <sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 400 MHz) δ 8.16 (d, 2H, *J* = 8.2), 7.92 (d, 2H, *J* = 8.4), 7.21 (d, 2H, *J* = 8.5), 6.93 (d, 2H, *J* = 8.5), 4.25 (t, 2H, *J* = 6.5), 3.1 (dd, 1H, *J* = 14.4, 4.8), 3.0 (t, 2H, *J* = 6.5), 2.76 (dd, 1H, *J* = 14.4, 7.9), 2.46 (s, 3H); low resolution MS (ES<sup>+</sup>)*m/e* 434.9 (MH<sup>+</sup>).

**Intermediate 51: (2S)-2-amino-3-(4-{2-[2-(4-isopropoxyphenyl)-5-methyl-1,3-oxazol-4-yl]ethoxy}phenyl)propanoic acid**

Intermediate 51 was prepared from 598 mg of N-(Boc)-L-Tyrosine methyl ester and 530 mg of Intermediate 14 to yield 540 mg (63% overall yield) of the title compound as a solid. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 400 MHz) δ 7.77 (d, 2H, J = 8.7), 7.12 (d, 2H, J = 8.6), 6.9 (d, 2H, J = 8.9), 6.82 (d, 2H, J = 8.5), 4.65 (hept, 1H, J = 6.0), 4.13 (t, 2H, J = 6.6), 3.0 (dd, 1H, J = 14.4, 4.5), 2.85 (t, 2H, J = 6.4), 2.76 (dd, 1H, J = 14.4, 8.0), 2.31 (s, 3H), 1.25 (d, 6H, J = 6.0); low resolution MS (ES<sup>+</sup>) m / e 425.2 (MH<sup>+</sup>).

**Intermediate 52: (2S)-2-amino-3-(4-[2-(5-ethyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]phenyl)propanoic acid**

To a mixture of 640 mg (2.95 mmol, 1.1 eq.) of alcohol intermediate 8B, 790 mg (2.67 mmol, 1.0 eq.) of N-(Boc)-L-Tyrosine methyl ester, and 773 mg (2.95 mmol, 1.1 eq.) of triphenylphosphine in 15 mL of anhydrous toluene at 50 °C was added 702 mg (3.47 mmol, 1.3 eq) of DIAD as a solution in 10 mL of toluene. Stirred 2.5 hr at 50 °C. Cooled, concentrated, purified by loading directly onto a silica gel column. Elution with EtOAc/Hexanes (1/20 to 1/1) gave 1.15 grams (87% yield) of BOC-protected intermediate methyl ester of sufficient purity to carry forward into the following deprotection reactions [<sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 300 MHz) δ 8.07-8.03 (m, 2H), 7.47 (m, 3H), 7.03 (d, 2H, J = 8.5), 6.85 (d, 2H, J = 8.5), 4.58-4.51 (m, 1H), 4.26 (t, 2H, J = 6.5), 3.72 (s, 3H), 3.06-3.0 (m, 4H), 2.78 (q, 2H, J = 7.5), 1.43 (s, 9H), 1.34 (t, 3H, J = 7.6)];

**Deprotections:** To a THF/H<sub>2</sub>O (15 mL/5 mL) solution of 900 mg of the above protected amino acid at 0 °C was added 2.64 mL of a 1.0 M LiOH in water solution. Stirred 2.5 hrs. Diluted the mixture with 30 mL of EtOAc and the pH was adjusted downward to ~6 by the addition of 0.1 M HCl solution. The aqueous was extracted with EtOAc. The organics were dried (MgSO<sub>4</sub>) and concentrated, then taken into DCM (15 mL) at 0 °C and treated with 7.5 mL of TFA. The solution was concentrated after 1 hr, then concentrated several times from DCM to yield 1.4 g of crude intermediate 52 (TFA salt). This solid was stirred with 50 mL of Et<sub>2</sub>O for 30 min. The solids were collected via

filtration and dried under vacuum for 12 h to yield 740 mg (67%) of the title compound as an off white solid:  $^1\text{H}$  NMR (DMSO- $d_6$ , 300 MHz)  $\delta$  8.18 (br s, 2H), 7.93-7.90 (m, 2H), 7.51-7.48 (m, 3H), 7.15 (d, 2H,  $J=8.5$ ), 6.89 (d, 2H,  $J=8.4$ ), 4.18 (t, 2H,  $J=7.5$ ), 4.11 (m, 1H), 3.01-2.99 (m, 2H), 2.95-2.90 (m, 2H), 2.75 (q, 2H,  $J=7.5$ ), 1.23 (t, 3H,  $J=7.5$ ); low resolution MS (ES $^+$ )  $m/e$  381.07 (MH $^+$ ).

**Intermediate 53: (2S)-2-amino-3-{4-[2-(2-phenyl-5-propyl-1,3-oxazol-4-yl)ethoxy]phenyl}propanoic acid**

Amino acid Intermediate 53 was prepared as described for the preparation of Intermediate 46. From 460 mg of N-(Boc)-L-Tyrosine methyl ester and 360 mg of Intermediate 8C was prepared 410 mg (67% overall yield) of the title compound as a solid:  $^1\text{H}$  NMR (DMSO- $d_6$ , 300 MHz)  $\delta$  7.90-7.87 (m, 2H), 7.48-7.46 (m, 3H), 7.13 (d, 2H,  $J=8.4$ ), 6.85 (d, 2H,  $J=8.4$ ), 4.17 (t, 2H,  $J=6.3$ ), 4.0 (t, 1H,  $J=6.3$ ), 2.99 (t, 1H,  $J=6.3$ ), 2.92 (t, 1H,  $J=6.3$ ), 2.69 (t, 2H,  $J=7.2$ ), 1.66 (q, 2H,  $J=7.5$ ), 0.93 (t, 3H,  $J=7.2$ ); low resolution MS (ES $^+$ )  $m/e$  395.09 (MH $^+$ ); TLC  $R_f$  = 0.24 (10/1, DCM/MeOH).

**Intermediate 54: (2S)-2-amino-3-{4-[(5-methyl-2-phenyl-1,3-oxazol-4-yl)methoxy]phenyl}propanoic acid**

Amino acid Intermediate 54 was prepared as described for the preparation of Intermediate 46. From 187 mg of N-(Boc)-L-Tyrosine methyl ester and 120 mg of Intermediate 8D (with DIAD replacing DEAD) was prepared 116 mg (53% overall yield) of the title compound as a solid.  $^1\text{H}$  NMR (DMSO- $d_6$ , 300 MHz)  $\delta$  7.89-7.85 (m, 2H), 7.43-7.36 (m, 3H), 7.12 (d, 2H,  $J=8.5$ ), 6.9 (d, 2H,  $J=8.5$ ), 4.9 (s, 2H), 3.63 (m, 1H), 3.0 (dd, 1H,  $J=14.4$ , 4.5), 2.85 (dd, 1H,  $J=14.4$ , 8.1), 2.37 (s, 3H); low resolution MS (ES $^+$ )  $m/e$  353.07 (MH $^+$ ).

**Intermediate 55: (2S)-2-amino-3-{4-[3-(5-methyl-2-phenyl-1,3-oxazol-4-yl)propoxy]phenyl}propanoic acid**

Amino acid Intermediate 55 was prepared as described for the preparation of Intermediate 46. From 503 mg of N-(Boc)-L-Tyrosine methyl ester and 370 mg of Intermediate 8E to yield 443 mg (68% overall yield) of the title compound as a solid. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 300 MHz) δ 7.90-7.86 (m, 2H), 7.52-7.44 (m, 3H), 7.14 (d, 2H, J=8.5), 6.86 (d, 2H, J=8.4), 3.94 (t, 2H, J=6.2), 3.75 (m, 1H), 3.03 (dd, 1H, J=14.3, 5.1), 2.90 (dd, 1H, J=14.3, 7.2), 2.59 (t, 2H, J=7.3), 2.27 (s, 3H), 2.0 (m, 2H); TLC R<sub>f</sub> = 0.22 (10/1, DCM/MeOH); low resolution MS (ES+) m/e 381.12 (MH<sup>+</sup>).

**Intermediate 56: (2S)-2-amino-3-{4-[2-(5-ethyl-2-(4-fluorophenyl)-1,3-thiazol-4-yl)ethoxy]phenyl}propanoic acid**

Intermediate 56 was prepared as described above for the preparation of intermediate 52. From 534 mg of N-(Boc)-L-Tyrosine methyl ester and 500 mg of Intermediate 11B was prepared 900 mg of BOC-protected intermediate methyl ester (94% yield; <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) δ 7.9 (dd, 2H, 8.7, 5.2), 7.13 (t, 2H, J=8.6), 7.04 (d, 2H, J=8.5), 6.85 (d, 2H, J=8.6), 4.97 (d, 1H, J=7.9), 4.55 (m, 1H), 4.32 (t, 2H, J=6.8), 3.73 (s, 3H), 3.21 (t, 2H, J=6.7), 3.10-3.02 (m, 2H), 2.89 (q, 2H, J=7.5), 1.44 (s, 9H), 1.36 (t, 3H, J=7.5); From 850 mg of protected intermediate was prepared 900 mg of intermediate 56 as a solid. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 300 MHz) δ 8.0 (br s, 2H), 7.91-7.87 (m, 2H), 7.30 (dt, 2H, J=8.8, 2.1), 7.14 (d, 2H, J=8.6), 6.90 (d, 2H, J=8.6), 4.27 (t, 2H, J=6.5), 4.12 (m, 1H), 3.11 (t, 2H, J=6.5), 3.0 (m, 2H), 2.86 (q, 2H, J=7.5), 1.26 (t, 3H, J=7.5); low resolution MS (ES+) m/e 414.85 (MH<sup>+</sup>).

**Intermediate 57: (2S)-2-amino-3-{4-[2-(5-ethyl-2-phenyl-1,3-thiazol-4-yl)ethoxy]phenyl}propanoic acid**

Intermediate 57 was prepared as described above for the preparation of intermediate 52. From 575 mg of N-(Boc)-L-Tyrosine methyl ester and 500 mg of Intermediate 9B was prepared 940 mg of BOC-protected intermediate methyl ester (94% yield; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 300 MHz) δ 7.93 (m, 2H), 7.47-7.41 (m, 3H), 7.04 (d, 2H, J=8.4), 6.86 (d, 2H, J=8.6), 4.96 (d, 1H, J=8.2), 4.56 (m, 1H), 4.34 (t, 2H, J=6.8), 3.73 (s, 3H), 3.23 (t, 2H, J=6.9), 3.04 (m,

5 2H), 2.90 (q, 2H,  $J=7.5$ ), 1.44 (s, 9H), 1.37 (t, 3H,  $J=7.5$ ); From 900 mg of  
protected intermediate was prepared 900 mg of intermediate 57 as a solid.  $^1\text{H}$   
10 NMR (DMSO- $d_6$ , 300 MHz)  $\delta$  8.16 (br s, 2H), 7.85 (m, 2H), 7.5-7.43 (m, 3H),  
7.14 (d, 2H,  $J=8.6$ ), 6.90 (d, 2H,  $J=8.6$ ), 4.27 (t, 2H,  $J=6.5$ ), 4.12 (m, 1H),  
5 3.12 (t, 2H,  $J=6.5$ ), 3.0 (m, 2H), 2.86 (q, 2H,  $J=7.5$ ), 1.26 (t, 3H,  $J=7.5$ ).

15 **Intermediate 58: (2S)-2-amino-3-(4-{2-[5-ethyl-2-(4-fluorophenyl)-1,3-oxazol-4-yl]ethoxy}phenyl)propanoic acid**

Intermediate 58 was prepared as described above for the preparation of  
10 intermediate 52. From 230 mg of N-(Boc)-L-Tyrosine methyl ester and 200 mg  
20 of Intermediate 10B was prepared 350 mg of BOC-protected intermediate  
methyl ester (88% yield;  $^1\text{H}$  NMR (DMSO- $d_6$ , 300 MHz)  $\delta$  8.07-8.01 (m, 2H),  
7.15 (t, 2H,  $J=8.7$ ), 7.04 (d, 2H,  $J=8.5$ ), 6.84 (d, 2H,  $J=8.5$ ), 4.50 (m, 1H),  
4.24 (t, 2H,  $J=6.5$ ), 3.72 (s, 3H), 3.04-2.99 (m, 2H), 2.77 (q, 2H,  $J=7.5$ ), 1.44  
25 (s, 9H), 1.33 (t, 3H,  $J=7.5$ ); From 340 mg of BOC-protected ester  
intermediate was prepared 350 mg of intermediate 58 as a crude off-white  
colored solid.  $^1\text{H}$  NMR (DMSO- $d_6$ , 400 MHz)  $\delta$  8.13 (br s, 2H), 7.91 (dd, 2H,  $J$   
30  $=8.8$ , 5.3), 7.3 (t, 2H,  $J=8.9$ ), 7.10 (d, 2H,  $J=8.6$ ), 6.85 (d, 2H,  $J=8.4$ ), 4.13 (t,  
2H,  $J=6.4$ ), 4.08 (m, 1H), 2.98-2.95 (m, 2H), 2.88 (t, 2H,  $J=6.5$ ), 2.70 (q, 2H,  
20  $J=7.4$ ), 1.19 (t, 3H,  $J=7.5$ ); low resolution MS (ES+)  $m/e$  399.13 (MH $^+$ ).

35 **Intermediate 59: (2S)-2-amino-3-(4-{3-[5-ethyl-2-(4-fluorophenyl)-1,3-oxazol-4-yl]propoxy}phenyl)propanoic acid**

Intermediate 59 was prepared as described above for the preparation of  
40 intermediate 52. From 370 mg of N-(Boc)-L-Tyrosine methyl ester and 310 mg  
of Intermediate 10C was prepared 575 mg of BOC-protected intermediate  
methyl ester (87% yield;  $^1\text{H}$  NMR (CDCl $_3$ , 400 MHz)  $\delta$  7.95 (dd, 2H,  $J=8.6$ ,  
5.4), 7.09 (t, 2H,  $J=8.7$ ), 6.99 (d, 2H,  $J=8.4$ ), 6.79 (d, 2H,  $J=8.5$ ), 4.50 (m,  
45 1H), 3.92 (t, 2H,  $J=7.1$ ), 3.68 (s, 3H), 3.05-2.93 (m, 2H), 2.69-2.59 (m, 2H),  
2.16-2.08 (m, 2H), 1.39 (s, 9H), 1.17 (t, 3H,  $J=7.6$ ); From 562 mg of BOC-  
30 protected ester intermediate was prepared the crude TFA salt of intermediate  
59 as a solid (87%);  $^1\text{H}$  NMR (DMSO- $d_6$ , 400 MHz)  $\delta$  7.97-7.9 (m, 2H), 7.33 (t,

2H,  $J = 8.8$ ), 7.14 (d, 2H,  $J = 8.3$ ), 6.88 (d, 2H,  $J = 8.1$ ), 3.97 (m, 3H), 3.25-3.01 (m, 2H), 2.68-2.60 (m, 4H), 2.04-1.99 (m, 2H), 1.16 (t, 3H,  $J = 7.4$ ); low resolution MS (ES+)  $m/e$  413.18 ( $MH^+$ ).

**Intermediate 60: (2S)-2-amino-3-{4-[3-(5-ethyl-2-phenyl-1,3-oxazol-4-yl)propoxy]phenyl}propanoic acid**

Intermediate 60 was prepared as described above for the preparation of intermediate 52. From 266 mg of N-(Boc)-L-Tyrosine methyl ester and 208 mg of Intermediate 8F was prepared 360 mg of BOC-protected intermediate methyl ester (79% yield;  $^1H$  NMR ( $CDCl_3$ , 300 MHz)  $\delta$  8.06-8.03 (m, 2H), 7.46-7.44 (m, 3H), 7.04 (d, 2H,  $J = 8.4$ ), 6.85 (d, 2H,  $J = 8.4$ ), 4.97 (d, 1H,  $J = 7.6$ ), 4.57 (m, 1H), 3.99 (t, 2H,  $J = 6.0$ ), 3.74 (s, 3H), 3.05 (m, 2H), 2.77-2.65 (m, 4H), 2.19 (t, 2H,  $J = 6.6$ ), 1.45 (s, 9H), 1.24 (t, 3H,  $J = 7.5$ ); From 352 mg of BOC-protected ester intermediate was prepared the crude TFA salt of intermediate 60. The TFA salt was suspended into water and the pH adjusted to  $\sim 7.0$  with 10% NaOH. The resulting solids were filtered and dried to yield intermediate 60;  $^1H$  NMR ( $DMSO-d_6$ , 400 MHz)  $\delta$  7.88-7.85 (m, 2H), 7.48-7.43 (m, 3H), 7.11 (d, 2H,  $J = 8.5$ ), 6.82 (d, 2H,  $J = 8.5$ ), 3.92 (t, 2H,  $J = 6.2$ ), 3.55 (m, 1H), 3.0 (dd, 1H,  $J = 14.3, 4.8$ ), 2.8 (dd, 1H,  $J = 14.3, 7.7$ ), 2.67-2.56 (m, 4H), 2.02-1.96 (m, 2H), 1.13 (t, 3H,  $J = 7.4$ ); low resolution MS (ES+)  $m/e$  392.9 ( $MH^+$ ); TLC  $R_f = 0.44$  (4/1, DCM/MeOH).

**Intermediate 61: (2S)-2-amino-3-{4-[5-ethyl-2-phenyl-1,3-oxazol-4-yl)methoxy]phenyl}propanoic acid**

Intermediate 61 was prepared as described above for the preparation of intermediate 52. From 465 mg of N-(Boc)-L-Tyrosine methyl ester and 320 mg of Intermediate 8G was prepared 520 mg of BOC-protected intermediate methyl ester (69% yield;  $^1H$  NMR ( $CDCl_3$ , 300 MHz)  $\delta$  8.09-8.06 (m, 2H), 7.47-7.46 (m, 3H), 7.08 (d, 2H,  $J = 8.7$ ), 6.97 (d, 2H,  $J = 8.7$ ), 5.02 (s, 2H), 5.01 (s, 1H), 4.59-4.55 (m, 1H), 3.74 (s, 3H), 3.08-3.02 (m, 2H), 2.83 (q, 2H,  $J = 7.5$ ), 1.45 (s, 9H), 1.32 (t, 3H,  $J = 7.5$ ); From 510 mg of BOC-protected ester intermediate was prepared the crude TFA salt of intermediate 61. The TFA salt was suspended into water and the pH adjusted to  $\sim 7.0$  with 10% NaOH.

5 The resulting solids were filtered and dried to yield 315 mg of intermediate 61;  
1H NMR (DMSO-d<sub>6</sub>, 300 MHz) δ 7.96-7.94 (m, 2H), 7.52-7.51 (m, 3H), 7.18  
(d, 2H, J = 8.4), 6.95 (d, 2H, J = 8.4), 4.97 (s, 3H), 3.10-3.05 (m, 1H), 2.86-2.79  
10 (m, 4H), 1.22 (t, 3H, J = 7.5); low resolution MS (ES+) m / e 366.86 (MH<sup>+</sup>); TLC  
5 R<sub>f</sub> = 0.52 (4/1, DCM/MeOH).

15 **Intermediate 62: (2S)-2-amino-3-{4-[(5-ethyl-2-phenyl-1,3-thiazol-4-yl)methoxy]phenyl}propanoic acid**

Intermediate 62 was prepared as described above for the preparation of  
10 intermediate 52. From 3.27 g of N-(Boc)-L-Tyrosine methyl ester and 2.43 g of  
20 Intermediate 9C was prepared 5.07 g of BOC-protected intermediate methyl  
ester (92% yield; 1H NMR (CDCl<sub>3</sub>, 300 MHz) δ 7.89-7.86 (m, 2H), 7.42-7.38  
(m, 3H), 7.01 (d, 2H, J = 6.3), 6.94 (d, 2H, J = 6.3), 5.11 (s, 2H), 4.94 (d, 1H, J  
= 6.0), 4.53-4.50 (m, 1H), 3.68 (s, 3H), 3.02-2.98 (m, 2H), 2.90 (q, 2H, J = 5.7),  
25 1.39 (s, 9H), 1.30 (t, 3H, J = 5.7); From 5.05 g of BOC-protected ester  
15 intermediate was prepared the crude TFA salt of intermediate 62 as in  
example 52. The TFA salt was suspended into water and the pH adjusted to  
30 ~7.0 with 10% NaOH. The resulting solids were filtered and dried to yield 4.28  
g of intermediate 62; 1H NMR (DMSO-d<sub>6</sub>, 400 MHz) δ 7.85-7.83 (m, 2H), 7.45-  
20 7.43 (m, 3H), 7.16-7.14 (m, 2H), 6.96-6.94 (m, 2H), 5.07 (s, 2H), 3.53 (s, 2H),  
3.05-3.02 (m, 1H), 2.91-2.81 (m, 4H), 1.21 (t, 3H, J = 7.4); low resolution MS  
35 (ES+) m / e 382.94 (MH<sup>+</sup>); TLC R<sub>f</sub> = 0.33 (4/1, DCM/MeOH).

40 **Intermediate 63: (2S)-2-amino-3-{4-[(5-ethyl-2-(4-fluorophenyl)-1,3-oxazol-4-yl)methoxy]phenyl}propanoic acid**

Intermediate 63 was prepared as described above for the preparation of  
intermediate 52. From 962 mg of N-(Boc)-L-Tyrosine methyl ester and 720 mg  
of Intermediate 10D was prepared 1.32 g of BOC-protected intermediate  
45 methyl ester (81% yield; 1H NMR (CDCl<sub>3</sub>, 300 MHz) δ 8.07-8.03 (m, 2H), 7.19-  
30 7.13 (m, 2H), 7.08 (d, 2H, J = 8.4), 6.97 (d, 2H, J = 8.4), 4.99-4.97 (m, 3H),  
4.58-4.57 (m, 1H), 3.74 (s, 3H), 3.09-3.03 (m, 2H), 2.82 (q, 2H, J = 7.5), 1.59  
50 (s, 9H), 1.30 (t, 3H, J = 7.5); From 1.31 g of BOC-protected ester intermediate



was prepared the crude TFA salt of intermediate 63. The TFA salt was suspended into water and the pH adjusted to ~7.0 with 10% NaOH. The resulting solids were filtered and dried to yield 1.01 g of the title compound; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 300 MHz) δ 8.01-7.97 (m, 2H), 7.36 (t, 2H, J = 9.0), 7.18 (d, 2H, J = 8.4), 6.96 (d, 2H, J = 8.4), 4.97 (s, 2H), 3.53-3.51 (m, 2H), 3.09-3.04 (m, 2H), 2.84 (q, 2H, J = 7.5), 1.22 (t, 3H, J = 7.5); low resolution MS (ES+) m/e 384.87 (MH<sup>+</sup>); TLC R<sub>f</sub> = 0.44 (4/1, DCM/MeOH).

**Example 1: (2S)-2-(((Z)-1-methyl-3-oxo-3-phenyl-1-propenyl)amino)-3-(4-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]phenyl)propanoic acid**

A suspension of 3.0 g (8.2 mmol) of Intermediate 45 and 1.86 g (11.5 mmol) of benzoylacetone in 26 mL of MeOH and 6 mL of trimethylorthoformate was refluxed for 16 h with dissolution occurring sometime after 4 hrs. The solution was cooled, concentrated, and the resulting crude products were purified by silica gel chromatography eluting with 1% to 100% MeOH in CH<sub>2</sub>Cl<sub>2</sub>. Column fractions homogeneous by TLC were combined and concentrated to give 1.8 g (43% yield) of Example 1 as a tan-colored solid: TLC (EtOAc/MeOH (7:3): R<sub>f</sub> = 0.19; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 400 MHz) δ 11.40 (d, 1 H, J = 8.89), 7.88 (d, 2H, J = 7.86), 7.77 (d, 2H, J = 7.35), 7.46 (m, 3H), 7.39 (m, 3H), 7.10 (d, 2H, J = 8.55), 6.80 (d, 2H, J = 8.20), 5.78 (s, 1H), 4.14 (m, 3H), 3.10 (m, 1H), 2.87 (m, 2H), 2.77 (m, 1H), 2.34 (s, 3H), 1.71 (s, 3H); low resolution MS (ES+) m/e 511.1 (MH<sup>+</sup>); RP-HPLC (Vydac C-18, 25cm x 4.6mm; 30-100% CH<sub>3</sub>CN in H<sub>2</sub>O) with 0.1% HCO<sub>2</sub>H buffer: 30 minutes; 1 mL/min: t<sub>r</sub> = 18.14 min (t<sub>0</sub> = 1.43); Daicel AD(2) (25cm x 4.6mm; 20% IPA in Hexane) with 0.1% TFA buffer: 15 minutes; 0.8 mL/min: t<sub>r</sub> = 6.87 min (t<sub>0</sub> = 1.43), 99.9% ee.

**Example 2: (2S)-3-(4-[2-(2-(4-fluorophenyl)-5-methyl-1,3-oxazol-4-yl)ethoxy]phenyl)-2-(((Z)-1-methyl-3-oxo-3-[4-(trifluoromethyl)phenyl]-1-propenyl)amino)propanoic acid**

A mixture of 75 mg (0.2 mmol) of Intermediate 46, 45 mg (0.2 mmol) of Intermediate 24, and 57 μL (0.41 mmol) of triethylamine in 0.5 mL of trimethylorthoformate and 2.5 mL of MeOH was refluxed for 10 hrs. The

solvents were evaporated under reduced pressure and the residue purified directly by silica gel chromatography. Elution with 5%-20% MeOH in CH<sub>2</sub>Cl<sub>2</sub> gave 30 mg (26% yield) of Example 2 as a solid (foam); TLC (DCM/MeOH, 4/1): R<sub>f</sub> = 0.58; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 300 MHz) δ 11.59 (d, 1H, J = 9.8), 8.03 (m, 4H), 7.81 (d, 2H, J = 8.8), 7.38 (t, 2H, J = 9.2), 7.19 (d, 2H, J = 8.8), 6.88 (d, 2H, J = 8.8), 5.76 (s, 1H), 4.21 (t, 2H, J = 6.6), 4.17 (m, 1H), 3.20 (m, 1H), 2.96 (t, 2H, J = 6.6), 2.82 (m, 1H), 2.39 (s, 3H), 1.76 (s, 3H); low resolution MS (ES+) m/e 597.0 (MH<sup>+</sup>).

**Example 3: (2S)-3-(4-{2-[2-(4-isopropoxyphenyl)-5-methyl-1,3-oxazol-4-yl]ethoxy}phenyl)-2-(((Z)-1-methyl-3-oxo-3-[4-(trifluoromethyl)phenyl]-1-propenyl)amino)propanoic acid**

The title compound was prepared (as described above for the preparation of Example 2) from 75 mg (0.18 mmol) of Intermediate 51 and 41 mg (0.18 mmol) of Intermediate 24 to yield 48 mg (43% yield) of Example 3: TLC (DCM/MeOH, 4/1): R<sub>f</sub> = 0.42; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 400 MHz) δ 11.5 (d, 1H, J = 8.9), 7.95 (d, 2H, J = 7.2), 7.77 (d, 2H, J = 8.7), 7.72 (d, 2H, J = 8.2), 7.1 (d, 2H, J = 8.4), 6.97 (d, 2H, J = 8.9), 6.79 (d, 2H, J = 8.5), 5.58 (s, 1H), 4.65 (m, 1H), 4.11 (t, 2H, J = 6.5), 4.1 (m, 1H), 3.16 (dd, 1H, J = 13.7, 3.6), 2.84 (t, 2H, J = 6.5), 2.74 (dd, 1H, 13.7, 9.0), 2.28 (s, 3H), 1.67 (s, 3H), 1.25 (d, 6H, J = 6.0); low resolution MS (ES+) m/e 637.1 (MH<sup>+</sup>);

**Example 4: (2S)-3-(4-{2-[2-(4-methoxyphenyl)-5-methyl-1,3-oxazol-4-yl]ethoxy}phenyl)-2-(((Z)-1-methyl-3-oxo-3-[4-(trifluoromethyl)phenyl]-1-propenyl)amino)propanoic acid**

The title compound was prepared (as described above for the preparation of Example 2) from 100 mg (0.25 mmol) of Intermediate 49 and 58 mg (0.25 mmol) of Intermediate 24 to yield 66 mg (43% yield) of Example 4: TLC (DCM/MeOH, 4/1): R<sub>f</sub> = 0.48; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 400 MHz) δ 11.5 (d, 1H, J = 8.9), 7.95 (d, 2H, J = 8.2), 7.80 (d, 2H, J = 8.7), 7.72 (d, 2H, J = 8.4), 7.1 (d, 2H, J = 8.4), 7.0 (d, 2H, J = 8.9), 6.79 (d, 2H, J = 8.5), 5.58 (s, 1H), 4.11 (t, 2H, J = 6.5), 4.1 (m, 1H), 3.78 (s, 3H), 3.15 (dd, 1H, J = 13.6, 3.6), 2.84 (t, 2H,

$J = 66$ ), 2.75 (dd, 1H, 13.7, 9.0), 2.28 (s, 3H), 1.67 (s, 3H); low resolution MS (ES+)  $m/e$  609.0 (MH<sup>+</sup>);

**Example 5: (2S)-2-([(Z)-1-ethyl-3-oxo-3-phenyl-1-propenyl]amino)-3-[4-(2-{5-methyl-2-[4-(trifluoromethyl)phenyl]-1,3-oxazol-4-yl}ethoxy)phenyl]propanoic acid**

The title compound was prepared (as described above for the preparation of Example 2) from 40 mg (0.092 mmol) of Intermediate 50 and 17 mg (0.092 mmol) of Intermediate 16 to yield 21 mg of Example 5: TLC (DCM/MeOH, 4/1):  $R_f = 0.50$ ; <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>, 400 MHz)  $\delta$  11.59 (d, 1H,  $J = 9.8$ ), 8.09 (d, 2H,  $J = 9.0$ ), 7.85 (d, 2H,  $J = 8.8$ ), 7.78 (m, 2H), 7.39 (m, 3H), 7.12 (d, 2H,  $J = 8.0$ ), 6.80 (d, 2H,  $J = 8.0$ ), 5.54 (s, 1H), 4.15 (t, 2H,  $J = 6.6$ ), 4.17 (m, 1H), 3.11 (m, 1H), 2.90 (t, 2H,  $J = 6.6$ ), 2.72 (m, 1H), 2.33 (s, 3H), 1.98 (m, 2H), 0.9 (t, 3H,  $J = 7.5$ ); low resolution MS (ES+)  $m/e$  593.1 (MH<sup>+</sup>).

**Example 6: (2S)-2-([(Z)-1-ethyl-3-(4-fluorophenyl)-3-oxo-1-propenyl]amino)-3-[4-(2-{2-(4-methoxyphenyl)-5-methyl-1,3-oxazol-4-yl}ethoxy)phenyl]propanoic acid**

The title compound was prepared (as described above for the preparation of Example 2) from 100 mg (0.25 mmol) of Intermediate 49 and 62 mg (0.32 mmol) of Intermediate 32 to yield 90 mg of Example 6: TLC (DCM/MeOH, 4/1):  $R_f = 0.46$ ; <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>, 400 MHz)  $\delta$  11.45 (d, 1H,  $J = 9.6$ ), 7.84 (m, 4H), 7.19 (t, 2H,  $J = 8.8$ ), 7.10 (d, 2H,  $J = 8.4$ ), 7.01 (d, 2H,  $J = 8.8$ ), 6.78 (d, 2H,  $J = 8.4$ ), 5.51 (s, 1H), 4.11 (t, 2H,  $J = 6.8$ ), 4.03 (m, 1H), 3.11 (m, 1H), 2.84 (t, 2H,  $J = 6.8$ ), 2.71 (dd, 1H,  $J = 14.0, 9.2$ ), 2.31 (s, 3H), 1.98 (m, 2H), 0.90 (t, 3H,  $J = 7.2$ ); low resolution MS (ES-)  $m/e$  571.2 (M-H).

**Example 7: (2S)-2-([(Z)-1-methyl-3-oxo-3-phenyl-1-propenyl]amino)-3-[4-(2-{5-methyl-2-[4-(trifluoromethyl)phenyl]-1,3-oxazol-4-yl}ethoxy)phenyl]propanoic acid**

The title compound was prepared (as described above for the preparation of Example 2) from 40 mg (0.092 mmol) of Intermediate 50 and

15 mg (0.092 mmol) of benzoylacetone to yield 32 mg of Example 7: TLC (DCM/MeOH, 4/1):  $R_f = 0.48$ ;  $^1\text{H NMR}$  (DMSO- $d_6$ , 300 MHz)  $\delta$  11.49 (d, 1H,  $J = 9.8$ ), 8.17 (d, 2H,  $J = 8.9$ ), 7.9 (d, 2H,  $J = 8.9$ ), 7.85 (m, 2H), 7.44 (m, 3H), 7.19 (d, 2H,  $J = 8.8$ ), 6.89 (d, 2H,  $J = 8.8$ ), 5.60 (s, 1H), 4.23 (t, 2H,  $J = 6.0$ ), 4.16 (m, 1H), 3.21 (m, 2H), 2.99 (t, 2H,  $J = 6.0$ ), 2.82 (dd, 1H,  $J = 14.2, 9.8$ ), 2.42 (s, 3H), 1.78 (s, 3H); low resolution MS (ES+)  $m/e$  579.0 ( $\text{MH}^+$ ).

**Example 8: (2S)-2-([(Z)-1-ethyl-3-oxo-3-phenyl-1-propenyl]amino)-3-(4-{2-[2-(4-fluorophenyl)-5-methyl-1,3-oxazol-4-yl]ethoxy}phenyl)propanoic acid**

The title compound was prepared (as described above for the preparation of Example 2) from 75 mg (0.195 mmol) of Intermediate 46 and 34 mg (0.2 mmol) of Intermediate 16 to yield 32 mg of Example 8: TLC (DCM/MeOH, 4/1):  $R_f = 0.63$ ;  $^1\text{H NMR}$  (DMSO- $d_6$ , 300 MHz)  $\delta$  11.57 (d, 1H,  $J = 9.6$ ), 7.97 (m, 2H), 7.86 (d, 1H,  $J = 7.2$ ), 7.42 (m, 5H), 7.18 (d, 2H, 8.4), 6.87 (d, 2H,  $J = 8.4$ ), 5.61 (s, 1H), 4.20 (t, 2H,  $J = 6.6$ ), 4.14 (m, 1H), 3.22 (m, 2H), 2.94 (t, 2H,  $J = 6.6$ ), 2.83 (dd, 1H,  $J = 13.8, 9.3$ ); low resolution MS (ES+)  $m/e$  543 ( $\text{MH}^+$ ).

**Example 9: (2S)-2-([(Z)-1-ethyl-3-(4-fluorophenyl)-3-oxo-1-propenyl]amino)-3-(4-{2-[2-(4-fluorophenyl)-5-methyl-1,3-oxazol-4-yl]ethoxy}phenyl)propanoic acid**

The title compound was prepared (as described above for the preparation of Example 2) from 500 mg (1.56 mmol) of Intermediate 46 and 315 mg (1.56 mmol) of Intermediate 32 to yield 412 mg of Example 9: TLC (DCM/MeOH, 4/1):  $R_f = 0.53$ ;  $^1\text{H NMR}$  (DMSO- $d_6$ , 300 MHz)  $\delta$  11.53 (d, 1H,  $J = 9.3$ ), 7.99 (m, 2H), 7.95 (m, 2H), 7.39 (t, 2H,  $J = 8.7$ ), 7.26 (t, 2H,  $J = 8.7$ ), 7.17 (d, 2H,  $J = 8.1$ ), 6.86 (d, 2H,  $J = 8.1$ ), 5.60 (s, 1H), 4.20 (t, 2H,  $J = 6.6$ ), 4.12 (m, 1H), 3.21 (m, 2H), 2.94 (t, 2H,  $J = 6.6$ ), 2.78 (dd, 1H,  $J = 13.8, 8.4$ ), 2.38 (s, 3H), 2.05 (m, 2H), 0.96 (t, 3H,  $J = 7.5$ ); low resolution MS (ES+)  $m/e$  561.4 ( $\text{MH}^+$ ).

**Example 10: (2S)-2-((Z)-1-methyl-3-oxo-3-(4-(trifluoromethyl)phenyl)-1-propenyl)amino-3-(4-(2-(5-methyl-2-phenyl-1,3-thiazol-4-yl)ethoxy)phenyl)propanoic acid**

The title compound was prepared (as described above for the preparation of Example 2) from 290 mg (0.76 mmol) of Intermediate 47 and 150 mg (0.76 mmol) of Intermediate 24 to yield 140 mg (31% yield) of Example 10: TLC (DCM/MeOH, 4/1):  $R_f = 0.47$ ;  $^1\text{H}$  NMR (DMSO- $d_6$ , 300 MHz)  $\delta$  11.52 (d, 1H,  $J = 8.9$ ), 7.96 (d, 2H,  $J = 8.1$ ), 7.80 (m, 2H), 7.72 (d, 2H,  $J = 8.4$ ), 7.42 (m, 3H), 7.10 (d, 2H,  $J = 8.4$ ), 6.80 (d, 2H,  $J = 8.4$ ), 5.61 (s, 1H), 4.20 (t, 2H,  $J = 6.6$ ), 4.15 (m, 1H), 3.17 (m, 1H), 3.05 (t, 2H,  $J = 6.5$ ), 2.77 (dd, 1H,  $J = 13.6, 8.6$ ), 2.41 (s, 3H), 1.69 (s, 3H); low resolution MS (ES+)  $m/e$  595.0 ( $\text{MH}^+$ ).

**Example 11: (2S)-2-((Z)-3-(4-fluorophenyl)-1-methyl-3-oxo-1-propenyl)amino-3-(4-(2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy)phenyl)propanoic acid**

A solution of 100 mg (0.27 mmol) of Intermediate 45 and 49 mg (0.27 mmol) of Intermediate 26 in 2.5 mL MeOH and 0.5 mL trimethyl orthoformate was refluxed at 80 °C for 16h. The reaction solution was concentrated and purified by flash silica gel chromatography using EtOAc / MeOH 7 / 3 as eluent to afford 83 mg of Example 11: TLC (EtOAc/MeOH, 7/3):  $R_f = 0.28$ ;  $^1\text{H}$  NMR (DMSO- $d_6$ )  $\delta$  11.38 (d, 1 H,  $J = 7.00$ ), 7.84 (m, 4H), 7.45 (m, 3H), 7.16 (t, 2H,  $J = 8.72$ ), 7.09 (d, 2H,  $J = 7.69$ ), 6.77 (d, 2H,  $J = 7.69$ ), 5.49 (s, 1H), 4.11 (m, 2H), 4.01 (m, br, 1H), 3.13 (m, 1H), 2.86 (m, 2H), 2.72 (s, br, 1H), 2.30 (s, 3H), 1.62 (s, 3H); low resolution MS (ES)  $m/e$  529 ( $\text{MH}^+$ ); RP-HPLC (Vydac C-18, 25cm x 4.6mm; 30-100%  $\text{CH}_3\text{CN}$  in  $\text{H}_2\text{O}$ ) with 0.1%  $\text{HCO}_2\text{H}$  buffer: 30 minutes; 1 mL/min:  $t_r = 20.10$  min ( $t_0 = 1.53$ ).

**Example 12: (2S)-2-((Z)-1-methyl-3-oxo-3-(2,3,4-trifluorophenyl)-1-propenyl)amino-3-(4-(2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy)phenyl)propanoic acid**

A mixture of 680 mg (1.85 mmol) of Intermediate 45, 367 mg (1.85 mmol) of Intermediate 36, and 0.67 mL (3.7 mmol) of DIEA in 15 mL of MeOH was refluxed overnight (14 h). The solvent was evaporated under reduced pressure and the residue purified directly by silica gel chromatography.

Elution with 5%-20% MeOH in CH<sub>2</sub>Cl<sub>2</sub> gave 612 mg (58% yield) of the title compound as a solid: TLC (DCM/MeOH, 4/1):  $R_f$  = 0.50; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 300 MHz) δ 11.35 (d, 1H,  $J$  = 9.0), 7.87 (m, 2H), 7.48 (m, 4H), 7.3 (q, 1H,  $J$  = 5.4), 7.10 (d, 2H,  $J$  = 8.7), 6.78 (d, 2H,  $J$  = 8.7), 5.25 (s, 1H), 4.18 (m, 3H), 3.19 (m, 2H), 2.86 (t, 2H,  $J$  = 6.3), 2.74 (dd, 1H,  $J$  = 13.8, 9.0), 2.32 (s, 3H), 1.64 (s, 3H); low resolution MS (ES+)  $m/e$  564.9 (MH<sup>+</sup>);

**Example 13: (2S)-2-(((Z)-1-methyl-3-(4-nitrophenyl)-3-oxo-1-propenyl)amino)-3-{4-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]phenyl}propanoic acid**

The title compound was prepared (as described above for the preparation of Example 12) from 120 mg (0.33 mmol) of Intermediate 45 and 62 mg (0.33 mmol) of Intermediate 41 to yield 105 mg (58% yield) of Example 13: TLC (DCM/MeOH, 4:1):  $R_f$  = 0.60; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 300 MHz) δ 11.66 (d, 1H,  $J$  = 9.0), 8.29 (d, 2H,  $J$  = 8.4), 8.08 (d, 2H,  $J$  = 8.4), 7.97 (m, 2H), 7.56 (m, 3H), 7.19 (d, 2H,  $J$  = 8.0), 6.88 (d, 2H,  $J$  = 8.0), 5.72 (s, 1H), 4.18 (m, 3H), 3.23 (m, 1H), 2.96 (t, 2H,  $J$  = 6.2), 2.85 (dd, 1H,  $J$  = 13.4, 9.6), 2.39 (s, 3H), 1.79 (s, 3H); low resolution MS (ES+)  $m/e$  556.4 (MH<sup>+</sup>).

**Example 14: (2S)-2-(((Z)-1-methyl-3-oxo-3-[4-(trifluoromethyl)phenyl]-1-propenyl)amino)-3-{4-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]phenyl}propanoic acid**

A solution of 676 mg (1.85 mmol) of Intermediate 45 and 510 mg (2.22 mmol) of Intermediate 24 in 30 mL MeOH and 6 mL trimethyl orthoformate with 4 Å sieves (300 mg) was refluxed at 80°C for 16h. The reaction solution was concentrated and purified by flash silica gel column chromatography using DCM / MeOH 15 / 1 followed by DCM / MeOH 10 / 1 as eluent to afford 329 mg of Example 14: TLC (DCM/MeOH (9:1):  $R_f$  = 0.26; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 300 MHz) δ 11.59 (d, 1H,  $J$  = 9.0), 8.03 (d, 2H,  $J$  = 8.1), 7.95 (m, 2H), 7.80

(d, 2H,  $J=8.1$ ), 7.56 (m, 3H), 7.19 (d, 2H,  $J=8.4$ ), 6.87 (d, 2H,  $J=8.4$ ), 5.66 (s, 1H), 4.18 (m, 3H), 3.23 (m, 2H), 2.95 (t, 2H,  $J=6.6$ ), 2.82 (dd, 1H,  $J=13.8$ , 9.0), 2.39 (s, 3H), 1.75 (s, 3H); low resolution MS (ES+)  $m/e$  579 ( $MH^+$ ); RP-HPLC (Vydac C-18, 25cm x 4.6mm; 30-100%  $CH_3CN$  in  $H_2O$ ) with 0.1%  $HCO_2H$  buffer: 30 minutes; 1 mL/min:  $t_r=21.47$  min ( $t_0=1.53$ ).

**Example 15: (2S)-2-[(Z)-1-ethyl-3-(4-fluorophenyl)-3-oxo-1-propenyl]amino-3-(4-[2-[2-(4-isopropoxyphenyl)-5-methyl-1,3-oxazol-4-yl]ethoxy]phenyl)propanoic acid**

The title compound was prepared (as described above for the preparation of Example 2) from 75 mg (0.177 mmol) of Intermediate 51 and 39 mg (0.194 mmol) of Intermediate 32 to yield 70 mg of Example 15: TLC (DCM/MeOH (4:1):  $R_f=0.55$ ;  $^1H$  NMR (DMSO- $d_6$ , 300 MHz)  $\delta$  11.53 (d, 1H,  $J=9.6$ ), 7.89 (dd, 2H,  $J=8.8$ , 5.7), 7.85 (d, 2H,  $J=8.8$ ), 7.26 (t, 2H,  $J=8.8$ ), 7.18 (d, 2H,  $J=8.5$ ), 7.05 (d, 2H,  $J=8.8$ ), 6.86 (d, 2H,  $J=8.5$ ), 5.60 (s, 1H), 4.73 (hept, 1H,  $J=6.0$ ), 4.19 (t, 2H,  $J=6.7$ ), 4.13 (br s, 1H), 3.19 (m, 1H), 2.92 (t, 2H,  $J=6.6$ ), 2.80 (dd, 1H,  $J=14.0$ , 9.2), 2.36 (s, 3H), 2.07 (m, 2H), 1.34 (d, 6H,  $J=6.0$ ), 0.99 (t, 3H,  $J=7.5$ ); low resolution MS (ES+)  $m/e$  601.1 ( $MH^+$ ).

**Example 16: (2S)-2-[(Z)-1-methyl-3-oxo-3-(2,4,5-trifluorophenyl)-1-propenyl]amino-3-(4-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]phenyl)propanoic acid**

The title compound was prepared (as described above for the preparation of Example 12) from 677 mg (1.85 mmol) of Intermediate 45 and 366 mg (1.85 mmol) of Intermediate 38 to yield 482 mg (46% yield) of Example 16. TLC (DCM/MeOH (4:1):  $R_f=0.55$ ;  $^1H$  NMR (DMSO- $d_6$ , 300 MHz)  $\delta$  11.39 (d, 1H,  $J=8.9$ ), 7.86 (m, 2H), 7.69-7.59 (m, 1H), 7.54-7.40 (m, 4H), 7.10 (d, 2H,  $J=8.5$ ), 6.79 (d, 2H,  $J=8.5$ ), 5.3 (s, 1H), 4.12 (t, 2H,  $J=6.5$ ), 4.1 (m, 1H), 3.16 (m, 1H), 2.86 (t, 2H,  $J=6.5$ ), 2.76 (dd, 1H,  $J=13.7$ , 8.9), 2.3 (s, 3H), 1.64 (s, 3H); low resolution MS (ES+)  $m/e$  565.1 ( $MH^+$ ).

**Example 17: (2S)-2-(((Z)-1-ethyl-3-oxo-3-phenyl-1-propenyl)amino)-3-(4-(2-(2-(4-isopropoxyphenyl)-5-methyl-1,3-oxazol-4-yl)ethoxy)phenyl)propanoic acid**

The title compound was prepared (as described above for the preparation of Example 2) from 75 mg (0.177 mmol) of Intermediate 51 and 34 mg (0.194 mmol) of Intermediate 16 to yield 30 mg of Example 17: TLC (DCM/MeOH (4:1):  $R_f$  = 0.63;  $^1\text{H}$  NMR (DMSO- $d_6$ , 400 MHz)  $\delta$  11.49 (d, 1H,  $J$  = 9.6), 7.77 (m, 4H), 7.37 (m, 3H), 7.10 (d, 2H,  $J$  = 8.8), 6.97 (d, 2H,  $J$  = 9.2), 6.79 (d, 2H,  $J$  = 9.2), 5.54 (s, 1H), 4.66 (septuplet, 1H,  $J$  = 6.0), 4.12 (t, 2H,  $J$  = 6.8), 4.04 (m, 1H), 3.35 (m, 1H), 3.11 (m, 1H), 2.84 (t, 2H,  $J$  = 6.8), 2.72 (dd, 1H,  $J$  = 13.9, 9.2), 2.28 (s, 3H), 1.99 (m, 2H), 1.26 (d, 6H,  $J$  = 6.0), 0.91 (t, 3H,  $J$  = 7.6); low resolution MS (ES+)  $m/e$  583.1 ( $\text{MH}^+$ ).

**Example 18: (2S)-2-(((Z)-1-ethyl-3-oxo-3-(4-(trifluoromethyl)phenyl)-1-propenyl)amino)-3-(4-(2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy)phenyl)propanoic acid**

The title compound was prepared (as described above for the preparation of Example 2) from 197 mg (0.41 mmol) of Intermediate 45 (as the TFA salt) and 100 mg (0.41 mmol) of Intermediate 25 to yield 110 mg (45% yield) of Example 18: TLC (DCM/MeOH (4:1):  $R_f$  = 0.60;  $^1\text{H}$  NMR (DMSO- $d_6$ , 400 MHz)  $\delta$  11.45 (d, 1H,  $J$  = 9.2), 7.85 (m, 4H), 7.47 (m, 3H), 7.17 (t, 2H,  $J$  = 8.8), 7.09 (d, 2H,  $J$  = 8.4), 6.78 (d, 2H,  $J$  = 8.4), 5.51 (s, 1H), 4.12 (t, 2H,  $J$  = 6.4), 4.02 (m, 1H), 3.10 (m, 2H), 2.86 (t, 2H,  $J$  = 6.4), 2.71 (dd, 1H,  $J$  = 13.9, 9.2), 2.31 (s, 3H), 1.98 (s, 3H), 0.90 (t, 3H,  $J$  = 7.6); low resolution MS (ES+)  $m/e$  593 ( $\text{MH}^+$ ).

**Example 19: (2S)-2-(((Z)-1-methyl-3-(4-methylphenyl)-3-oxo-1-propenyl)amino)-3-(4-(2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy)phenyl)propanoic acid**

The title compound was prepared (as described above for the preparation of Example 2) from 790 mg (1.64 mmol) of Intermediate 45 (as the TFA salt) and 0.29 g (1.64 mmol) of Intermediate 27 (with 660  $\mu\text{L}$  of DCA replacing  $\text{Et}_3\text{N}$ ) to yield crude product contaminated with DCA). A second



silica gel chromatography column eluting with 8:1 DCM:MeOH gave 70 mg of Example 19 as a beige-colored solid: TLC (DCM/MeOH (4:1):  $R_f$  = 0.58;  $^1\text{H}$  NMR (DMSO- $d_6$ , 400 MHz)  $\delta$  11.43 (d, 1H,  $J$  = 9.0), 7.96 (m, 2H), 7.74 (d, 2H,  $J$  = 8.1), 7.54 (m, 3H), 7.24 (d, 2H,  $J$  = 8.1), 7.18 (d, 2H,  $J$  = 8.4), 6.87 (d, 2H,  $J$  = 8.4), 5.57 (s, 1H), 4.21 (t, 2H,  $J$  = 6.6), 4.08 (m, 1H), 3.23 (m, 2H), 2.95 (t, 2H,  $J$  = 6.6), 2.80 (dd, 1H,  $J$  = 13.8, 9.0), 2.39 (s, 3H), 2.37 (s, 3H), 1.71 (s, 3H); low resolution MS (ES+)  $m/e$  525.2 ( $\text{MH}^+$ ).

**Example 20: (2S)-2-([(Z)-1-ethyl-3-oxo-3-phenyl-1-propenyl]amino)-3-(4-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]phenyl)propanoic acid**

The title compound was prepared (as described above for the preparation of Example 2) from 1.03 g (2.1 mmol) of Intermediate 45 (as the TFA salt) and 740 mg (4.2 mmol) of Intermediate 16 to yield 500 mg of Example 20 as a white solid: TLC (EtOAc/MeOH, 7/3):  $R_f$  = 0.35;  $^1\text{H}$  NMR (DMSO- $d_6$ , 400 MHz)  $\delta$  11.55 (d, 1H,  $J$  = 9.6), 7.93 (d, 2H,  $J$  = 7.8), 7.84 (d, 2H,  $J$  = 7.8), 7.57 (m, 3H), 7.52 (m, 3H), 7.18 (d, 2H,  $J$  = 8.4), 6.87 (d, 2H,  $J$  = 8.4), 5.58 (s, 1H), 4.21 (t, 2H,  $J$  = 6.6), 4.16 (m, 1H), 3.21 (m, 1H), 2.95 (t, 2H,  $J$  = 6.6), 2.80 (dd, 1H,  $J$  = 13.5, 9.0), 2.42 (s, 3H), 2.08 (m, 2H), 1.04 (t, 3H,  $J$  = 7.8); low resolution MS (ES+)  $m/e$  525 ( $\text{MH}^+$ ).

**Example 21: (2S)-3-(4-(2-[2-(4-fluorophenyl)-5-methyl-1,3-thiazol-4-yl]ethoxy)phenyl)-2-([(Z)-3-oxo-3-phenyl-1-(trifluoromethyl)-1-propenyl]amino)propanoic acid**

A solution of 120 mg (0.31 mmol) of Intermediate 48, 62 mg (0.31 mmol) of Intermediate 44, and 110  $\mu\text{L}$  (2 equivalents) of DIEA in MeOH (3.5 mL) was refluxed for 10 hrs. The solvents were removed under reduced pressure and the crude products were purified by silica gel chromatography (gradient, 15/1 to 10/1  $\text{CH}_2\text{Cl}_2/\text{MeOH}$ ) to give 78 mg (46% yield) of Example 21 as a solid: TLC (DCM/MeOH (4:1):  $R_f$  = 0.62;  $^1\text{H}$  NMR (DMSO- $d_6$ , 400 MHz)  $\delta$  10.9 (br s, 1H), 7.9-7.83 (m, 4H), 7.6-7.43 (m, 3H), 7.27 (t, 2H,  $J$  = 8.8), 7.05 (d, 2H,  $J$  = 8.4), 6.78 (d, 2H,  $J$  = 8.4), 6.13 (s, 1H), 4.2 (t, 2H,  $J$

=6.6), 4.08 (br s, 1H), 3.1 (m, 1H), 3.08 (t, 2H,  $J=6.7$ ), 2.86 (dd, 1H,  $J=13.6$ , 7.7), 2.37 (s, 3H); low resolution MS (ES+)  $m/e$  599.0 (MH<sup>+</sup>).

**Example 22: (2S)-2-(((Z)-1-ethyl-3-oxo-3-(4-(trifluoromethyl)phenyl)-1-propenyl)amino)-3-(4-(2-(2-(4-fluorophenyl)-5-methyl-1,3-thiazol-4-yl)ethoxy)phenyl)propanoic acid**

The title compound was prepared (as described above for the preparation of Example 2) from 75 mg (0.19 mmol) of Intermediate 48 and 46 mg (0.19 mmol) of Intermediate 25 to yield 26 mg of Example 22: TLC (DCM/MeOH (4:1):  $R_f = 0.58$ ; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 400 MHz)  $\delta$  11.59 (d, 1H,  $J=9.3$ ), 7.97 (d, 2H,  $J=8.1$ ), 7.85 (m, 2H), 7.72 (d, 2H,  $J=8.4$ ), 7.29 (m, 2H), 7.10 (t, 2H,  $J=8.4$ ), 6.79 (d, 2H,  $J=8.4$ ), 5.60 (s, 1H), 4.20 (t, 2H,  $J=6.6$ ), 4.12 (m, 1H), 3.16 (m, 1H), 3.06 (t, 2H,  $J=6.6$ ), 2.75 (dd, 1H,  $J=13.8$ , 9.0), 2.38 (s, 3H), 2.02 (m, 2H), 0.92 (t, 3H,  $J=7.5$ ); low resolution MS (ES+)  $m/e$  627.0 (MH<sup>+</sup>).

**Example 23: (2S)-2-(((Z)-1-butyl-3-oxo-3-phenyl-1-propenyl)amino)-3-(4-(2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy)phenyl)propanoic acid**

The title compound was prepared (as described above for the preparation of Example 2) from 1.0 g (2.05 mmol) of Intermediate 45 (as the TFA salt) and 0.86 g (4.2 mmol) of Intermediate 18 to yield 290 mg (25% yield) of Example 23 as a solid: TLC (EtOAc/MeOH, 7/3):  $R_f = 0.55$ ; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 400 MHz)  $\delta$  11.59 (d, 1H,  $J=9.7$ ), 7.86 (m, 2H), 7.76 (m, 2H), 7.48-7.34 (m, 6H), 7.10 (d, 2H,  $J=8.5$ ), 7.10 (t, 2H,  $J=8.4$ ), 6.8 (d, 2H,  $J=8.6$ ), 5.52 (s, 1H), 4.12 (t, 2H,  $J=6.5$ ), 4.02 (m, 1H), 3.16 (m, 1H), 2.86 (t, 2H,  $J=6.5$ ), 2.75 (dd, 1H,  $J=13.5$ , 9.5), 2.31 (s, 3H), 1.9 (m, 2H), 1.3-1.1 (m, 4H), 0.73 (t, 3H,  $J=6.9$ ); low resolution MS (ES+)  $m/e$  553.2 (MH<sup>+</sup>).

**Example 24: (2S)-2-(((Z)-3-(4-chlorophenyl)-1-methyl-3-oxo-1-propenyl)amino)-3-(4-(2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy)phenyl)propanoic acid**

The title compound was prepared (as described above for the preparation of Example 2) from 100 mg (0.27 mmol) of Intermediate 45 and

5 54 mg (0.27 mmol) of Intermediate 28 to yield 76 mg of Example 24: TLC  
(DCM/MeOH (4:1):  $R_f$  = 0.52;  $^1\text{H}$  NMR (DMSO- $d_6$ , 300 MHz)  $\delta$  11.52 (d, 1H,  $J$   
=9.6), 7.97 (m, 2H), 7.87 (d, 2H,  $J$  =8.6), 7.55 (m, 5H), 7.18 (d, 2H,  $J$  =8.5),  
10 6.86 (d, 2H,  $J$  =8.5), 5.59 (s, 1H), 4.21 (t, 2H,  $J$  =6.1), 4.09 (m, 1H), 3.22 (m,  
5 1H), 2.96 (t, 2H,  $J$  =6.1), 2.78 (dd, 1H,  $J$  =13.9, 9.4), 2.39 (s, 3H), 1.73 (s, 3H).

15 **Example 25: (2S)-2-(((Z)-1-methyl-3-(3-nitrophenyl)-3-oxo-1-  
propenyl)amino)-3-(4-[2-(5-methyl-2-phenyl-1,3-oxazol-4-  
yl)ethoxy]phenyl)propanoic acid**

19 The title compound was prepared (as described above for the  
20 preparation of Example 12) from 120 mg (0.25 mmol) of Intermediate 45 and  
48 mg (0.25 mmol) of Intermediate 42 to yield 67 mg of Example 25: TLC  
(DCM/MeOH (4:1):  $R_f$  = 0.52;  $^1\text{H}$  NMR (DMSO- $d_6$ , 400 MHz)  $\delta$  11.52 (d, 1H,  $J$   
=9.1), 8.52 (s, 1H), 8.24 (d, 1H,  $J$  =8.2), 8.21 (d, 2H,  $J$  =7.7), 7.87 (d, 2H,  $J$   
25 =7.7), 7.66 (t, 1H,  $J$  =7.9), 7.45 (m, 3H), 7.10 (d, 2H,  $J$  =8.2), 6.79 (d, 2H,  $J$   
15 =8.3), 5.74 (s, 1H), 4.13 (t, 2H,  $J$  =6.6), 4.07 (br s, 1H), 3.15 (m, 1H), 2.87 (t,  
2H,  $J$  =6.4), 2.75 (m, 1H), 2.31 (s, 3H), 1.68 (s, 3H); low resolution MS  
(ES $^+$ ) $m/e$  556.3 (MH $^+$ ).

20 **Example 26: (2S)-2-(((Z)-3-[2-fluoro-3-(trifluoromethyl)phenyl]-1-methyl-  
3-oxo-1-propenyl)amino)-3-(4-[2-(5-methyl-2-phenyl-1,3-oxazol-4-  
yl)ethoxy]phenyl)propanoic acid**

The title compound was prepared (as described above for the  
preparation of Example 12) from 190 mg (0.52 mmol) of Intermediate 45 and  
40 120 mg (0.52 mmol) of Intermediate 37 to yield 102 mg (35% yield) of  
25 Example 26:  $^1\text{H}$  NMR (DMSO- $d_6$ , 400 MHz)  $\delta$  11.36 (d, 1H,  $J$  =8.9), 7.9 (m,  
3H), 7.77 (t, 1H,  $J$  =6.9), 7.46 (m, 3H), 7.42 (m, 1H), 7.11 (d, 2H,  $J$  =8.5), 6.81  
(d, 2H,  $J$  =8.5), 5.22 (s, 1H), 4.14 (t, 2H,  $J$  =6.6), 4.1 (m, 1H), 3.13 (m, 1H),  
45 2.87 (t, 2H,  $J$  =6.6), 2.78 (dd, 1H,  $J$  =13.7, 8.5), 2.31 (s, 3H), 1.68 (s, 3H); low  
30 resolution MS (ES $^+$ ) $m/e$  597.4 (MH $^+$ ).

**Example 27: (2S)-2-(((Z)-3-(4-isopropoxyphenyl)-1-methyl-3-oxo-1-propenyl)amino)-3-(4-{2-[5-methyl-2-phenyl-1,3-oxazol-4-yl]ethoxy}phenyl)propanoic acid**

The title compound was prepared (as described above for the preparation of Example 2) from 435 mg (0.91 mmol) of Intermediate 45 (as the TFA salt) and 200 mg (0.91 mmol) of Intermediate 29 to yield 125 mg of Example 27 as a beige glassy solid: TLC (DCM/MeOH (4:1):  $R_f$  = 0.50;  $^1\text{H}$  NMR (DMSO- $d_6$ , 400 MHz)  $\delta$  11.29 (d, 1H,  $J$  = 8.8), 7.88 (m, 2H), 7.70 (d, 2H,  $J$  = 8.8), 7.48 (m, 3H), 7.09 (d, 2H,  $J$  = 8.0), 6.85 (d, 2H,  $J$  = 8.8), 6.78 (d, 2H,  $J$  = 8.0), 5.46 (s, 1H), 4.62 (septuplet, 1H,  $J$  = 6.4), 4.12 (t, 2H,  $J$  = 6.4), 4.0 (m, 1H), 3.14 (m, 2H), 2.87 (t, 2H,  $J$  = 6.4), 2.70 (dd, 1H,  $J$  = 13.2, 9.6), 2.31 (s, 3H), 1.61 (s, 3H), 1.21 (d, 6H,  $J$  = 6.4); low resolution MS (ES+)  $m/e$  569 (MH $^+$ ).

**Example 28: (2S)-2-(((Z)-3-(2-chlorophenyl)-1-methyl-3-oxo-1-propenyl)amino)-3-(4-{2-[5-methyl-2-phenyl-1,3-oxazol-4-yl]ethoxy}phenyl)propanoic acid**

The title compound was prepared (as described above for the preparation of Example 2) from 100 mg (0.27 mmol) of Intermediate 45 and 54 mg (0.27 mmol) of Intermediate 29 to yield 44 mg of Example 27: TLC (DCM/MeOH (4:1):  $R_f$  = 0.47;  $^1\text{H}$  NMR (DMSO- $d_6$ , 400 MHz)  $\delta$  11.12 (d, 1H,  $J$  = 9.0), 7.91 (m, 2H), 7.44 (m, 3H), 7.39 (m, 1H), 7.33 (m, 3H), 7.17 (d, 2H,  $J$  = 7.2), 6.80 (d, 2H,  $J$  = 7.2), 4.97 (s, 1H), 4.14 (m, 3H), 3.16 (m, 1H), 2.88 (m, 2H), 2.77 (m, 1H), 2.32 (s, 3H), 1.61 (s, 3H); low resolution MS (ES-)  $m/e$  545.0 (M-H).

**Example 29: (2S)-2-(((Z)-3-(2-furyl)-1-methyl-3-oxo-1-propenyl)amino)-3-(4-{2-[5-methyl-2-phenyl-1,3-oxazol-4-yl]ethoxy}phenyl)propanoic acid**

The title compound was prepared (as described above for the preparation of Example 2) from 630 mg (1.31 mmol) of Intermediate 45 (as the TFA salt) and 200 mg (1.31 mmol) of 1-(2-furyl)-1,3-butanedione (purchased from Acros Organics) to yield 230 mg (35% yield) of Example 29 as a light brown solid: TLC (DCM/MeOH (4:1):  $R_f$  = 0.62;  $^1\text{H}$  NMR (DMSO- $d_6$ , 300

5 MHz)  $\delta$  11.07 (d, 1H,  $J$  = 8.7), 7.88 (m, 2H), 7.69 (d, 1H,  $J$  = 1.0), 7.48 (m, 3H),  
7.09 (d, 2H,  $J$  = 8.5), 6.86 (m, 3H), 6.52 (d, 1H,  $J$  = 1.5), 5.33 (s, 1H), 4.13 (t,  
10 2H,  $J$  = 6.6), 4.03 (m, 1H), 3.14 (m, 1H), 2.87 (t, 2H,  $J$  = 6.6), 2.70 (dd, 1H,  $J$   
= 13.6, 9.0), 2.34 (s, 3H), 1.62 (s, 3H); low resolution MS (ES+)  $m/e$  501.2  
5 (MH<sup>+</sup>).

15 **Example 30: (2S)-2-(((Z)-1-methyl-3-oxo-3-(2-pyrazinyl)-1-propenyl)amino)-3-(4-{2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy}phenyl)propanoic acid**

10 The title compound was prepared (as described above for the  
preparation of Example 2) from 292 mg (0.61 mmol) of Intermediate 45 (as  
20 the TFA salt) and 100 mg (0.61 mmol) of Intermediate 35 to yield 135 mg  
(43%) of Example 30: TLC (DCM/MeOH (4:1):  $R_f$  = 0.22; <sup>1</sup>H NMR (DMSO- $d_6$ ,  
400 MHz)  $\delta$  11.58 (d, 2H,  $J$  = 8.8), 9.08 (s, 1H), 8.68 (s, 1H), 8.60 (s, 1H), 7.87  
25 (d, 2H,  $J$  = 7.6), 7.45 (m, 3H), 7.11 (d, 2H,  $J$  = 8.0), 6.81 (d, 2H,  $J$  = 8.0), 6.04 (s,  
15 1H), 4.13 (m, 3H), 3.17 (m, 1H), 2.87 (t, 2H,  $J$  = 6.4), 2.78 (dd, 1H,  $J$  = 13.2,  
9.2), 2.31 (s, 3H), 1.71 (s, 3H); low resolution MS (ES-)  $m/e$  511.1 (M-H).

30 **Example 31: (2S)-2-(((Z)-3-(2,4-difluorophenyl)-1-methyl-3-oxo-1-propenyl)amino)-3-(4-{2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy}phenyl)propanoic acid**

35 The title compound was prepared (as described above for the  
preparation of Example 12) from 120 mg (0.33 mmol) of Intermediate 45 and  
59 mg (0.33 mmol) of Intermediate 39 to yield 63 mg (35% yield) of Example  
25 31: TLC (DCM/MeOH (4:1):  $R_f$  = 0.72; <sup>1</sup>H NMR (DMSO- $d_6$ , 400 MHz)  $\delta$  11.37  
40 (d, 1H,  $J$  = 9.4), 7.87 (m, 2H), 7.72 (q, 1H,  $J$  = 6.4), 7.43 (m, 3H), 7.14 (m, 4H),  
6.79 (d, 2H,  $J$  = 8.8), 5.25 (s, 1H), 4.13 (m, 3H), 3.18 (m, 1H), 2.85 (t, 2H,  $J$   
= 6.0), 2.77 (dd, 1H,  $J$  = 14.0, 9.8), 2.29 (s, 3H), 1.62 (s, 3H); low resolution  
45 MS (ES-)  $m/e$  545.1 (M-H).

30 **Example 32: (2S)-2-(((Z)-1-methyl-3-oxo-3-(1,3-thiazol-2-yl)-1-propenyl)amino)-3-(4-{2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy}phenyl)propanoic acid**

5 The title compound was prepared (as described above for the preparation of Example 12) from 200 mg (0.43 mmol) of Intermediate 45 (as the TFA salt) and 65 mg (0.43 mmol) of Intermediate 43 to yield 202 mg of crude product. This material was re-columned on silica gel eluting with  
10 DCM/MeOH (6/1 to 4/1 gradient) to yield 22 mg of Example 32. : TLC (DCM/MeOH, 4/1):  $R_f = 0.42$ ;  $^1\text{H NMR}$  (DMSO- $d_6$ , 300 MHz)  $\delta$  11.25 (d, 1H,  $J = 9.0$ ), 7.94 (m, 4H), 7.54 (m, 3H), 7.17 (d, 2H,  $J = 8.4$ ), 6.88 (d, 2H,  $J = 8.4$ ), 5.84 (s, 1H), 4.22 (t, 2H,  $J = 6.6$ ), 4.18 (m, 1H), 3.1 (dd, 1H,  $J = 14.0, 4.0$ ), 2.96 (t, 2H,  $J = 6.6$ ), 2.83 (dd, 1H,  $J = 13.8, 9.0$ ), 2.43 (s, 3H), 1.76 (s, 3H); low  
15 resolution MS (ES+)  $m/e$  518.1 ( $\text{MH}^+$ ).  
20

**Example 33: (2S)-2-(((Z)-1-methyl-3-oxo-3-(3-thienyl)-1-propenyl)amino)-3-(4-(2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy)phenyl)propanoic acid**

25 The title compound was prepared (as described above for the preparation of Example 2) from 428 mg (0.89 mmol) of Intermediate 45 (as the TFA salt) and 150 mg (0.89 mmol) of Intermediate 31 to yield 210 mg of Example 33: TLC (DCM/MeOH, 4/1):  $R_f = 0.60$ ;  $^1\text{H NMR}$  (DMSO- $d_6$ , 400 MHz)  $\delta$  11.8 (d, 1H,  $J = 9.2$ ), 7.88 (m, 3H), 7.42 (m, 5H), 7.08 (d, 2H,  $J = 8.4$ ), 6.78 (d, 2H,  $J = 8.4$ ), 5.39 (s, 1H), 4.12 (t, 2H,  $J = 6.4$ ), 4.01 (m, 1H), 3.23 (m, 1H),  
30 3.09 (m, 1H), 2.86 (t, 2H,  $J = 6.4$ ), 2.69 (dd, 1H,  $J = 13.6, 9.2$ ), 2.31 (s, 3H), 1.60 (s, 3H); low resolution MS (ES+)  $m/e$  517.1 ( $\text{MH}^+$ ).  
35

**Example 34: (2S)-2-(((Z)-1-methyl-3-oxo-3-(2-pyridinyl)-1-propenyl)amino)-3-(4-(2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy)phenyl)propanoic acid**

40 The title compound was prepared (as described above for the preparation of Example 2) from 441 mg (0.92 mmol) of Intermediate 45 (as the TFA salt) and 150 mg (0.92 mmol) of Intermediate 35 to yield 240 mg of Example 34: TLC (DCM/MeOH, 4/1):  $R_f = 0.46$ ;  $^1\text{H NMR}$  (DMSO- $d_6$ , 400 MHz)  $\delta$  11.49 (d, 1H,  $J = 9.2$ ), 8.52 (d, 1H,  $J = 4.8$ ), 7.90 (m, 4H), 7.47 (m, 4H), 7.11 (d, 2H,  $J = 8.4$ ), 6.80 (d, 2H,  $J = 8.4$ ), 6.13 (s, 1H), 4.12 (m, 3H), 3.16 (m, 2H),  
45 2.87 (t, 2H,  $J = 6.4$ ), 2.77 (dd, 1H,  $J = 13.6, 9.2$ ), 2.34 (s, 3H), 1.71 (s, 3H); low  
50 resolution MS (ES-)  $m/e$  510.1 ( $\text{M-H}$ ).  
55

**Example 35: (2S)-2-(((Z)-1-ethyl-3-(4-fluorophenyl)-3-oxo-1-propenyl)amino)-3-(4-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]phenyl)propanoic acid**

The title compound was prepared (as described above for the preparation of Example 2) from 100 mg (0.27 mmol) of Intermediate 45 and 49 mg (0.27 mmol) of Intermediate 32 to yield 83 mg of Example 35: TLC (EtOAc/MeOH (7:3):  $R_f$  = 0.28;  $^1\text{H}$  NMR (DMSO- $d_6$ , 400 MHz)  $\delta$  11.38 (d, 1 H,  $J$  = 7.00), 7.84 (m, 4H), 7.45 (m, 3H), 7.16 (t, 2H,  $J$  = 8.72), 7.09 (d, 2H,  $J$  = 7.69), 6.77 (d, 2H,  $J$  = 7.69), 5.49 (s, 1H), 4.11 (m, 2H), 4.01 (m, br, 1H), 3.13 (m, 1H), 2.86 (m, 2H), 2.72 (s, br, 1H), 2.30 (s, 3H), 1.62 (s, 3H); low resolution MS (ES+)  $m/e$  529 ( $\text{MH}^+$ ); RP-HPLC (Vydac C-18, 25cm x 4.6mm; 30-100%  $\text{CH}_3\text{CN}$  in  $\text{H}_2\text{O}$ ) with 0.1%  $\text{HCO}_2\text{H}$  buffer: 30 minutes; 1 mL/min:  $t_r$  = 20.10 min ( $t_0$  = 1.53).

**Example 36: (2S)-2-(((Z)-1-methyl-3-oxo-3-phenyl-1-propenyl)amino)-3-(4-[2-(5-methyl-2-phenyl-1,3-thiazol-4-yl)ethoxy]phenyl)propanoic acid**

The title compound was prepared (as described above for the preparation of Example 2) from 0.5 g (1.31 mmol) of Intermediate 46 and 215 mg (1.31 mmol) of benzoyl acetone and 260  $\mu\text{L}$  (1.31 mmol) of dicyclohexylamine (replacing TEA) to yield a total of 245 mg of Example 36 as a dull yellow solid. TLC (EtOAc/MeOH (3:1):  $R_f$  = 0.56;  $^1\text{H}$  NMR (DMSO- $d_6$ , 400 MHz)  $\delta$  11.4 (d, 1H,  $J$  = 9.1), 7.81 (m, 2H), 7.75 (m, 2H), 7.42 (m, 3H), 7.36 (m, 3H), 7.0 (d, 2H,  $J$  = 8.4), 6.78 (d, 2H,  $J$  = 8.6), 5.51 (s, 1H), 4.55 (m, 1H), 4.20 (t, 2H,  $J$  = 6.7), 4.0 (br s, 1H), 3.14-3.04 (m, 3H), 2.7 (m, 1H), 2.38 (s, 3H), 1.68 (s, 3H); low resolution MS (ES+)  $m/e$  527.1 ( $\text{MH}^+$ ).

**Example 37: (2S)-2-(((Z)-3-(2-fluorophenyl)-1-methyl-3-oxo-1-propenyl)amino)-3-(4-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]phenyl)propanoic acid**

The title compound was prepared (as described above for the preparation of Example 2) from 100 mg (0.27 mmol) of Intermediate 45 and

5 58 mg (0.32 mmol) of Intermediate 20 to yield 55 mg of Example 37: TLC  
(EtOAc/MeOH (7:3):  $R_f$  = 0.25;  $^1\text{H}$  NMR (DMSO- $d_6$ , 400 MHz)  $\delta$  11.42 (d, 1 H,  
10  $J$  = 8.93), 7.95 (m, 2H), 7.70 (dd, 1H,  $J$  = 1.78, 7.69), 7.51 (m, 4H), 7.23 (m,  
4H), 6.88 (d, 2H,  $J$  = 8.50), 5.36 (s, 1H), 4.22 (m, 2H), 4.16 (brs, 1H), 3.18 (m,  
5 1H), 2.94 (m, 2H), 2.84 (m, 1H), 2.40 (s, 3H), 1.70 (s, 3H); low resolution MS  
(ES-)  $m/e$  545.1 (M-H); RP-HPLC (Vydac C-18, 25cm x 4.6mm; 30-100%  
15  $\text{CH}_3\text{CN}$  in  $\text{H}_2\text{O}$ ) with 0.1%  $\text{HCO}_2\text{H}$  buffer: 30 minutes; 1 mL/min:  $t_r$  = 19.47 min  
( $t_0$  = 1.43).

10 **Example 38: (2S)-2-([(Z)-3-(2,3-difluorophenyl)-1-methyl-3-oxo-1-**  
20 **propenyl]amino)-3-{4-[2-(5-methyl-2-phenyl-1,3-oxazol-4-**  
**yl)ethoxy]phenyl}propanoic acid**

The title compound was prepared (as described above for the  
25 preparation of Example 12) from 120 mg (0.33 mmol) of Intermediate 45 and  
15 59 mg (0.33 mmol) of Intermediate 40 to yield 32 mg (18% yield) of Example  
38: TLC (DCM/MeOH (4:1):  $R_f$  = 0.72;  $^1\text{H}$  NMR (DMSO- $d_6$ , 300 MHz)  $\delta$  11.43  
(d, 1H,  $J$  = 9.8), 7.97 (m, 2H), 7.56 (m, 5h), 7.28 (m, 1H), 7.19 (d, 2H,  $J$  = 8.0),  
30 6.91 (d, 2H,  $J$  = 8.0), 5.34 (s, 1H), 4.21 (m, 3H), 3.21 (m, 2H), 2.98 (t, 2H,  $J$   
= 6.6), 2.84 (m, 1H), 2.40 (s, 3H), 1.78 (s, 3H); low resolution MS (ES+)  $m/e$   
20 547 ( $\text{MH}^+$ ).

35 **Example 39: (2S)-2-([(Z)-3-(2-hydroxyphenyl)-1-methyl-3-oxo-1-**  
**propenyl]amino)-3-{4-[2-(5-methyl-2-phenyl-1,3-oxazol-4-**  
**yl)ethoxy]phenyl}propanoic acid**

40 25 A solution of 100 mg (0.27 mmol) of Intermediate 45 and 58 mg (0.33  
mmol) of 2-(acetoacetyl)phenol (Aldrich Chemical Co.) in 5.0 mL MeOH, 1.0  
mL trimethyl orthoformate, and 4A sieves (50 mg) was refluxed at 80°C for  
16h. The reaction solution was concentrated and purified by silica gel flash  
45 column chromatography using DCM / MeOH 20 / 1 to 10/1 as eluent to afford  
30 135 mg of Example 39: TLC (DCM/MeOH (10:1):  $R_f$  = 0.32;  $^1\text{H}$  NMR (DMSO-  
 $d_6$ )  $\delta$  11.16 (d, 1 H,  $J$  = 8.72), 7.96 (m, 2H), 7.78 (m, 1H), 7.54 (m, 3H), 7.33  
(m, 1H), 7.20 (m, 2H), 6.84 (m, 4H), 5.69 (s, 1H), 4.26 (m, 2H), 4.17 (brs, 1H),  
50 3.20 (dd, 1H,  $J$  = 3.93, 13.67), 2.95 (m, 2H), 2.81 (m, 1H), 2.40 (s, 3H), 1.75



(s, 3H); low resolution MS (ES+)  $m/e$  527.0 ( $MH^+$ ); RP-HPLC (Vydac C-18, 25cm x 4.6mm; 50-100%  $CH_3CN$  in  $H_2O$ ) with 0.1%  $HCO_2H$  buffer: 30 minutes; 1 mL/min:  $t_r$  = 15.97 min ( $t_0$  = 1.43).

**Example 40: (2S)-3-{4-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]phenyl}-2-[(Z)-3-oxo-3-phenyl-1-propenyl]amino}propanoic acid**

The title compound was prepared as described above for the preparation of Example 2, from 1.0 g (2.05 mmol) of Intermediate 45 (as the TFA salt) and 0.8 g (4.2 mmol) of Intermediate 17 to yield 400 mg of Example 40 as a white solid:  $^1H$  NMR ( $DMSO-d_6$ , 400 MHz)  $\delta$  11.55 (d, 1H,  $J$  = 9.6), 7.87 (m, 2H), 7.75 (m, 2H), 7.45 (m, 3H), 7.35 (m, 3H), 7.10 (d, 2H,  $J$  = 8.6), 6.80 (d, 2H,  $J$  = 8.5), 5.52 (s, 1H), 4.12 (t, 2H,  $J$  = 6.6), 4.07 (m, 1H), 3.21 (dd, 1H,  $J$  = 13.7, 3.7), 2.85 (t, 2H,  $J$  = 6.5), 2.73 (dd, 1H,  $J$  = 13.7, 8.6), 2.3 (s, 3H), 1.90 (m, 2H), 1.4-1.25 (m, 2H), 0.76 (t, 3H,  $J$  = 7.4); low resolution MS (ES+)  $m/e$  539.2 ( $MH^+$ ).

**Example 41: (2S)-2-[(Z)-3-(4-methoxyphenyl)-1-methyl-3-oxo-1-propenyl]amino-3-{4-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]phenyl}propanoic acid**

The title compound was prepared (as described above for the preparation of Example 2) from 100 mg (0.27 mmol) of Intermediate 45 and 52 mg (0.27 mmol) of Intermediate 22 to yield 126 mg of Example 41. TLC (EtOAc/MeOH (7:3):  $R_f$  = 0.44:  $^1H$  NMR ( $DMSO-d_6$ , 400 MHz)  $\delta$  11.30 (d, 1H,  $J$  = 7.00), 7.87 (m, 2H), 7.73 (d, 2H,  $J$  = 8.72), 7.45 (m, 3H), 7.10 (d, 2H,  $J$  = 8.37), 6.88 (d, 2H,  $J$  = 8.89), 6.77 (d, 2H,  $J$  = 8.37), 5.74 (s, 1H), 4.11 (m, 2H), 4.01 (m, br, 1H), 3.13 (m, 1H), 2.86 (m, 2H), 2.72 (m, 1H), 2.31 (s, 3H), 1.65 (s, 3H); low resolution MS (ES-)  $m/e$  539 ( $M-H$ ); RP-HPLC (Vydac C-18, 25cm x 4.6mm; 10-50%  $CH_3CN$  in  $H_2O$ ) with 0.1%  $HCO_2H$  buffer: 30 minutes

**Example 42: (2S)-3-{4-[2-[2-(4-methoxyphenyl)-5-methyl-1,3-oxazol-4-yl]ethoxy]phenyl}-2-[(Z)-1-methyl-3-oxo-3-phenyl-1-propenyl]amino}propanoic acid**

The title compound was prepared (as described above for the preparation of Example 2) from 75 mg (0.19 mmol) of Intermediate 49 and 31 mg (0.19 mmol) of benzoylacetone to yield 30 mg of Example 42: TLC (DCM/MeOH (4:1):  $R_f$  = 0.45;  $^1\text{H}$  NMR (DMSO- $d_6$ , 400 MHz)  $\delta$  11.5 (d, 1H,  $J$  = 9.1), 7.89 (d, 2H,  $J$  = 8.8), 7.84 (m, 2H), 7.45 (m, 3H), 7.18 (d, 2H,  $J$  = 8.4); 7.09 (d, 2H,  $J$  = 8.8), 6.88 (d, 2H,  $J$  = 8.6), 5.63 (s, 1H), 4.2 (t, 2H,  $J$  = 6.5), 4.1 (m, 1H), 3.86 (s, 3H), 3.2 (m, 1H), 2.93 (t, 2H,  $J$  = 6.6), 2.83 (dd, 1H,  $J$  = 13.2, 6.4), 2.37 (s, 3H), 1.76 (s, 3H); low resolution MS (ES-)  $m/e$  539.2 (MH-).

**Example 43: (2S)-2-(((Z)-3-cyclohexyl-1-methyl-3-oxo-1-propenyl)amino)-3-(4-{2-[5-methyl-2-phenyl-1,3-oxazol-4-yl]ethoxy}phenyl)propanoic acid**

The title compound was prepared (as described above for the preparation of Example 2) from 428 mg (0.89 mmol) of Intermediate 45 (as the TFA salt) and 150 mg (0.89 mmol) of Intermediate 33 to yield 245 mg (53% yield) of Example 43 as a solid beige-colored glass: TLC (DCM/MeOH (4:1):  $R_f$  = 0.69;  $^1\text{H}$  NMR (DMSO- $d_6$ , 400 MHz)  $\delta$  10.79 (d, 1H,  $J$  = 9.0), 7.89 (m, 2H), 7.50-7.41 (m, 3H), 7.05 (d, 2H,  $J$  = 8.4), 6.77 (d, 2H,  $J$  = 8.4), 4.7 (s, 1H), 4.12 (t, 2H,  $J$  = 6.5), 3.9 (br s, 1H), 3.04 (dd, 1H,  $J$  = 13.6, 4.0), 2.87 (t, 2H,  $J$  = 6.5), 2.64 (dd, 1H,  $J$  = 13.6, 8.9), 2.32 (s, 3H), 1.9 (m, 1H), 1.6 (m, 4H), 1.5 (s, 3H), 1.25-1.05 (m, 4H); low resolution MS (ES+)  $m/e$  517.3 (MH $^+$ ).

**Example 44: (2S)-3-(4-{2-[2-(4-isopropoxyphenyl)-5-methyl-1,3-oxazol-4-yl]ethoxy}phenyl)-2-(((Z)-1-methyl-3-oxo-3-phenyl-1-propenyl)amino)propanoic acid**

The title compound was prepared (as described above for the preparation of Example 2) from 75 mg (0.18 mmol) of Intermediate 51 and 32 mg (0.194 mmol) of benzoylacetone to yield 55 mg of Example 44: TLC (DCM/MeOH (4:1):  $R_f$  = 0.50;  $^1\text{H}$  NMR (DMSO- $d_6$ , 400 MHz)  $\delta$  11.5 (d, 1H,  $J$  = 9.0), 7.85 (m, 4H), 7.45 (m, 3H), 7.18 (d, 2H,  $J$  = 8.4); 7.06 (d, 2H,  $J$  = 8.8), 6.87 (d, 2H,  $J$  = 8.5), 5.82 (s, 1H), 4.74 (m, 1H), 4.19 (t, 2H,  $J$  = 6.6), 4.1 (m, 1H), 3.19 (m, 1H), 2.92 (t, 2H,  $J$  = 6.6), 2.81 (dd, 1H,  $J$  = 13.9, 9.1), 2.36 (s,

3H), 1.74 (s, 3H), 1.34 (d, 6H,  $J=6.0$ ); low resolution MS (ES-)  $m/e$  567.2 (M-)

**Example 45: (2S)-2-(((Z)-1-heptyl-3-oxo-3-phenyl-1-propenyl)amino)-3-(4-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]phenyl)propanoic acid**

The title compound was prepared as described above for the preparation of Example 2, from 102 mg (0.28 mmol) of Intermediate 45 and 69 mg (0.28 mmol) of Intermediate 19 to yield 77 mg of Example 45: TLC (EtOAc/MeOH, 7/3):  $R_f = 0.56$ ;  $^1\text{H}$  NMR (DMSO- $d_6$ , 400 MHz)  $\delta$  11.55 (d, 1H,  $J=9.6$ ), 7.86 (d, 2H,  $J=6.8$ ), 7.76 (d, 2H,  $J=6.4$ ), 7.46 (m, 3H), 7.38 (m, 3H), 7.10 (d, 2H,  $J=8.4$ ), 6.78 (d, 2H,  $J=8.4$ ), 5.5 (s, 1H), 4.10 (t, 2H,  $J=6.6$ ), 4.0 (m, 1H), 3.16 (m, 1H), 2.86 (t, 2H,  $J=6.5$ ), 2.69 (dd, 1H,  $J=13.6, 9.6$ ), 2.31 (s, 3H), 1.87 (m, 2H), 1.25-1.1 (m, 10H), 0.78 (t, 3H,  $J=6.8$ ); low resolution MS (ES+)  $m/e$  595.1 (MH $^+$ ).

**Example 46: (2S)-2-(((Z)-1-methyl-3-(3-methylphenyl)-3-oxo-1-propenyl)amino)-3-(4-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]phenyl)propanoic acid**

The title compound was prepared (as described above for the preparation of Example 2) from 100 mg (0.27 mmol) of Intermediate 45 and 48 mg (0.27 mmol) of Intermediate 21 to yield 104 mg of Example 46.  $^1\text{H}$  NMR (DMSO- $d_6$ , 400 MHz)  $\delta$  11.47 (d, 1H,  $J=9.06$ ), 7.95 (m, 2H), 7.65 (m, 2H), 7.54 (m, 3H), 7.31 (m, 2H), 7.19 (d, 2H,  $J=8.51$ ), 6.88 (d, 2H,  $J=8.51$ ), 5.64 (s, 1H), 4.23 (m, 1H), 4.23 (m, 3H), 3.21 (m, 1H), 2.96 (m, 2H), 2.86 (m, 1H), 2.60 (s, 3H), 1.80 (s, 3H); low resolution MS (ES)  $m/e$  525.2 (MH $^+$ ); RP-HPLC (Vydac C-18, 25cm x 4.6mm; 50-100% CH $_3$ CN in H $_2$ O) with 0.1% HCO $_2$ H buffer: 30 minutes; 1 mL/min:  $t_r = 15.57$  min ( $t_0 = 1.43$ ).

**Example 47: (2S)-2-(((Z)-1-ethyl-3-oxo-3-phenyl-1-propenyl)amino)-3-(4-[2-[2-(4-methoxyphenyl)-5-methyl-1,3-oxazol-4-yl)ethoxy]phenyl)propanoic acid**

The title compound was prepared (as described above for the preparation of Example 2) from 100 mg (0.25 mmol) of Intermediate 49 and

45 mg (0.25 mmol) of Intermediate 16 to yield 95 mg (68% yield) of Example 47 as a solid: TLC  $R_f$  = 0.2-0.3 (slight streak, 4/1,  $\text{CH}_2\text{Cl}_2/\text{MeOH}$ );  $^1\text{H}$  NMR ( $\text{DMSO}-d_6$ , 400 MHz)  $\delta$  11.5 (d, 1H,  $J$  = 9.0), 7.85 (m, 4H), 7.45 (m, 3H), 7.18 (d, 2H,  $J$  = 8.4); 7.06 (d, 2H,  $J$  = 8.8), 6.87 (d, 2H,  $J$  = 8.5), 5.82 (s, 1H), 4.74 (m, 1H), 4.19 (t, 2H,  $J$  = 6.6), 4.1 (m, 1H), 3.19 (m, 1H), 2.92 (t, 2H,  $J$  = 6.6), 2.81 (dd, 1H,  $J$  = 13.9, 9.1), 2.36 (s, 3H), 1.74 (s, 3H), 1; low resolution MS ( $\text{ES}^+$ )  $m/e$  555.1 ( $\text{MH}^+$ );

**Example 48: (2S)-3-{4-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]phenyl}-2-[(Z)-3-oxo-3-phenyl-1-(trifluoromethyl)-1-propenyl]amino}propanoic acid**

The title compound was prepared (as described above for the preparation of Example 12) from 1.3 g (3.55 mmol) of Intermediate 45 and 0.7 g (3.55 mmol) of Intermediate 44 to yield 0.84 g (42% yield) of Example 48 as a solid: TLC ( $\text{DCM}/\text{MeOH}$  (4:1):  $R_f$  = 0.55;  $^1\text{H}$  NMR ( $\text{DMSO}-d_6$ , 300 MHz)  $\delta$  10.99 (s, 1H), 7.94 (m, 4H), 7.68 (m, 6H), 7.14 (d, 2H,  $J$  = 8.4), 6.84 (d, 2H,  $J$  = 8.4), 6.21 (s, 1H), 4.19 (m, 3H), 3.17 (m, 1H), 2.92-2.85 (m, 3H), 2.36 (s, 3H); low resolution MS ( $\text{ES}^+$ )  $m/e$  565.0 ( $\text{MH}^+$ ).

**Example 49 (2S)-3-{4-[2-(5-methyl-2-phenyl-1,3-thiazol-4-yl)ethoxy]phenyl}-2-[(Z)-3-oxo-3-phenyl-1-(trifluoromethyl)-1-propenyl]amino}propanoic acid**

The title compound was prepared (as described above for the preparation of Example 12) from 120 mg (0.31 mmol) of Intermediate 46 and 62 mg (0.31 mmol) of Intermediate 44 to yield 78 mg of Example 49: TLC ( $\text{DCM}/\text{MeOH}$  (4:1):  $R_f$  = 0.58;  $^1\text{H}$  NMR ( $\text{DMSO}-d_6$ , 300 MHz)  $\delta$  10.92 (br s, 1H), 7.88 (d, 2H,  $J$  =), 7.81 (m, 2H), 7.56-7.37 (m, 6H), 7.01 (d, 2H,  $J$  = 8.5), 6.78 (d, 2H,  $J$  = 8.3), 6.13 (s, 1H), 4.19 (t, 2H,  $J$  = 6.5), 4.10 (br s, 1H), 3.15-3.11 (m, 1H), 3.05 (t, 2H,  $J$  = 6.6), 2.87 (dd, 1H,  $J$  = 13.7, 7.7), 2.37 (s, 3H); low resolution MS ( $\text{ES}^+$ )  $m/e$  581.0 ( $\text{MH}^+$ ).

**Example 50: (2S)-2-(((Z)-1-ethyl-3-oxo-3-(4-(trifluoromethyl)phenyl)-1-propenyl)amino)-3-(4-(2-(2-(4-fluorophenyl)-5-methyl-1,3-oxazol-4-yl)ethoxy)phenyl)propanoic acid**

The title compound was prepared (as described above for the preparation of Example 2) from 150 mg (0.39 mmol) of Intermediate 46 and 95 mg (0.39 mmol) of Intermediate 25 to yield 79 mg (33% yield) of Example 50: TLC (DCM/MeOH, 4/1):  $R_f$  = 0.62;  $^1\text{H}$  NMR (DMSO- $d_6$ , 300 MHz)  $\delta$  11.59 (d, 1H,  $J$  = 9.0), 7.97-7.88 (m, 4H), 7.72 (d, 2H,  $J$  = 8.4), 7.29 (t, 2H,  $J$  = 9.0), 7.11 (d, 2H,  $J$  = 8.4), 6.78 (d, 2H,  $J$  = 8.4), 5.59 (s, 1H), 4.11 (m, 3H), 3.17 (m, 2H), 2.85 (t, 2H,  $J$  = 6.3), 2.77 (m, 1H), 2.29 (s, 3H), 2.03 (m, 2H), 0.91 (t, 3H,  $J$  = 7.5); low resolution MS ( $\text{ES}^+$ )  $m/e$  611.2 ( $\text{MH}^+$ ).

**Example 51: (2S)-2-(((Z)-1-ethyl-3-oxo-3-(4-(trifluoromethyl)phenyl)-1-propenyl)amino)-3-(4-(2-(2-(4-isopropoxyphenyl)-5-methyl-1,3-oxazol-4-yl)ethoxy)phenyl)propanoic acid**

The title compound was prepared (as described above for the preparation of Example 2) from 151 mg (0.39 mmol) of Intermediate 51 and 92 mg (0.39 mmol) of Intermediate 25 to yield 91 mg (36% yield) of Example 51: TLC (DCM/MeOH, 4/1):  $R_f$  = 0.60;  $^1\text{H}$  NMR (DMSO- $d_6$ , 300 MHz)  $\delta$  11.6 (d, 1H,  $J$  = 9.6), 7.96 (d, 2H,  $J$  = 8.1), 7.78-7.70 (m, 4H), 7.10 (d, 2H,  $J$  = 8.4), 6.96 (d, 2H,  $J$  = 8.7), 6.78 (d, 2H,  $J$  = 8.4), 5.59 (s, 1H), 4.62 (sept, 1H,  $J$  = 6.0), 4.13 (m, 3H), 3.16 (m, 1H), 2.84-2.72 (m, 3H), 2.26 (s, 3H), 2.05-1.96 (m, 2H), 1.24 (d, 6H,  $J$  = 6.0), 0.91 (t, 3H,  $J$  = 7.5); low resolution MS ( $\text{ES}^+$ )  $m/e$  651.3 ( $\text{MH}^+$ ).

**Example 52: (2S)-2-(((Z)-1-ethyl-3-oxo-3-(4-(trifluoromethyl)phenyl)-1-propenyl)amino)-3-(4-(2-(2-(4-methoxyphenyl)-5-methyl-1,3-oxazol-4-yl)ethoxy)phenyl)propanoic acid**

The title compound was prepared (as described above for the preparation of Example 2) from 150 mg (0.38 mmol) of Intermediate 49 and 93 mg (0.38 mmol) of Intermediate 25 to yield 45 mg (19% yield) of Example 52: TLC (DCM/MeOH, 4/1):  $R_f$  = 0.63;  $^1\text{H}$  NMR (DMSO- $d_6$ , 300 MHz)  $\delta$  11.6

(d, 1H,  $J=9.6$ ), 7.96 (d, 2H,  $J=8.1$ ), 7.81-7.70 (m, 4H), 7.11 (d, 2H,  $J=8.4$ ), 7.0 (d, 2H,  $J=8.7$ ), 6.78 (d, 2H,  $J=8.4$ ), 5.6 (s, 1H), 4.18-4.07 (m, 3H), 3.77 (s, 3H), 3.17 (m, 1H), 2.85-2.72 (m, 3H), 2.26 (s, 3H), 2.05-1.96 (m, 2H), 0.91 (t, 3H,  $J=7.5$ ); low resolution MS ( $ES^+$ )  $m/e$  623.2 ( $MH^+$ ).

**Example 53: (2S)-3-{4-[2-[2-(4-fluorophenyl)-5-methyl-1,3-oxazol-4-yl]ethoxy]phenyl}-2-[(Z)-3-oxo-3-phenyl-1-(trifluoromethyl)-1-propenyl]amino}propanoic acid**

The title compound was prepared (as described above for the preparation of Example 12) from 203 mg (0.53 mmol) of Intermediate 46 and 130 mg (0.63 mmol) of Intermediate 44 to yield 154 mg (50% yield) of Example 53: TLC (DCM/MeOH, 4/1):  $R_f = 0.65$ ;  $^1H$  NMR (DMSO- $d_6$ , 400 MHz)  $\delta$  11.5 (d, 1H,  $J=8.9$ ), 7.95 (d, 2H,  $J=7.2$ ), 7.77 (d, 2H,  $J=8.7$ ), 7.72 (d, 2H,  $J=8.2$ ), 7.1 (d, 2H,  $J=8.4$ ), 6.97 (d, 2H,  $J=8.9$ ), 6.79 (d, 2H,  $J=8.5$ ), 5.58 (s, 1H), 4.65 (m, 1H), 4.11 (t, 2H,  $J=6.5$ ), 4.1 (m, 1H), 3.16 (dd, 1H,  $J=13.7$ , 3.6), 2.84 (t, 2H,  $J=6.5$ ), 2.74 (dd, 1H, 13.7, 9.0), 2.28 (s, 3H), 1.67 (s, 3H), 1.25 (d, 6H,  $J=6.0$ ); low resolution MS ( $ES^+$ )  $m/e$  637.1 ( $MH^+$ )

**Example 54: (2S)-2-[(Z)-1-methyl-3-oxo-3-phenyl-1-propenyl]amino}-3-{4-[(5-methyl-2-phenyl-1,3-oxazol-4-yl)methoxy]phenyl}propanoic acid**

The title compound was prepared (as described above for the preparation of example 1) from a suspension of 113 mg (0.32 mmol) of Intermediate 54 and 52 mg (0.32 mmol) of benzoylacetone to yield 44 mg (28%) of example 54: TLC (DCM/MeOH, 4/1):  $R_f = 0.55$ ;  $^1H$  NMR (DMSO- $d_6$ , 300 MHz)  $\delta$  11.43 (d, 1H,  $J=9.0$ ), 7.92-7.88 (m, 2H), 7.8 (m, 2H), 7.48 (m, 3H), 7.38-7.38 (m, 3H), 7.16 (d, 2H,  $J=8.4$ ), 6.9 (d, 2H,  $J=8.4$ ), 5.53 (s, 1H), 4.91 (s, 1H), 4.15 (m, 1H), 3.2 (m, 2H), 2.83-2.74 (m, 1H), 2.37 (s, 3H), 1.67 (s, 3H); low resolution MS ( $ES^+$ )  $m/e$  497.07 ( $MH^+$ ).

**Example 55: (2S)-3-{4-[2-[5-ethyl-2-phenyl-1,3-oxazol-4-yl]ethoxy]phenyl}-2-[(Z)-1-methyl-3-oxo-3-phenyl-1-propenyl]amino}propanoic acid**

5 The title compound was prepared (as described above for the preparation of example 1) from a suspension of 205 mg (0.54 mmol) of Intermediate 52 and 87 mg (0.54 mmol) of benzoylacetone to yield 145 mg (51%) of example 55: TLC (DCM/MeOH, 4/1):  $R_f$  = 0.60;  $^1\text{H}$  NMR (DMSO- $d_6$ , 300 MHz)  $\delta$  11.42 (d, 1H,  $J$  = 9.0), 7.89-7.86 (m, 2H), 7.77-7.74 (m, 2H), 7.48-7.32 (m, 6H), 7.11 (d, 2H,  $J$  = 8.4), 6.77 (d, 2H,  $J$  = 8.4), 5.53 (s, 1H), 4.1 (t, 1H,  $J$  = 6.3), 3.2-3.14 (m, 2H), 2.86 (t, 2H,  $J$  = 6.3), 2.79-2.74 (m, 1H), 2.67 (q, 2H,  $J$  = 7.5), 1.65 (s, 3H), 1.17 (t, 3H,  $J$  = 7.5); low resolution MS ( $\text{ES}^+$ )  $m/e$  525.2 ( $\text{MH}^+$ ).

10  
20 **Example 56: (2S)-2-([(Z)-1-methyl-3-oxo-3-phenyl-1-propenyl]amino)-3-{4-[2-(2-phenyl-5-propyl-1,3-oxazol-4-yl)ethoxy]phenyl}propanoic acid**

The title compound was prepared (as described above for the preparation of example 1) from a suspension of 204 mg (0.52 mmol) of Intermediate 53 and 84 mg (0.52 mmol) of benzoylacetone to yield 137 mg (49%) of example 56: TLC (DCM/MeOH, 4/1):  $R_f$  = 0.60;  $^1\text{H}$  NMR (DMSO- $d_6$ , 300 MHz)  $\delta$  11.42 (d, 1H,  $J$  = 9.0), 7.89-7.86 (m, 2H), 7.77-7.74 (m, 2H), 7.49-7.32 (m, 6H), 7.11 (d, 2H,  $J$  = 8.4), 6.77 (d, 2H,  $J$  = 8.4), 5.53 (s, 1H), 4.1 (t, 1H,  $J$  = 6.3), 3.19-3.13 (m, 2H), 2.86 (t, 2H,  $J$  = 6.3), 2.78-2.71 (m, 1H), 2.64 (t, 2H,  $J$  = 7.2), 1.65-1.54 (m, 5H), 0.88 (t, 3H,  $J$  = 7.5); low resolution MS ( $\text{ES}^+$ )  $m/e$  539.19 ( $\text{MH}^+$ ).

35  
40 **Example 57: (2S)-2-([(Z)-1-methyl-3-oxo-3-phenyl-1-propenyl]amino)-3-{4-[3-(5-methyl-2-phenyl-1,3-oxazol-4-yl)propoxy]phenyl}propanoic acid**

The title compound was prepared (as described above for the preparation of example 1) from a suspension of 205 mg (0.54 mmol) of Intermediate 55 and 88 mg (0.54 mmol) of benzoylacetone to yield 142 mg (50% yield) of example 57: TLC (DCM/MeOH, 4/1):  $R_f$  = 0.66;  $^1\text{H}$  NMR (DMSO- $d_6$ , 300 MHz)  $\delta$  11.44 (d, 1H,  $J$  = 9.0), 7.88-7.85 (m, 2H), 7.78-7.75 (m, 2H), 7.49-7.43 (m, 3H), 7.37-7.32 (m, 3H), 7.12 (d, 2H,  $J$  = 8.4), 6.78 (d, 2H,  $J$  = 8.4), 5.54 (s, 1H), 4.13 (m, 1H), 3.87 (t, 1H,  $J$  = 6.0), 3.22-3.16 (m, 1H), 2.77

(d, 1H,  $J = 14.4, 9.2$ ), 2.56 (t, 2H,  $J = 7.2$ ), 2.22 (s, 3H), 2.00-1.93 (m, 2H), 1.66 (s, 3H); low resolution MS (ES<sup>+</sup>)  $m/e$  525.15 (MH<sup>+</sup>).

**Example 58: (2S)-3-{4-[2-(5-ethyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]phenyl}-2-[(Z)-3-oxo-3-phenyl-1-(trifluoromethyl)-1-propenyl]amino}propanoic acid**

A mixture of 100 mg (0.26 mmol, 1.0 equivalent) of Intermediate 52, 110 mg (0.39 mmol, 1.5 equivalents) of Intermediate 44, and 0.14 mL (0.8 mmol, 3.0 equivalents) of DIEA in 2 mL of MeOH was stirred for 2.5 hours at ambient temperature (22 °C). The solvent was evaporated under reduced pressure. The residue was taken into EtOAc (20 mL) and washed with 10 mL 0.1 N HCl solution and 10 mL brine. The organics were dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, concentrated, and the residue purified directly by silica gel chromatography. Elution with 10%-20% MeOH in EtOAc gave 55 mg (36% yield) of the title compound as a solid after drying under vacuum for several hours: TLC (DCM/MeOH, 4/1):  $R_f = 0.65$ ; <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>, 400 MHz)  $\delta$  10.95 (br s, 1H), 7.88-7.93 (m, 4H), 7.52-7.41 (m, 6H), 7.72 (d, 2H,  $J = 8.2$ ), 7.02 (d, 2H,  $J = 8.4$ ), 6.73 (d, 2H,  $J = 8.5$ ), 6.06 (s, 1H), 4.09 (t, 2H,  $J = 6.5$ ), 3.99 (m, 1H), 3.0 (m, 1H, NH), 2.86-2.79 (m, 4H), 2.66 (q, 2H, 7.5), 1.15 (t, 3H,  $J = 7.5$ ); low resolution MS (ES<sup>+</sup>)  $m/e$  579.08 (MH<sup>+</sup>)

**Example 59: (2S)-3-{4-[3-(5-methyl-2-phenyl-1,3-oxazol-4-yl)propoxy]phenyl}-2-[(Z)-3-oxo-3-phenyl-1-(trifluoromethyl)-1-propenyl]amino}propanoic acid**

The title compound was prepared (as described above for the preparation of Example 58) from 100 mg (0.26 mmol) of Intermediate 55 and 110 mg (0.39 mmol) of Intermediate 44 to yield 65 mg (42% yield) of Example 59: TLC (DCM/MeOH, 4/1):  $R_f = 0.58$ ; <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>, 400 MHz)  $\delta$  10.74 (d, 1H,  $J = 9.4$ ), 7.90 (d, 2H,  $J = 7.5$ ), 7.86-7.82 (m, 2H), 7.54 (m, 1H), 7.48-7.40 (m, 5H), 7.09 (d, 2H,  $J = 8.4$ ), 6.81 (d, 2H,  $J = 8.4$ ), 6.25 (s, 1H), 4.3 (br s, 1H), 3.89 (t, 2H,  $J = 6.1$ ), 3.09 (dd, 1H,  $J = 14.0, 4.8$ ), 2.96 (dd, 1H, 14.0, 7.9), 2.2 (s, 3H), 2.0-1.92 (m, 2H); low resolution MS (ES<sup>+</sup>)  $m/e$  579.08 (MH<sup>+</sup>)



**Example 60: (2S)-3-(4-{2-[5-ethyl-2-phenyl-1,3-thiazol-4-yl]ethoxy}phenyl)-2-[(Z)-3-oxo-3-phenyl-1-(trifluoromethyl)-1-propenyl]amino}propanoic acid**

The title compound was prepared (as described above for the preparation of Example 58) from 180 mg (0.35 mmol) of Intermediate 57 (as its TFA salt) and 148 mg (0.53 mmol) of Intermediate 44 to yield 113 mg (54% yield) of Example 60: TLC (DCM/MeOH, 85/15):  $R_f = 0.50$ ;  $^1\text{H}$  NMR (DMSO- $d_6$ , 300 MHz)  $\delta$  10.72 (d, 1H,  $J = 10.3$ ), 7.94 (d, 2H,  $J = 7.3$ ), 7.87-7.83 (m, 2H), 7.67-7.39 (m, 5H), 7.13 (d, 2H,  $J = 8.6$ ), 6.86 (d, 2H,  $J = 8.5$ ), 6.32 (s, 1H), 4.26 (t, 2H,  $J = 6.5$ ), 3.17-3.01 (m, 4H), 2.81 (q, 2H,  $J = 7.4$ ), 1.22 (t, 3H,  $J = 7.4$ ); low resolution MS (ES+)  $m/e$  594.9 ( $\text{MH}^+$ )

**Example 61: (2S)-3-(4-{2-[5-ethyl-2-(4-fluorophenyl)-1,3-oxazol-4-yl]ethoxy}phenyl)-2-[(Z)-3-oxo-3-phenyl-1-(trifluoromethyl)-1-propenyl]amino}propanoic acid**

The title compound was prepared (as described above for the preparation of Example 58) from 130 mg (0.26 mmol) of Intermediate 58, 106 mg (0.38 mmol) of Intermediate 44, and 0.14 mL diisopropylethylamine in 5 mL MeOH. The crude product was recrystallized from hot  $\text{CH}_3\text{CN}$  to yield 67 mg (43% yield) of Example 61: TLC (DCM/MeOH, 85/15):  $R_f = 0.44$ ;  $^1\text{H}$  NMR (DMSO- $d_6$ , 300 MHz)  $\delta$  10.72 (d, 1H,  $J = 10.3$ ), 7.97-7.92 (m, 4H), 7.62-7.56 (m, 1H), 7.52-7.46 (m, 3H), 7.32 (t, 2H,  $J = 8.7$ ), 7.13 (d, 2H,  $J = 8.6$ ), 6.85 (d, 2H,  $J = 8.6$ ), 6.32 (s, 1H), 4.35 (m, 1H), 4.16 (t, 2H,  $J = 6.5$ ), 3.13 (dd, 1H,  $J = 14.0, 4.8$ ), 3.02 (dd, 1H,  $J = 14.0, 7.7$ ), 2.89 (t, 2H,  $J = 6.5$ ), 2.69 (q, 2H,  $J = 7.4$ ), 1.19 (t, 3H,  $J = 7.5$ ); low resolution MS (ES+)  $m/e$  596.9 ( $\text{MH}^+$ ).

**Example 62: (2S)-3-(4-{3-[5-ethyl-2-(4-fluorophenyl)-1,3-oxazol-4-yl]propoxy}phenyl)-2-[(Z)-3-oxo-3-phenyl-1-(trifluoromethyl)-1-propenyl]amino}propanoic acid**

The title compound was prepared (as described above for the preparation of Example 58) from 200 mg (0.38 mmol) of Intermediate 59, 127 mg (0.57 mmol) of Intermediate 44, and 0.2 mL diisopropylethylamine in 2.5 mL MeOH. The crude product was recrystallized from hot  $\text{CH}_3\text{CN}$  to yield 106

mg (46% yield) of Example 62: TLC (DCM/MeOH, 4/1):  $R_f$  = 0.61;  $^1\text{H}$  NMR (DMSO- $d_6$ , 300 MHz)  $\delta$  10.72 (d, 1H,  $J$  = 10.2), 7.95-7.91 (m, 4H), 7.76-7.61 (m, 3H), 7.31 (t, 2H,  $J$  = 8.7), 7.13 (d, 2H,  $J$  = 8.1), 6.85 (d, 2H,  $J$  = 8.4), 6.32 (s, 1H), 4.42-4.37 (m, 1H), 3.92 (t, 1H,  $J$  = 5.7), 3.15-2.98 (m, 3H), 2.60-2.65 (m, 4H), 2.0 (t, 2H,  $J$  = 6.3), 1.10 (t, 3H,  $J$  = 7.5); low resolution MS (ES $^+$ )  $m/e$  610.96 (MH $^+$ ).

**Example 63: (2S)-3-(4-(2-(5-ethyl-2-(4-fluorophenyl)-1,3-thiazol-4-yl)ethoxy)phenyl)-2-(((E)-3-oxo-3-phenyl-1-(trifluoromethyl)-1-propenyl)amino)propanoic acid**

The title compound was prepared (as described above for the preparation of Example 58) from 130 mg (0.25 mmol) of Intermediate 56, 103 mg (0.37 mmol) of Intermediate 44, and 0.17 mL diisopropylethylamine in 5 mL MeOH. The crude product was recrystallized from hot CH<sub>3</sub>CN to yield 110 mg (72% yield) of Example 63: TLC (DCM/MeOH, 85/15):  $R_f$  = 0.45;  $^1\text{H}$  NMR (DMSO- $d_6$ , 300 MHz)  $\delta$  10.72 (d, 1H,  $J$  = 10.2), 7.96-7.86 (m, 4H), 7.59 (m, 1H), 7.52-7.46 (m, 3H), 7.28 (t, 2H,  $J$  = 8.8), 7.13 (d, 2H,  $J$  = 8.6), 6.85 (d, 2H,  $J$  = 8.5), 6.32 (s, 1H), 4.38 (m, 1H), 4.25 (t, 2H,  $J$  = 6.5), 3.31-2.98 (m, 4H), 2.81 (q, 2H,  $J$  = 7.4), 1.22 (t, 3H,  $J$  = 7.4); low resolution MS (ES $^+$ )  $m/e$  612.9 (MH $^+$ ).

**Example 64: (2S)-3-(4-(3-(5-ethyl-2-(4-fluorophenyl)-1,3-oxazol-4-yl)propoxy)phenyl)-2-(((Z)-1-ethyl-3-oxo-3-(4-(trifluoromethyl)phenyl)-1-propenyl)amino)propanoic acid**

The title compound was prepared (as described above for the preparation of Example 2) from 95 mg (0.18 mmol) of Intermediate 59 (TFA Salt), 44 mg (0.18 mmol) of Intermediate 25, and 0.055 mL triethylamine in MeOH (1.5 mL) and trimethylorthoformate (0.5 mL) to yield 44 mg (38% yield) of Example 64 after silica gel chromatography (5%-15% MeOH in DCM elution): TLC (DCM/MeOH, 4/1):  $R_f$  = 0.53;  $^1\text{H}$  NMR (DMSO- $d_6$ , 300 MHz)  $\delta$

11.63 (d, 1H,  $J=9.3$ ), 7.99-7.89 (m, 4H), 7.73 (d, 2H,  $J=8.1$ ), 7.30 (t, 2H,  $J=8.7$ ), 7.13 (d, 2H,  $J=8.1$ ), 6.79 (d, 2H,  $J=8.1$ ), 5.62 (s, 1H), 4.20 (m, 1H), 3.87 (m, 2H), 3.20 (m, 2H), 2.77 (m, 2H), 2.61-2.54 (m, 4H), 2.05-1.96 (m, 4H), 1.10 (t, 3H,  $J=7.2$ ), 0.92 (t, 3H,  $J=7.5$ ); low resolution MS ( $ES^+$ )  $m/e$  638.93 ( $MH^+$ ).

**Example 65: (2S)-2-(((Z)-1-ethyl-3-oxo-3-[4-(trifluoromethyl)phenyl]-1-propenyl)amino)-3-[4-[3-(5-ethyl-2-phenyl-1,3-oxazol-4-yl)propoxy]phenyl]propanoic acid**

The title compound was prepared (as described above for the preparation of Example 2) from 50 mg (0.13 mmol) of Intermediate 60, 31 mg (0.13 mmol) of Intermediate 25, and 0.044 mL triethylamine in MeOH (1.0 mL) and trimethylorthoformate (0.5 mL) to yield 22 mg (28% yield) of Example 65 after silica gel chromatography (5%-15% MeOH in DCM elution): TLC (DCM/MeOH, 4/1):  $R_f = 0.50$ ;  $^1H$  NMR (DMSO- $d_6$ , 300 MHz)  $\delta$  11.70 (d, 1H,  $J=9.6$ ), 8.05 (d, 2H,  $J=7.8$ ), 7.95 (d, 2H,  $J=7.8$ ), 7.81 (d, 2H,  $J=8.1$ ), 7.55-7.52 (m, 3H), 7.20 (d, 2H,  $J=8.4$ ), 6.86 (d, 2H,  $J=8.4$ ), 5.69 (s, 1H), 4.28-4.22 (m, 1H), 3.96 (d, 2H,  $J=6.0$ ), 3.28-3.20 (m, 2H), 2.89-2.81 (m, 1H), 2.72-2.63 (m, 4H), 2.05-1.96 (m, 4H), 1.18 (t, 3H,  $J=7.5$ ), 0.99 (t, 3H,  $J=7.5$ ); low resolution MS ( $ES^+$ )  $m/e$  620.98 ( $MH^+$ ).

**Example 66: (2S)-3-[4-[3-[5-ethyl-2-(4-fluorophenyl)-1,3-oxazol-4-yl]propoxy]phenyl]-2-(((Z)-1-methyl-3-oxo-3-[4-(trifluoromethyl)phenyl]-1-propenyl)amino)propanoic acid**

The title compound was prepared (as described above for the preparation of Example 2) from 90 mg (0.17 mmol) of Intermediate 59, 40 mg (0.17 mmol) of Intermediate 24, and 0.05 mL triethylamine in MeOH (1.5 mL) and trimethylorthoformate (0.5 mL) to yield 54 mg (51% yield) of Example 66 after silica gel chromatography (5%-15% MeOH in DCM): TLC (DCM/MeOH, 4/1):  $R_f = 0.60$ ;  $^1H$  NMR (DMSO- $d_6$ , 300 MHz)  $\delta$  11.53 (d, 1H,  $J=9.0$ ), 7.98-7.90 (m, 4H), 7.73 (d, 2H,  $J=8.4$ ), 7.30 (t, 2H,  $J=8.7$ ), 7.13 (d, 2H,  $J=8.4$ ),

6.79 (d, 2H,  $J=8.4$ ), 5.61 (s, 1H), 4.23-4.19 (m, 1H), 3.89 (t, 2H,  $J=6.0$ ), 3.22-3.17 (m, 2H), 2.83-2.75 (m, 2H), 2.64-2.55 (m, 4H), 1.97 (t, 2H,  $J=6.3$ ), 1.71 (s, 3H), 1.10 (t, 3H,  $J=7.5$ ); low resolution MS ( $ES^+$ )  $m/e$  625.15 ( $MH^+$ ).

**Example 67: (2S)-3-{4-[3-(5-ethyl-2-phenyl-1,3-oxazol-4-yl)propoxy]phenyl}-2-[(Z)-1-methyl-3-oxo-3-[4-(trifluoromethyl)phenyl]-1-propenyl]amino}propanoic acid**

The title compound was prepared (as described above for the preparation of Example 2) from 57 mg (0.14 mmol) of Intermediate 60, 33 mg (0.14 mmol) of Intermediate 24, and 0.042 mL triethylamine in MeOH (1.5 mL) and trimethylorthoformate (0.5 mL) to yield 24 mg (28% yield) of Example 67 after silica gel chromatography (5%-15% MeOH in DCM): TLC (DCM/MeOH, 4/1):  $R_f = 0.62$ ;  $^1H$  NMR (DMSO- $d_6$ , 300 MHz)  $\delta$  11.53 (d, 1H,  $J=9.0$ ), 7.97 (d, 2H,  $J=8.4$ ), 7.89 (d, 2H,  $J=7.8$ ), 7.73 (d, 2H,  $J=8.4$ ), 7.46 (m, 3H), 7.12 (d, 2H,  $J=8.1$ ), 6.79 (d, 2H,  $J=8.4$ ), 5.62 (s, 1H), 4.25-4.19 (m, 1H), 3.90 (t, 2H,  $J=6.0$ ), 3.21-3.15 (m, 1H), 2.58-2.83 (m, 5H), 1.98 (t, 2H,  $J=5.7$ ), 1.72 (s, 3H), 1.11 (t, 3H,  $J=7.5$ ); low resolution MS ( $ES^+$ )  $m/e$  607.18 ( $MH^+$ ).

**Example 68: (2S)-3-{4-[3-(5-ethyl-2-phenyl-1,3-oxazol-4-yl)propoxy]phenyl}-2-[(Z)-3-oxo-3-phenyl-1-(trifluoromethyl)-1-propenyl]amino}propanoic acid**

The title compound was prepared (as described above for the preparation of Example 58) from 100 mg (0.25 mmol) of Intermediate 60, 85 mg (0.38 mmol) of Intermediate 44, and 0.14 mL diisopropylethylamine in 2.0 mL MeOH. Attempted recrystallization from hot  $CH_3CN$  without success. Silica gel chromatography with MeOH/DCM (10%-20%) gave 68 mg (45% yield) of Example 68: TLC (DCM/MeOH, 4/1):  $R_f = 0.60$ ;  $^1H$  NMR (DMSO- $d_6$ , 300 MHz)  $\delta$  10.96 (br s, 1H), 7.90-7.87 (m, 4H), 7.54-7.46 (m, 6H), 7.07 (d, 2H,  $J=8.4$ ), 6.78 (d, 2H,  $J=8.1$ ), 6.12 (s, 1H), 4.11-4.03 (m, 1H), 3.89 (t, 2H,  $J=6.0$ ), 3.12-3.07 (m, 1H), 2.91-2.83 (m, 1H), 2.66-2.57 (m, 4H), 1.98 (t, 2H,  $J=5.7$ ), 1.11 (t, 3H,  $J=7.5$ ); low resolution MS ( $ES^+$ )  $m/e$  593.02 ( $MH^+$ ).

**Example 69: (2S)-3-{4-[(5-ethyl-2-phenyl-1,3-thiazol-4-yl)methoxy]phenyl}-2-[(Z)-3-oxo-3-phenyl-1-(trifluoromethyl)-1-propenyl]amino}propanoic acid**

The title compound was prepared (as described above for the preparation of Example 58) from 1.5 g (3.02 mmol) of Intermediate 62, 1.01 g (4.53 mmol) of Intermediate 44, and 1.6 mL diisopropylethylamine in 20 mL MeOH. Attempted recrystallization from hot CH<sub>3</sub>CN without success. Silica gel chromatography with MeOH/DCM (10%-20%) gave 1.05 g (60% yield) of Example 69: TLC (DCM/MeOH, 4/1): R<sub>f</sub> = 0.43; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 300 MHz) δ 10.98 (br s, 1H), 7.91-7.85 (m, 4H), 7.55-7.45 (m, 6H), 7.11 (d, 2H, J = 8.4), 6.92 (d, 2H, J = 8.4), 6.15 (s, 1H), 5.07 (s, 2H), 4.13-4.06 (m, 1H), 3.16-3.09 (m, 1H), 2.95-2.85 (m, 4H), 1.21 (t, 3H, J = 7.5); low resolution MS (ES<sup>+</sup>)m/e 580.95 (MH<sup>+</sup>).

**Example 70: (2S)-3-{4-[(5-ethyl-2-phenyl-1,3-thiazol-4-yl)methoxy]phenyl}-2-[(Z)-1-methyl-3-oxo-3-[4-(trifluoromethyl)phenyl]-1-propenyl]amino}propanoic acid**

The title compound was prepared (as described above for the preparation of Example 2) from 1.35 g (2.72 mmol) of Intermediate 62, 630 mg (2.72 mmol) of Intermediate 24, and 0.8 mL triethylamine in MeOH (20 mL) and trimethylorthoformate (5 mL) to yield 1.04 g (65% yield) of Example 70 after silica gel chromatography (5%-15% MeOH in DCM): TLC (DCM/MeOH, 4/1): R<sub>f</sub> = 0.45; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 300 MHz) δ 11.57 (d, 1H, J = 8.7), 7.96 (d, 2H, J = 8.4), 7.85 (m, 2H), 7.71 (d, 2H, J = 8.4), 7.44 (m, 3H), 6.92 (d, 2H, J = 8.4), 5.6 (s, 1H), 5.05 (s, 2H), 4.21-4.27 (m, 1H), 3.27-3.22 (m, 1H), 2.89-2.81 (m, 4H), 1.72 (s, 3H), 1.19 (t, 3H, J = 7.5); low resolution MS (ES<sup>+</sup>)m/e 594.96 (MH<sup>+</sup>).

5 **Example 71: (2S)-3-{4-[(5-ethyl-2-phenyl-1,3-oxazol-4-yl)methoxy]phenyl}-2-[(Z)-3-oxo-3-phenyl-1-(trifluoromethyl)-1-propenyl]amino}propanoic acid**

10 The title compound was prepared (as described above for the preparation of Example 58) from 100 mg (0.27 mmol) of Intermediate 61, 91 mg (0.41 mmol) of Intermediate 44, and 0.14 mL diisopropylethylamine in 2 mL MeOH. Attempted recrystallization from hot CH<sub>3</sub>CN without success.

15 Silica gel chromatography with MeOH/DCM (10%-20%) gave 91 mg (59% yield) of Example 71: TLC (DCM/MeOH, 4/1): R<sub>f</sub> = 0.57; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 300 MHz) δ 10.96 (br s, 1H), 7.94-7.89 (m, 4H), 7.58-7.44 (m, 6H), 7.12 (d, 2H, J = 8.4), 6.89 (d, 2H, J = 8.1), 6.16 (s, 1H), 4.93 (s, 2H), 4.20-4.11 (m, 1H), 3.16-3.11 (m, 1H), 2.96-2.88 (m, 1H), 2.77 (q, 2H, J = 7.5), 1.17 (t, 3H, J = 7.5); low resolution MS (ES<sup>+</sup>) m/e 564.88 (MH<sup>+</sup>).

25 **Example 72: (2S)-3-{4-[(5-ethyl-2-(4-fluorophenyl)-1,3-oxazol-4-yl)methoxy]phenyl}-2-[(Z)-3-oxo-3-phenyl-1-(trifluoromethyl)-1-propenyl]amino}propanoic acid**

30 The title compound was prepared (as described above for the preparation of Example 58) from 700 mg (1.82 mmol) of Intermediate 63, 610 mg (2.73 mmol) of Intermediate 44, and 0.95 mL diisopropylethylamine in 12 mL MeOH. Silica gel chromatography with MeOH/DCM (10%-20%) gave 580 mg (55% yield) of Example 72: TLC (DCM/MeOH, 4/1): R<sub>f</sub> = 0.50; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 300 MHz) δ 10.95 (br s, 1H), 7.99-7.89 (m, 4H), 7.58-7.53 (m, 3H), 7.34 (t, 2H, J = 8.7), 7.12 (d, 2H, J = 8.1), 6.89 (d, 2H, J = 8.7), 6.16 (s, 1H), 4.92 (s, 2H), 4.21-4.11 (m, 1H), 3.16-3.11 (m, 2H), 2.96-2.91 (m, 1H), 2.79 (q, 2H, J = 7.5), 1.17 (t, 3H, J = 7.5); low resolution MS (ES<sup>+</sup>) m/e 582.92 (MH<sup>+</sup>).

45 **Example 73: (2S)-2-[(Z)-1-ethyl-3-oxo-3-{4-(trifluoromethyl)phenyl}-1-propenyl]amino}-3-{4-[(5-ethyl-2-phenyl-1,3-oxazol-4-yl)methoxy]phenyl}propanoic acid**

5 The title compound was prepared (as described above for the  
preparation of Example 2) from 100 mg (0.27 mmol) of Intermediate 61, 67  
mg (0.27 mmol) of Intermediate 25, and 0.08 mL triethylamine in MeOH (2.0  
10 mL) and trimethylorthoformate (0.5 mL) to yield 54 mg (51% yield) of Example  
73 after silica gel chromatography (5%-15% MeOH in DCM): TLC  
(DCM/MeOH, 4/1):  $R_f$  = 0.54;  $^1\text{H}$  NMR (DMSO- $d_6$ , 300 MHz)  $\delta$  11.63 (d, 1H,  $J$   
=9.3), 7.98-7.90 (m, 4H), 7.73 (d, 2H,  $J$ =8.1), 7.50-7.48 (m, 3H), 7.16 (d, 2H,  
15  $J$ =8.7), 6.90 (d, 2H,  $J$ =8.7), 5.6 (s, 1H), 4.93 (s, 2H), 4.24-4.18 (m, 1H), 3.24-  
3.19 (m, 1H), 2.84-2.72 (m, 4H), 2.08-1.98 (m, 2H), 1.16 (t, 3H,  $J$ =7.8), 0.91  
10 (t, 3H,  $J$ =7.5); low resolution MS ( $\text{ES}^+$ )  $m/e$  593.10 ( $\text{MH}^+$ ).

20  
**Example 74: (2S)-3-{4-[[5-ethyl-2-(4-fluorophenyl)-1,3-oxazol-4-  
yl]methoxy]phenyl}-2-[(Z)-1-ethyl-3-oxo-3-[4-(trifluoromethyl)phenyl]-1-  
propenyl]amino}propanoic acid**

25 The title compound was prepared (as described above for the  
preparation of Example 2) from 160 mg (0.42 mmol) of Intermediate 63, 102  
mg (0.42 mmol) of Intermediate 25, and 0.12 mL triethylamine in MeOH (3.0  
30 mL) and trimethylorthoformate (0.75 mL) to yield 117 mg (46% yield) of  
Example 74 after silica gel chromatography (5%-15% MeOH in DCM): TLC  
(DCM/MeOH, 4/1):  $R_f$  = 0.57;  $^1\text{H}$  NMR (DMSO- $d_6$ , 300 MHz)  $\delta$  11.63 (d, 1H,  $J$   
=9.6), 7.99-7.93 (m, 4H), 7.72 (d, 2H,  $J$ =8.1), 7.32 (t, 2H,  $J$ =8.7), 7.16 (d, 2H,  
35  $J$ =8.7), 6.90 (d, 2H,  $J$ =8.1), 5.6 (s, 1H), 4.91 (s, 2H), 4.26-4.20 (m, 1H), 3.24-  
3.19 (m, 1H), 2.84-2.72 (m, 4H), 2.07-2.00 (m, 2H), 1.15 (t, 3H,  $J$ =7.5), 0.91  
25 (t, 3H,  $J$ =7.5); low resolution MS ( $\text{ES}^+$ )  $m/e$  611.10 ( $\text{MH}^+$ ).

40  
**Example 75: (2S)-3-{4-[[5-ethyl-2-phenyl-1,3-oxazol-4-  
yl]methoxy]phenyl}-2-[(Z)-1-methyl-3-oxo-3-[4-(trifluoromethyl)phenyl]-  
1-propenyl]amino}propanoic acid**

45 The title compound was prepared (as described above for the  
preparation of Example 2) from 98 mg (0.27 mmol) of Intermediate 61, 62 mg  
50 (0.27 mmol) of Intermediate 24, and 0.08 mL triethylamine in MeOH (2.0 mL)

and trimethylorthoformate (0.5 mL) to yield 95 mg (61% yield) of Example 75 after silica gel chromatography (5%-15% MeOH in DCM): TLC (DCM/MeOH, 4/1):  $R_f$  = 0.59;  $^1\text{H}$  NMR (DMSO- $d_6$ , 300 MHz)  $\delta$  11.55 (d, 1H,  $J$  = 9.0), 7.98-7.90 (m, 4H), 7.73 (d, 2H,  $J$  = 8.4), 7.51-7.48 (m, 3H), 7.17 (d, 2H,  $J$  = 8.4), 6.91 (d, 2H,  $J$  = 8.4), 5.61 (s, 1H), 4.93 (s, 2H), 4.24-4.18 (m, 1H), 3.23-3.16 (m, 1H), 2.86-2.73 (m, 4H), 1.72 (s, 3H), 1.17 (t, 3H,  $J$  = 7.5); low resolution MS ( $\text{ES}^+$ )  $m/e$  579.05 ( $\text{MH}^+$ ).

**Example 76: (2S)-3-{4-[[5-ethyl-2-(4-fluorophenyl)-1,3-oxazol-4-yl]methoxy]phenyl}-2-[[{(Z)-1-methyl-3-oxo-3-[4-(trifluoromethyl)phenyl]-1-propenyl]amino}propanoic acid**

The title compound was prepared (as described above for the preparation of Example 2) from 176 mg (0.46 mmol) of Intermediate 63, 105 mg (0.46 mmol) of Intermediate 24, and 0.14 mL triethylamine in MeOH (3.0 mL) and trimethylorthoformate (0.75 mL) to yield 152 mg (56% yield) of Example 76 after silica gel chromatography (5%-15% MeOH in DCM): TLC (DCM/MeOH, 4/1):  $R_f$  = 0.55;  $^1\text{H}$  NMR (DMSO- $d_6$ , 300 MHz)  $\delta$  11.55 (d, 1H,  $J$  = 9.0), 7.99-7.94 (m, 4H), 7.73 (d, 2H,  $J$  = 8.4), 7.33 (t, 2H,  $J$  = 8.7), 7.16 (d, 2H,  $J$  = 8.4), 6.90 (d, 2H,  $J$  = 8.7), 5.61 (s, 1H), 4.93 (s, 2H), 4.23-4.18 (m, 1H), 3.22-3.18 (m, 1H), 2.85-2.73 (m, 4H), 1.72 (s, 3H), 1.17 (t, 3H,  $J$  = 7.5); low resolution MS ( $\text{ES}^+$ )  $m/e$  597.09 ( $\text{MH}^+$ ).

**Example 77: (2S)-2-[[{(Z)-1-ethyl-3-oxo-3-[4-(trifluoromethyl)phenyl]-1-propenyl]amino}-3-[4-[[5-ethyl-2-phenyl-1,3-thiazol-4-yl]methoxy]phenyl]propanoic acid**

The title compound was prepared (as described above for the preparation of Example 2) from 1.44 g (2.90 mmol) of Intermediate 62, 710 mg (2.90 mmol) of Intermediate 25, and 0.85 mL triethylamine in MeOH (20 mL) and trimethylorthoformate (5 mL) to yield 985 mg (56% yield) of Example 77 after silica gel chromatography (5%-15% MeOH in DCM): TLC (DCM/MeOH, 4/1):  $R_f$  = 0.45;  $^1\text{H}$  NMR (DMSO- $d_6$ , 300 MHz)  $\delta$  11.65 (d, 1H,  $J$



=9.3), 7.97 (d, 2H,  $J=8.1$ ), 7.85 (m, 2H), 7.71 (d, 2H,  $J=8.1$ ), 7.44 (m, 3H), 7.44 (d, 2H,  $J=8.1$ ), 7.16 (d, 2H,  $J=8.1$ ), 5.61 (s, 1H), 5.05 (s, 2H), 4.26-4.19 (m, 1H), 3.26-3.21 (m, 1H), 2.89-2.77 (m, 4H), 2.07-2.0 (m, 2H), 1.19 (t, 3H,  $J=7.5$ ), 0.91 (t, 3H,  $J=7.5$ ); low resolution MS ( $ES^+$ )  $m/e$  609.07 ( $MH^+$ ).

**Example 78: (2S)-3-(4-{2-[5-ethyl-2-(4-fluorophenyl)-1,3-oxazol-4-yl]ethoxy}phenyl)-2-(((Z)-1-ethyl-3-oxo-3-[4-(trifluoromethyl)phenyl]-1-propenyl)amino)propanoic acid**

The title compound was prepared (as described above for the preparation of Example 2) from 150 mg (0.29 mmol) of Intermediate 58 (TFA Salt), 108 mg (0.3 mmol) of Intermediate 25, and 0.1 mL triethylamine in MeOH and trimethylorthoformate to yield 70 mg (38% yield) of Example 78: TLC (DCM/MeOH, 90/10):  $R_f = 0.26$ ;  $^1H$  NMR (DMSO- $d_6$ , 300 MHz)  $\delta$  11.62 (d, 1H,  $J=9.3$ ), 7.99-7.93 (m, 4H), 7.74 (d, 2H,  $J=8.4$ ), 7.32 (t, 2H,  $J=8.9$ ), 7.13 (d, 2H,  $J=8.4$ ), 6.80 (d, 2H,  $J=8.5$ ), 5.61 (s, 1H), 4.14 (m, 3H), 3.4-3.1 (m, 2H, obscured by water peak), 2.88 (t, 2H,  $J=6.4$ ), 2.70 (q, 2H,  $J=7.4$ ), 2.02 (m, 2H), 1.19 (t, 3H,  $J=7.5$ ), 0.94 (t, 3H,  $J=7.5$ ); low resolution MS ( $ES^+$ )  $m/e$  625.2 ( $MH^+$ ).

**Example 79: (2S)-3-(4-{2-[5-ethyl-2-(4-fluorophenyl)-1,3-thiazol-4-yl]ethoxy}phenyl)-2-(((Z)-1-ethyl-3-oxo-3-[4-(trifluoromethyl)phenyl]-1-propenyl)amino)propanoic acid**

The title compound was prepared (as described above for the preparation of Example 2) from 150 mg (0.28 mmol) of Intermediate 56 (TFA Salt), 108 mg (0.3 mmol) of Intermediate 25, and 0.1 mL triethylamine in MeOH and trimethylorthoformate to yield 60 mg (33% yield) of Example 79: TLC (DCM/MeOH, 90/10):  $R_f = 0.26$ ;  $^1H$  NMR (DMSO- $d_6$ , 300 MHz)  $\delta$  11.63 (d, 1H,  $J=9.4$ ), 7.98 (d, 2H,  $J=8.0$ ), 7.9-7.85 (m, 2H), 7.74 (d, 2H,  $J=8.3$ ), 7.27 (t, 2H,  $J=8.7$ ), 7.13 (d, 2H,  $J=8.4$ ), 6.80 (d, 2H,  $J=8.3$ ), 5.62 (s, 1H), 4.25 (m, 3H), 3.4-3.2 (m, 2H, obscured by water peak), 3.07 (t, 2H,  $J=6.3$ ).

2.85-2.77 (m, 2H), 2.06-1.99 (m, 2H), 1.21 (t, 3H,  $J = 7.4$ ), 0.93 (t, 3H,  $J = 7.4$ );  
low resolution MS ( $ES^+$ )  $m/e$  641.12 ( $MH^+$ ).

**Example 80: (2S)-2-((Z)-1-ethyl-3-oxo-3-(4-(trifluoromethyl)phenyl)-1-propenyl)amino)-3-(4-(2-(5-ethyl-2-phenyl-1,3-thiazol-4-yl)ethoxy)phenyl)propanoic acid**

The title compound was prepared (as described above for the preparation of Example 2) from 150 mg (0.29 mmol) of Intermediate 57 (TFA Salt), 108 mg (0.3 mmol) of Intermediate 25, and 0.1 mL triethylamine in MeOH and trimethylorthoformate to yield 40 mg (22% yield) of Example 80: TLC (DCM/MeOH, 90/10):  $R_f = 0.28$ ;  $^1H$  NMR (DMSO- $d_6$ , 300 MHz)  $\delta$  11.62 (d, 1H,  $J = 9.4$ ), 7.98 (d, 2H,  $J = 8.2$ ), 7.85 (m, 2H), 7.74 (d, 2H,  $J = 8.3$ ), 7.48-7.42 (m, 3H), 7.13 (d, 2H,  $J = 8.4$ ), 6.80 (d, 2H,  $J = 8.3$ ), 5.63 (s, 1H), 4.23 (t, 2H,  $J = 6.5$ ), 4.2 (m, 1H), 3.2-3.05 (m, 3H), 2.89-2.75 (m, 3H), 2.07-2.03 (m, 2H), 1.23 (t, 3H,  $J = 7.5$ ), 0.94 (t, 3H,  $J = 7.5$ ); low resolution MS ( $ES^+$ )  $m/e$  622.91 ( $MH^+$ ).

The following Intermediates A-G were prepared to make radioligand for the binding assay described below.

**Intermediate A: 2-(4-(2-(Phenylmethyloxycarbonylamino)ethyl)phenoxy)-2-methylbutanoic acid**

A solution of 4-(2-(phenylmethyloxycarbonylamino)ethyl)phenol (5.74 g; 21.16 mmole) in 2-butanone (17 mL) and chloroform (6 g) was added dropwise to a mixture of sodium hydroxide (9.0 g; 225 mmole) and 2-butanone (67 mL) whilst keeping the reaction temperature below 30°C. The mixture was allowed to stir at 30°C for 4h. Ether (100 mL) was added and the resultant solid was collected by filtration and washed with ether (100 mL). The solid was dissolved in water (70 mL) and any residual ether removed by evaporation. 1N Hydrochloric acid was added to adjust the pH to 1, and the resulting oil was extracted with dichloromethane (3 x 50 mL). The combined extracts were dried ( $Na_2SO_4$ ) and evaporated to afford a yellow oil (3.82 g; 49%).  $^1H$ -NMR ( $CDCl_3$ )  $\delta$  7.26 (s, 5H), 7.09 (d, 2H,  $J = 7.9$  Hz), 6.88 (d, 2H,  $J = 8.4$  Hz), 5.09 (s, 2H), 4.75 (br s, 1H), 3.42-3.44 (m, 2H), 2.75 (t, 2H,  $J = 6.7$

Hz), 1.92-2.00 (m, 2H), 1.47 (s, 3H), 1.04 (t, 3H, J=2.6 Hz). Mass spectrometry ES<sup>+</sup>, m/e (M+H)<sup>+</sup>=372.

**Intermediate B: Methyl 2-(4-(2-**

**(phenylmethyloxycarbonylamino)ethyl)phenoxy)-2-methyl butyrate**

A solution of Intermediate A (2.0 g; 5.38 mmole) in dimethylformamide (12 mL) was treated with potassium carbonate (2.23 g; 16.14 mmole) and methyl iodide (1.54 g; 10.76 mmole) and the resulting mixture stirred at 23°C for 2h. The mixture was filtered and the solid collected was washed with ethyl acetate (70 mL). The filtrate was washed with brine (4 x 50 mL), dried (Na<sub>2</sub>SO<sub>4</sub>) and evaporated. The residue was purified by chromatography on silica gel using hexane then 33% ethyl acetate-hexane as eluent to afford a colorless oil (1.27 g; 61%).

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ 7.31 (m, 5H), 7.06 (d, 2H, J=8.4 Hz), 6.68 (d, 2H, J=8.4 Hz), 4.98 (s, 2H), 3.67 (s, 3H), 3.15 (m, 2H), 2.62 (t, 2H, J=7.1 Hz), 1.86 (m, 2H), 1.38 (s, 3H), 0.86 (t, 3H, J=7.3 Hz). Mass spectrometry ES<sup>+</sup>, m/e (M+Na)<sup>+</sup> = 408.

**Intermediate C: Methyl 2-(4-(2-aminoethyl)phenoxy)-2-methyl butyrate acetate salt**

A solution of Intermediate B (1.27 g; 3.29 mmole) in methanol (50 mL) and acetic acid (0.4 g) was treated with 10% palladium on carbon and shaken in a hydrogen atmosphere (50 psi) for 2h. The catalyst was filtered through celite and the solvent was evaporated to afford a yellow oil in quantitative yield (1.04 g).

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 7.06 (d, 2H, J=8.4 Hz), 6.77 (d, 2H, J=8.4 Hz), 6.70 (br s, 2H), 3.76 (s, 3H), 3.02 (br s, 2H), 2.82 (m, 2H), 1.99 (s, 3H), 1.92 (m, 2H), 1.48 (s, 3H), 0.96 (t, 3H, J=7.4 Hz). Mass spectrometry ES<sup>+</sup>, m/e (M+H)<sup>+</sup>=252.

**Intermediate D: Methyl 2-(4-(2-(2,4-dinitrophenylsulfonylamino)ethyl)phenoxy)-2-methyl butyrate**

A solution of Intermediate C (2 g; 6.42 mmole) in CH<sub>2</sub>Cl<sub>2</sub> (40 mL) was treated with saturated sodium bicarbonate solution and the organic layer was separated. The aqueous layer was extracted with CH<sub>2</sub>Cl<sub>2</sub> (5 x 50 mL) and the combined organic layers were dried (Na<sub>2</sub>SO<sub>4</sub>) and evaporated to afford the free base as a yellow oil (1.61 g; 100%). This was dissolved in CH<sub>2</sub>Cl<sub>2</sub> (40 mL) and treated with pyridine

(0.45 g; 5.61 mmole) and 2,4-dinitrophenylsulfonyl chloride (1.5 g; 5.61 mmole), and the mixture was allowed to stir at 23°C for 3h. Water (60 mL) was added and the organic layer separated, washed with water (3 X 40 mL) and saturated sodium bicarbonate (40 mL). The organic layer was dried (Na<sub>2</sub>SO<sub>4</sub>) and evaporated and the residue purified by chromatography using 15-20% EtOAc-Hexane as eluent to afford a light yellow solid (1.38 g; 51%). <sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 8.63 (d, 1H, J=2.3 Hz), 8.49 (dd, 1H, J=8.4 Hz, J'=2.3 Hz), 8.07 (d, 1H, J=8.4 Hz), 6.89 (d, 2H, J=8.4 Hz), 6.54 (d, 2H, J=8.4 Hz), 5.34 (t, 1H, J=5.3 Hz), 3.78 (s, 3H), 3.48 (q, 2H, J=8.3 Hz), 2.75 (t, 2H, J=6.6 Hz), 1.92 (m, 2H), 1.42 (s, 3H), 0.93 (t, 3H, J=7.5 Hz).

**Intermediate E: Methyl 2-(4-(2-((2,4-dinitrophenylsulfonyl)(hept-2-en-1-yl))amino)ethyl)phenoxy)-2-methyl butyrate**

A solution of Intermediate D (315 mg; 0.654 mmole) in THF (15 mL) was treated with triphenylphosphine (343 mg; 1.308 mmole), hept-2-en-1-ol (150 mg; 1.308 mmole) and diethylazodicarboxylate (228 mg; 1.308 mmole) and the mixture allowed to stir at 23°C for 1h. The solvent was evaporated and the residue purified by chromatography using 10-15% EtOAc-Hexane as eluent to afford a semi-solid (400 mg; >100%). TLC and NMR shows that the desired compound is present along with 1,2-(diethoxycarbonyl)hydrazine.

**Intermediate F: Methyl 2-(4-(2-(hept-2-en-1-ylamino)ethyl)phenoxy)-2-methyl butanoate**

A solution of Intermediate E (400 mg; 0.654 mmole) in CH<sub>2</sub>Cl<sub>2</sub> (5 mL) was treated with triethylamine (132 mg; 1.308 mmole) and mercaptoacetic acid (78 mg; 0.85 mmole) and the mixture was allowed to stir at 23°C for 1h. The mixture was diluted with EtOAc (30 mL) and washed with water (3 X 20 mL) and aqueous sodium bicarbonate (30 mL). The organic layer was dried (Na<sub>2</sub>SO<sub>4</sub>), evaporated and the residue purified by chromatography using 10% EtOAc-Hexane then 50% EtOAc-Hexane then MeOH as eluent to afford an oil (177 mg; 78% from intermediate 24). <sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 7.06 (d, 2H, J=7.5 Hz), 6.75 (d, 2H, J=7.5 Hz), 5.59 (m, 2H), 3.76 (s, 3H), 3.30 (d, 2H, J=6.3 Hz), 2.87 (m, 4H), 1.96 (m, 4H), 1.47 (s, 3H), 1.28 (m, 5H), 0.96 (t, 3H, J=7.6 Hz), 0.86 (t, 3H, J=6.9 Hz).

**Intermediate G: Methyl 2-(4-(2-(1-hept-2-enyl-3-(2,4-difluorophenyl)ureido)ethyl)phenoxy)-2-methylbutyrate**

A solution of Intermediate F (157 mg; 0.452 mmole) in methylene chloride (5 mL) was treated with 2,4-difluorophenylisocyanate (140 mg; 0.904 mmole) and the mixture allowed to stand at 23°C for 18h. The solvent was evaporated and the residue purified by chromatography on silica gel using 10% then 15% ethyl acetate-hexane as eluent to afford a yellow semi-solid (212 mg; 93%). Contaminated with bis-(2,4-difluorophenyl)urea which co-elutes on column.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 8.85 (br s, 1H), 8.02 (m, 1H), 7.09 (d, 2H, J=8.4 Hz), 6.77-6.90 (m, 4H), 5.70 (m, 1H), 5.36 (m, 1H), 3.76 (s, 3H), 3.54 (t, 2H, J=7.3 Hz), 2.84 (t, 2H, J=7.1 Hz), 1.55 (br s, 1H), 1.46 (s, 3H), 1.25-1.35 (m, 5H), 0.96 (t, 3H, J=7.3 Hz), 0.88 (t, 3H, J=7.4 Hz). Mass spectrometry CI/AP<sup>+</sup>, m/e (M+H)<sup>+</sup>=503.

**2-(4-(2-(1-Hept-2-enyl-3-(2,4-difluorophenyl)ureido)ethyl) phenoxy)-2-methylbutanoic acid (Radioligand Precursor)**

A solution of Intermediate G (370 mg; 0.736 mmole) in methanol (15 mL) was treated with 1N NaOH (7.5 mL) and the mixture heated under reflux for 2h. The mixture was acidified with 1N HCl and extracted with ethyl acetate (3 x 25 mL). The combined organic layers were washed with brine, dried (Na<sub>2</sub>SO<sub>4</sub>) and evaporated. The residue was purified by chromatography on silica gel using 20% ethyl acetate-hexane then ethyl acetate as eluent to afford a tan oil (280 mg; 78%).

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ 7.95-8.09 (m, 1H), 7.14 (d, 2H, J=7.1 Hz), 6.90 (d, 2H, J=7.4 Hz), 6.81 (d, 2H, J=5.2 Hz), 5.66 (m, 1H), 5.37 (m, 1H), 3.56 (t, 2H, J=7.4 Hz), 2.87 (t, 2H, J=7.4 Hz), 2.00 (m, 4H), 1.44 (s, 3H), 1.27 (m, 6H), 1.03 (t, 3H, J=7.3 Hz), 0.88 (t, 3H, J=7.3 Hz). Mass spectrometry ES<sup>-</sup>, m/e (M+H)<sup>+</sup> = 489.

**Radioligand: 2-(4-(2-(2,3-Ditritio-1-heptyl-3-(2,4-difluorophenyl)ureido)ethyl) phenoxy)-2-methylbutanoic acid**

A solution of radioligand precursor prepared above (10 mg) in anhydrous DMF (3.5 mL) was transferred to a reaction vessel containing 10 % Pd/C (9.8 mg). The reaction vessel was evacuated and degassed via one freeze-thaw-evacuation cycle and then exposed to tritium gas (10.1 Ci). After 4h, the mixture was filtered through celite, evaporated and the residue

5 dissolved in acetonitrile. A portion of this solution (0.8 mL, 26.6 mCi) was  
purified by HPLC (Dynamax C8, 25 min gradient from 4:1  
acetonitrile:0.1%TFA to 9:1 acetonitrile: 0.1% TFA, 235 nm). Fractions  
10 containing pure material were combined and evaporated under nitrogen. The  
5 residue was redissolved in acetonitrile to provide a solution of the title  
compound (82.0 Ci/mmol, radiochemical purity, 99%).

15 **2-(4-(2-(1-Heptyl-3-(2,4-difluorophenyl)ureido)ethyl) phenoxy)-2-  
methylbutanoic acid**

10 The unlabelled ("cold") version of the above radioligand was prepared  
as a control. A solution of Intermediate G (10 mg) in anhydrous DMF (3.5 mL)  
20 was transferred to a reaction vessel containing 10 % Pd/C (9.8 mg). The  
reaction vessel was evacuated and degassed via one freeze-thaw-evacuation  
cycle and then exposed to hydrogen gas. After 4h, the mixture was filtered  
25 15 through celite and evaporated. The residue was purified by chromatography  
using 2% MeOH/CH<sub>2</sub>Cl<sub>2</sub> as eluent to afford a gum (7mg).

30 The radioligand prepared above was used in the PPAR $\alpha$  binding assay  
described below (in addition to <sup>3</sup>H-BRL 49653 for the PPAR $\gamma$  binding assay)  
20 to show that active compounds in the transfection assays were also ligands  
for PPAR $\alpha$  and PPAR $\gamma$ .

35 **Binding Assay:**

Compounds were tested for their ability to bind to hPPAR $\gamma$  or hPPAR $\alpha$   
25 using a Scintillation Proximity Assay (SPA). The PPAR ligand binding domain  
40 (LBD) was expressed in *E. coli* as polyHis tagged fusion proteins and purified.  
The LBD was then labeled with biotin and immobilized on streptavidin-  
modified scintillation proximity beads. The beads were then incubated with a  
45 constant amount of the appropriate radioligand (<sup>3</sup>H-BRL 49653 for PPAR $\gamma$  and  
30 2-(4-(2-(2,3-Ditritio-1-heptyl-3-(2,4-difluorophenyl)ureido)ethyl)phenoxy)-2-  
methylbutanoic acid for hPPAR $\alpha$ ) and variable concentrations of test  
50 compound, and after equilibration the radioactivity bound to the beads was

5 measured by a scintillation counter. The amount of nonspecific binding, as  
assessed by control wells containing 50  $\mu$ M of the corresponding unlabeled  
ligand, was subtracted from each data point. For each compound tested,  
10 plots of ligand concentration vs. CPM of radioligand bound were constructed  
and apparent  $K_i$  values were estimated from nonlinear least squares fit of the  
data assuming simple competitive binding. The details of this assay have  
15 been reported elsewhere (see, Blanchard, S. G. et. al. Development of a  
Scintillation Proximity Assay for Peroxisome Proliferator-Activated Receptor  
gamma Ligand Binding Domain. *Anal. Biochem.* **1998**, 257, 112-119).

10 Preferably, the compounds of this invention bind to both hPPAR $\gamma$  and  
hPPAR $\alpha$ . All of the above Examples 1-80 did bind to both hPPAR $\gamma$  and  
hPPAR $\alpha$ . Apparent  $pK_i$  values were  $>6.3$  for all Examples 1-80 in both the  
PPAR $\gamma$  and PPAR $\alpha$  binding assays described above ( $pK_i = -\log$  of the  
25 concentration of test compound required to achieve an apparent  $K_i$  value  
according to the equation  $K_i = IC_{50} / (1 + [L]/K_d)$ , where  $IC_{50}$  = the concentration  
of test compound required to inhibit 50% of the specific binding of the  
radioligand,  $[L]$  is the concentration of the radioligand used, and  $K_d$  is the  
30 dissociation constant for the radioligand at the receptor).

#### 20 Transfection assay:

35 Compounds were screened for functional potency in transient  
transfection assays in CV-1 cells for their ability to activate the PPAR  
subtypes (transactivation assay). A previously established chimeric receptor  
25 system was utilized to allow comparison of the relative transcriptional activity  
of the receptor subtypes on the same target gene and to prevent endogenous  
receptor activation from complicating the interpretation of results. See, for  
example, Lehmann, J. M.; Moore, L. B.; Smith-Oliver, T. A.; Wilkison, W. O.;  
45 Willson, T. M.; Kliewer, S. A., An antidiabetic thiazolidinedione is a high  
affinity ligand for peroxisome proliferator-activated receptor  $\gamma$  (PPAR $\gamma$ ), *J. Biol.*  
30 *Chem.*, **1995**, 270, 12953-6. The ligand binding domains for murine and  
human PPAR $\alpha$ , and PPAR $\gamma$ , were each fused to the yeast transcription factor  
50

5 GAL4 DNA binding domain. CV-1 cells were transiently transfected with  
expression vectors for the respective PPAR chimera along with a reporter  
construct containing five copies of the GAL4 DNA binding site driving  
10 expression of secreted placental alkaline phosphatase (SPAP) and  $\beta$ -  
5 galactosidase. After 16 h, the medium was exchanged to DME medium  
supplemented with 10% delipidated fetal calf serum and the test compound at  
the appropriate concentration. After an additional 24 h, cell extracts were  
15 prepared and assayed for alkaline phosphatase and  $\beta$ -galactosidase activity.  
Alkaline phosphatase activity was corrected for transfection efficiency using  
10 the  $\beta$ -galactosidase activity as an internal standard (see, for example, Kliewer,  
S. A., et. al. *Cell* 83, 813-819 (1995)). Rosiglitazone (BRL 49653) was used  
20 as a positive control in the hPPAR $\gamma$  assay. The positive control in the  
hPPAR $\alpha$  assay was 2-[4-(2-(3-(4-fluorophenyl)-1-heptylureido)ethyl)-  
phenoxy]-2-methylpropionic acid, which can be prepared as described in  
25 Brown, Peter J., et. al. *Synthesis* Issue 7, 778-782 (1997), or patent  
15 publication WO 9736579.

30 All of the above Examples showed at least 50% activation of hPPAR $\alpha$   
and hPPAR $\gamma$  at concentrations of  $10^{-7}$  M or less.

20 **In vivo evaluation:**

35 Male Zucker Diabetic Fatty rats were lightly anesthetized with  
isoflurane gas and bled by tail vein to obtain postprandial baseline  
concentrations for plasma glucose, serum lipids and insulin. Animals were  
baseline matched by plasma glucose and randomized into vehicle or  
40 25 treatment groups with compound administration by oral gavage, b.i.d.,  
beginning at 8.5 weeks of age. Selected compounds were administered as a  
suspension in 0.5% methylcellulose for 7 consecutive days. Changes in body  
weight and food consumption were monitored from representative animals  
45 from each group, over a 48 hr period. After 7 days of treatment animals were  
30 anesthetized, blood samples obtained and analyzed for plasma glucose and  
lactate, serum triglyceride, total and HDL cholesterol, non-esterified free fatty  
acids and insulin concentration. Livers were weighed and corrected for body  
50



5

weight. Values in Table 1 for % Glucose reduction represent a summary of the percent reduction from vehicle control animals at day 7 relative to normalization defined in this model as plasma glucose levels of 140 mg/dL.

10

15

20

25

30

35

40

45

50

55

5

Table 1. Biological activity

		<u>Example</u>	<u>% Glucose Reduction</u>
10	5	1	39
		7	34
		12	61
		16	59
15	10	20	50
		21	82
		22	49
20	15	23	86
		24	45
		40	79
25	20	48	81

25

30

35

40

45

50

55

## Claims

5

10

15

20

25

30

35

40


45

50

55

What is claimed is:

1. A compound of formula (I), or a tautomeric form, pharmaceutically acceptable salt, or solvate thereof,



(I)

wherein;

R<sup>1</sup> is hydrogen or C<sub>1-3</sub>alkyl;

R<sup>2</sup> is hydrogen, or C<sub>1-6</sub>alkyl optionally substituted by one or more halogens;

R<sup>3</sup> is C<sub>1-6</sub>alkyl, C<sub>4-7</sub>cycloalkyl or cycloalkenyl, -OC<sub>1-6</sub>alkyl, -NR'R' (where each R' is independently hydrogen or C<sub>1-3</sub>alkyl), a 5 or 6 membered heterocyclic group containing at least one oxygen, nitrogen, or sulfur ring atom (optionally substituted by one or more halogen, C<sub>1-6</sub>alkyl optionally substituted by one or more halogens, -OC<sub>1-6</sub>alkyl optionally substituted by one or more halogens, -CN, or -NO<sub>2</sub>), or phenyl (optionally substituted by one or more halogen, C<sub>1-6</sub>alkyl optionally substituted by one or more halogens, -OC<sub>1-6</sub>alkyl optionally substituted by one or more halogens, -CN, or -NO<sub>2</sub>);

R<sup>4</sup> is a 5 or 6 membered heterocyclic group containing at least one oxygen, nitrogen, or sulfur ring atom (optionally substituted by one or more halogen, C<sub>1-6</sub>alkyl optionally substituted by one or more halogens, -OC<sub>1-6</sub>alkyl optionally substituted by one or more halogens, -CN, or -NO<sub>2</sub>), or phenyl (optionally substituted by one or more halogen, C<sub>1-6</sub>alkyl optionally substituted by one or more halogens, -OC<sub>1-6</sub>alkyl optionally substituted by one or more halogens, -NR'R', -CN, or -NO<sub>2</sub>);

5           R<sup>5</sup> is hydrogen, halogen, or C<sub>1-3</sub>alkyl optionally substituted by one or more halogens;

          R<sup>6</sup> is hydrogen or C<sub>1-3</sub>alkyl;

10          X is O or S; and

5          n is 1, 2, or 3.

15           2. A compound of Claim 1 wherein R<sup>1</sup> is hydrogen or methyl.

          3. A compound of any of Claims 1-2 wherein R<sup>2</sup> is C<sub>1-8</sub>alkyl optionally substituted by one or more halogens.

20           4. A compound of any of Claims 1-3 wherein R<sup>3</sup> is, pyridine, pyrazine, thiophene, furan, thiazole, or phenyl (any of which may be optionally substituted by one or more halogen, C<sub>1-6</sub>alkyl optionally substituted by one or more halogens, -OC<sub>1-6</sub>alkyl optionally substituted by one or more halogens, -CN, or -NO<sub>2</sub>) or C<sub>4-7</sub>cycloalkyl.

30           5. A compound of Claim 4 wherein R<sup>3</sup> is phenyl (optionally substituted by one or more halogen, C<sub>1-6</sub>alkyl optionally substituted by one or more halogens, -OC<sub>1-6</sub>alkyl optionally substituted by one or more halogens, -CN, or -NO<sub>2</sub>).

35           6. A compound of any of Claims 1-5 wherein R<sup>4</sup> is phenyl (optionally substituted by one or more halogen, C<sub>1-6</sub>alkyl optionally substituted by one or more halogens, or -OC<sub>1-6</sub>alkyl optionally substituted by one or more halogens).

40           7. A compound of any of Claims 1-6 wherein R<sup>5</sup> is hydrogen, or C<sub>1-3</sub>alkyl optionally substituted by one or more halogens.

45           8. A compound of Claim 7 wherein R<sup>5</sup> is hydrogen.

50           9. A compound of any of Claims 1-8 wherein R<sup>6</sup> is methyl or ethyl.

10. A compound of any of Claims 1-9 wherein n is 2.

11. A compound of Claim 1 selected from the group consisting of:

- (2S)-2-(((Z)-1-methyl-3-oxo-3-phenyl-1-propenyl)amino)-3-(4-(2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy)phenyl)propanoic acid
- (2S)-3-(4-(2-(2-(4-fluorophenyl)-5-methyl-1,3-oxazol-4-yl)ethoxy)phenyl)-2-(((Z)-1-methyl-3-oxo-3-(4-(trifluoromethyl)phenyl)-1-propenyl)amino)propanoic acid
- (2S)-3-(4-(2-(2-(4-isopropoxyphenyl)-5-methyl-1,3-oxazol-4-yl)ethoxy)phenyl)-2-(((Z)-1-methyl-3-oxo-3-(4-(trifluoromethyl)phenyl)-1-propenyl)amino)propanoic acid
- (2S)-3-(4-(2-(2-(4-methoxyphenyl)-5-methyl-1,3-oxazol-4-yl)ethoxy)phenyl)-2-(((Z)-1-methyl-3-oxo-3-(4-(trifluoromethyl)phenyl)-1-propenyl)amino)propanoic acid
- (2S)-2-(((Z)-1-ethyl-3-oxo-3-phenyl-1-propenyl)amino)-3-(4-(2-(5-methyl-2-(4-(trifluoromethyl)phenyl)-1,3-oxazol-4-yl)ethoxy)phenyl)propanoic acid
- (2S)-2-(((Z)-1-ethyl-3-(4-fluorophenyl)-3-oxo-1-propenyl)amino)-3-(4-(2-(2-(4-methoxyphenyl)-5-methyl-1,3-oxazol-4-yl)ethoxy)phenyl)propanoic acid
- (2S)-2-(((Z)-1-methyl-3-oxo-3-phenyl-1-propenyl)amino)-3-(4-(2-(5-methyl-2-(4-(trifluoromethyl)phenyl)-1,3-oxazol-4-yl)ethoxy)phenyl)propanoic acid
- (2S)-2-(((Z)-1-ethyl-3-oxo-3-phenyl-1-propenyl)amino)-3-(4-(2-(2-(4-fluorophenyl)-5-methyl-1,3-oxazol-4-yl)ethoxy)phenyl)propanoic acid
- (2S)-2-(((Z)-1-ethyl-3-(4-fluorophenyl)-3-oxo-1-propenyl)amino)-3-(4-(2-(2-(4-fluorophenyl)-5-methyl-1,3-oxazol-4-yl)ethoxy)phenyl)propanoic acid
- (2S)-2-(((Z)-1-methyl-3-oxo-3-(4-(trifluoromethyl)phenyl)-1-propenyl)amino)-3-(4-(2-(5-methyl-2-phenyl-1,3-thiazol-4-yl)ethoxy)phenyl)propanoic acid
- (2S)-2-(((Z)-3-(4-fluorophenyl)-1-methyl-3-oxo-1-propenyl)amino)-3-(4-(2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy)phenyl)propanoic acid
- (2S)-2-(((Z)-1-methyl-3-oxo-3-(2,3,4-trifluorophenyl)-1-propenyl)amino)-3-(4-(2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy)phenyl)propanoic acid
- (2S)-2-(((Z)-1-methyl-3-(4-nitrophenyl)-3-oxo-1-propenyl)amino)-3-(4-(2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy)phenyl)propanoic acid

- 5 (2S)-2-(((Z)-1-methyl-3-oxo-3-[4-(trifluoromethyl)phenyl]-1-propenyl)amino)-3-[4-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]phenyl]propanoic acid
- 10 (2S)-2-(((Z)-1-ethyl-3-(4-fluorophenyl)-3-oxo-1-propenyl)amino)-3-[4-[2-[2-(4-isopropoxyphenyl)-5-methyl-1,3-oxazol-4-yl]ethoxy]phenyl]propanoic acid
- 5 (2S)-2-(((Z)-1-methyl-3-oxo-3-(2,4,5-trifluorophenyl)-1-propenyl)amino)-3-[4-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]phenyl]propanoic acid
- 15 (2S)-2-(((Z)-1-ethyl-3-oxo-3-phenyl-1-propenyl)amino)-3-[4-[2-[2-(4-isopropoxyphenyl)-5-methyl-1,3-oxazol-4-yl]ethoxy]phenyl]propanoic acid
- 10 (2S)-2-(((Z)-1-ethyl-3-oxo-3-[4-(trifluoromethyl)phenyl]-1-propenyl)amino)-3-[4-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]phenyl]propanoic acid
- 20 (2S)-2-(((Z)-1-methyl-3-(4-methylphenyl)-3-oxo-1-propenyl)amino)-3-[4-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]phenyl]propanoic acid
- 25 (2S)-2-(((Z)-1-ethyl-3-oxo-3-phenyl-1-propenyl)amino)-3-[4-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]phenyl]propanoic acid
- 15 (2S)-3-[4-[2-[2-(4-fluorophenyl)-5-methyl-1,3-thiazol-4-yl]ethoxy]phenyl]-2-(((Z)-3-oxo-3-phenyl-1-(trifluoromethyl)-1-propenyl)amino)propanoic acid
- 30 (2S)-2-(((Z)-1-ethyl-3-oxo-3-[4-(trifluoromethyl)phenyl]-1-propenyl)amino)-3-[4-[2-[2-(4-fluorophenyl)-5-methyl-1,3-thiazol-4-yl]ethoxy]phenyl]propanoic acid
- 20 (2S)-2-(((Z)-1-butyl-3-oxo-3-phenyl-1-propenyl)amino)-3-[4-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]phenyl]propanoic acid
- 35 (2S)-2-(((Z)-3-(4-chlorophenyl)-1-methyl-3-oxo-1-propenyl)amino)-3-[4-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]phenyl]propanoic acid
- 25 (2S)-2-(((Z)-1-methyl-3-(3-nitrophenyl)-3-oxo-1-propenyl)amino)-3-[4-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]phenyl]propanoic acid
- 40 (2S)-2-(((Z)-3-[2-fluoro-3-(trifluoromethyl)phenyl]-1-methyl-3-oxo-1-propenyl)amino)-3-[4-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]phenyl]propanoic acid
- 45 (2S)-2-(((Z)-3-(4-isopropoxyphenyl)-1-methyl-3-oxo-1-propenyl)amino)-3-[4-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]phenyl]propanoic acid
- 30 (2S)-2-(((Z)-3-(2-chlorophenyl)-1-methyl-3-oxo-1-propenyl)amino)-3-[4-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]phenyl]propanoic acid
- 50
- 55

- 5 (2S)-2-(((Z)-3-(2-furyl)-1-methyl-3-oxo-1-propenyl)amino)-3-{4-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]phenyl}propanoic acid
- (2S)-2-(((Z)-1-methyl-3-oxo-3-(2-pyrazinyl)-1-propenyl)amino)-3-{4-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]phenyl}propanoic acid
- 10 (2S)-2-(((Z)-3-(2,4-difluorophenyl)-1-methyl-3-oxo-1-propenyl)amino)-3-{4-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]phenyl}propanoic acid
- (2S)-2-(((Z)-1-methyl-3-oxo-3-(1,3-thiazol-2-yl)-1-propenyl)amino)-3-{4-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]phenyl}propanoic acid
- 15 (2S)-2-(((Z)-1-methyl-3-oxo-3-(3-thienyl)-1-propenyl)amino)-3-{4-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]phenyl}propanoic acid
- 20 (2S)-2-(((Z)-1-methyl-3-oxo-3-(2-pyridinyl)-1-propenyl)amino)-3-{4-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]phenyl}propanoic acid
- (2S)-2-(((Z)-1-ethyl-3-(4-fluorophenyl)-3-oxo-1-propenyl)amino)-3-{4-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]phenyl}propanoic acid
- 25 (2S)-2-(((Z)-1-methyl-3-oxo-3-phenyl-1-propenyl)amino)-3-{4-[2-(5-methyl-2-phenyl-1,3-thiazol-4-yl)ethoxy]phenyl}propanoic acid
- (2S)-2-(((Z)-3-(2-fluorophenyl)-1-methyl-3-oxo-1-propenyl)amino)-3-{4-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]phenyl}propanoic acid
- 30 (2S)-2-(((Z)-3-(2,3-difluorophenyl)-1-methyl-3-oxo-1-propenyl)amino)-3-{4-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]phenyl}propanoic acid
- 20 (2S)-2-(((Z)-3-(2-hydroxyphenyl)-1-methyl-3-oxo-1-propenyl)amino)-3-{4-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]phenyl}propanoic acid
- 35 (2S)-3-{4-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]phenyl}-2-(((Z)-3-oxo-3-phenyl-1-propyl-1-propenyl)amino)propanoic acid
- 25 (2S)-2-(((Z)-3-(4-methoxyphenyl)-1-methyl-3-oxo-1-propenyl)amino)-3-{4-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]phenyl}propanoic acid
- 40 (2S)-3-{4-[2-[2-(4-methoxyphenyl)-5-methyl-1,3-oxazol-4-yl]ethoxy]phenyl}-2-(((Z)-1-methyl-3-oxo-3-phenyl-1-propenyl)amino)propanoic acid
- 45 (2S)-2-(((Z)-3-cyclohexyl-1-methyl-3-oxo-1-propenyl)amino)-3-{4-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]phenyl}propanoic acid
- 30 (2S)-3-{4-[2-[2-(4-isopropoxyphenyl)-5-methyl-1,3-oxazol-4-yl]ethoxy]phenyl}-2-(((Z)-1-methyl-3-oxo-3-phenyl-1-propenyl)amino)propanoic acid
- 50



- 5 (2S)-2-(((Z)-1-heptyl-3-oxo-3-phenyl-1-propenyl)amino)-3-{4-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]phenyl}propanoic acid
- (2S)-2-(((Z)-1-methyl-3-(3-methylphenyl)-3-oxo-1-propenyl)amino)-3-{4-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]phenyl}propanoic acid
- 10 (2S)-2-(((Z)-1-ethyl-3-oxo-3-phenyl-1-propenyl)amino)-3-{4-[2-(2-(4-methoxyphenyl)-5-methyl-1,3-oxazol-4-yl)ethoxy]phenyl}propanoic acid
- (2S)-3-{4-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]phenyl}-2-(((Z)-3-oxo-3-phenyl-1-(trifluoromethyl)-1-propenyl)amino)propanoic acid
- 15 (2S)-3-{4-[2-(5-methyl-2-phenyl-1,3-thiazol-4-yl)ethoxy]phenyl}-2-(((Z)-3-oxo-3-phenyl-1-(trifluoromethyl)-1-propenyl)amino)propanoic acid
- 20 (2S)-2-(((Z)-1-ethyl-3-oxo-3-[4-(trifluoromethyl)phenyl]-1-propenyl)amino)-3-{4-[2-(2-(4-fluorophenyl)-5-methyl-1,3-oxazol-4-yl)ethoxy]phenyl}propanoic acid
- (2S)-2-(((Z)-1-ethyl-3-oxo-3-[4-(trifluoromethyl)phenyl]-1-propenyl)amino)-3-{4-[2-(2-(4-isopropoxyphenyl)-5-methyl-1,3-oxazol-4-yl)ethoxy]phenyl}propanoic acid
- 25 (2S)-2-(((Z)-1-ethyl-3-oxo-3-[4-(trifluoromethyl)phenyl]-1-propenyl)amino)-3-{4-[2-(2-(4-methoxyphenyl)-5-methyl-1,3-oxazol-4-yl)ethoxy]phenyl}propanoic acid
- 30 (2S)-3-{4-[2-(2-(4-fluorophenyl)-5-methyl-1,3-oxazol-4-yl)ethoxy]phenyl}-2-(((Z)-3-oxo-3-phenyl-1-(trifluoromethyl)-1-propenyl)amino)propanoic acid
- 35 (2S)-2-(((Z)-1-methyl-3-oxo-3-phenyl-1-propenyl)amino)-3-{4-[(5-methyl-2-phenyl-1,3-oxazol-4-yl)methoxy]phenyl}propanoic acid
- (2S)-3-{4-[2-(5-ethyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]phenyl}-2-(((Z)-1-methyl-3-oxo-3-phenyl-1-propenyl)amino)propanoic acid
- 40 (2S)-2-(((Z)-1-methyl-3-oxo-3-phenyl-1-propenyl)amino)-3-{4-[2-(2-phenyl-5-propyl-1,3-oxazol-4-yl)ethoxy]phenyl}propanoic acid
- (2S)-2-(((Z)-1-methyl-3-oxo-3-phenyl-1-propenyl)amino)-3-{4-[3-(5-methyl-2-phenyl-1,3-oxazol-4-yl)propoxy]phenyl}propanoic acid
- 45 (2S)-3-{4-[2-(5-ethyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]phenyl}-2-(((Z)-3-oxo-3-phenyl-1-(trifluoromethyl)-1-propenyl)amino)propanoic acid
- 50 (2S)-3-{4-[3-(5-methyl-2-phenyl-1,3-oxazol-4-yl)propoxy]phenyl}-2-(((Z)-3-oxo-3-phenyl-1-(trifluoromethyl)-1-propenyl)amino)propanoic acid

- 5 (2S)-3-{4-[2-(5-ethyl-2-phenyl-1,3-thiazol-4-yl)ethoxy]phenyl}-2-[[{(Z)-3-oxo-3-phenyl-1-(trifluoromethyl)-1-propenyl]amino}propanoic acid
- (2S)-3-{4-[2-[5-ethyl-2-(4-fluorophenyl)-1,3-oxazol-4-yl]ethoxy]phenyl}-2-[[{(Z)-  
10 3-oxo-3-phenyl-1-(trifluoromethyl)-1-propenyl]amino}propanoic acid
- 5 (2S)-3-{4-[3-[5-ethyl-2-(4-fluorophenyl)-1,3-oxazol-4-yl]propoxy]phenyl}-2-[[{(Z)-3-oxo-3-phenyl-1-(trifluoromethyl)-1-propenyl]amino}propanoic acid
- (2S)-3-{4-[2-[5-ethyl-2-(4-fluorophenyl)-1,3-thiazol-4-yl]ethoxy]phenyl}-2-[[{(Z)-  
15 3-oxo-3-phenyl-1-(trifluoromethyl)-1-propenyl]amino}propanoic acid
- (2S)-3-{4-[3-[5-ethyl-2-(4-fluorophenyl)-1,3-oxazol-4-yl]propoxy]phenyl}-2-  
10 [[{(Z)-1-ethyl-3-oxo-3-[4-(trifluoromethyl)phenyl]-1-propenyl]amino}propanoic acid
- 20 (2S)-2-[[{(Z)-1-ethyl-3-oxo-3-[4-(trifluoromethyl)phenyl]-1-propenyl]amino}-3-{4-[3-(5-ethyl-2-phenyl-1,3-oxazol-4-yl)propoxy]phenyl}propanoic acid
- (2S)-3-{4-[3-[5-ethyl-2-(4-fluorophenyl)-1,3-oxazol-4-yl]propoxy]phenyl}-2-  
25 [[{(Z)-1-methyl-3-oxo-3-[4-(trifluoromethyl)phenyl]-1-propenyl]amino}propanoic acid
- 30 (2S)-3-{4-[3-(5-ethyl-2-phenyl-1,3-oxazol-4-yl)propoxy]phenyl}-2-[[{(Z)-1-methyl-3-oxo-3-[4-(trifluoromethyl)phenyl]-1-propenyl]amino}propanoic acid
- (2S)-3-{4-[3-(5-ethyl-2-phenyl-1,3-oxazol-4-yl)propoxy]phenyl}-2-[[{(Z)-3-oxo-3-  
20 phenyl-1-(trifluoromethyl)-1-propenyl]amino}propanoic acid
- (2S)-3-{4-[[5-ethyl-2-phenyl-1,3-thiazol-4-yl]methoxy]phenyl}-2-[[{(Z)-3-oxo-3-  
35 phenyl-1-(trifluoromethyl)-1-propenyl]amino}propanoic acid
- (2S)-3-{4-[[5-ethyl-2-phenyl-1,3-thiazol-4-yl]methoxy]phenyl}-2-[[{(Z)-1-methyl-  
3-oxo-3-[4-(trifluoromethyl)phenyl]-1-propenyl]amino}propanoic acid
- 25 (2S)-3-{4-[[5-ethyl-2-phenyl-1,3-oxazol-4-yl]methoxy]phenyl}-2-[[{(Z)-3-oxo-3-phenyl-1-(trifluoromethyl)-1-propenyl]amino}propanoic acid
- 40 (2S)-3-{4-[[5-ethyl-2-(4-fluorophenyl)-1,3-oxazol-4-yl]methoxy]phenyl}-2-[[{(Z)-3-oxo-3-phenyl-1-(trifluoromethyl)-1-propenyl]amino}propanoic acid
- 45 (2S)-2-[[{(Z)-1-ethyl-3-oxo-3-[4-(trifluoromethyl)phenyl]-1-propenyl]amino}-3-{4-[[5-ethyl-2-phenyl-1,3-oxazol-4-yl]methoxy]phenyl}propanoic acid
- 30 (2S)-3-{4-[[5-ethyl-2-(4-fluorophenyl)-1,3-oxazol-4-yl]methoxy]phenyl}-2-[[{(Z)-1-ethyl-3-oxo-3-[4-(trifluoromethyl)phenyl]-1-propenyl]amino}propanoic acid
- 50

(2S)-3-[4-[(5-ethyl-2-phenyl-1,3-oxazol-4-yl)methoxy]phenyl]-2-[[[(Z)-1-methyl-3-oxo-3-[4-(trifluoromethyl)phenyl]-1-propenyl]amino]propanoic acid  
(2S)-3-[4-[(5-ethyl-2-(4-fluorophenyl)-1,3-oxazol-4-yl)methoxy]phenyl]-2-[[[(Z)-1-methyl-3-oxo-3-[4-(trifluoromethyl)phenyl]-1-propenyl]amino]propanoic acid  
(2S)-2-[[[(Z)-1-ethyl-3-oxo-3-[4-(trifluoromethyl)phenyl]-1-propenyl]amino]-3-[4-[(5-ethyl-2-phenyl-1,3-thiazol-4-yl)methoxy]phenyl]propanoic acid  
(2S)-3-[4-[2-[5-ethyl-2-(4-fluorophenyl)-1,3-oxazol-4-yl]ethoxy]phenyl]-2-[[[(Z)-1-ethyl-3-oxo-3-[4-(trifluoromethyl)phenyl]-1-propenyl]amino]propanoic acid  
(2S)-3-[4-[2-[5-ethyl-2-(4-fluorophenyl)-1,3-thiazol-4-yl]ethoxy]phenyl]-2-[[[(Z)-1-ethyl-3-oxo-3-[4-(trifluoromethyl)phenyl]-1-propenyl]amino]propanoic acid  
(2S)-2-[[[(Z)-1-ethyl-3-oxo-3-[4-(trifluoromethyl)phenyl]-1-propenyl]amino]-3-[4-[2-(5-ethyl-2-phenyl-1,3-thiazol-4-yl)ethoxy]phenyl]propanoic acid,  
and pharmaceutically acceptable salts and solvates thereof.

12. A compound of Claim 1 selected from the group consisting of  
(2S)-2-[[[(Z)-1-methyl-3-oxo-3-phenyl-1-propenyl]amino]-3-[4-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]phenyl]propanoic acid  
(2S)-2-[[[(Z)-1-ethyl-3-oxo-3-phenyl-1-propenyl]amino]-3-[4-[2-(5-methyl-2-(4-(trifluoromethyl)phenyl)-1,3-oxazol-4-yl)ethoxy]phenyl]propanoic acid  
(2S)-2-[[[(Z)-1-methyl-3-oxo-3-(2,3,4-trifluorophenyl)-1-propenyl]amino]-3-[4-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]phenyl]propanoic acid  
(2S)-2-[[[(Z)-1-methyl-3-(4-nitrophenyl)-3-oxo-1-propenyl]amino]-3-[4-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]phenyl]propanoic acid  
(2S)-2-[[[(Z)-1-methyl-3-oxo-3-[4-(trifluoromethyl)phenyl]-1-propenyl]amino]-3-[4-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]phenyl]propanoic acid  
(2S)-2-[[[(Z)-1-methyl-3-oxo-3-(2,4,5-trifluorophenyl)-1-propenyl]amino]-3-[4-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]phenyl]propanoic acid  
(2S)-2-[[[(Z)-1-ethyl-3-oxo-3-phenyl-1-propenyl]amino]-3-[4-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]phenyl]propanoic acid  
(2S)-3-[4-[2-[2-(4-fluorophenyl)-5-methyl-1,3-thiazol-4-yl]ethoxy]phenyl]-2-[[[(Z)-3-oxo-3-phenyl-1-(trifluoromethyl)-1-propenyl]amino]propanoic acid

- 5 (2S)-2-(((Z)-1-ethyl-3-oxo-3-[4-(trifluoromethyl)phenyl]-1-propenyl)amino)-3-(4-{2-[2-(4-fluorophenyl)-5-methyl-1,3-thiazol-4-yl]ethoxy}phenyl)propanoic acid
- 10 (2S)-2-(((Z)-1-butyl-3-oxo-3-phenyl-1-propenyl)amino)-3-{4-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]phenyl}propanoic acid
- 5 (2S)-2-(((Z)-3-(4-chlorophenyl)-1-methyl-3-oxo-1-propenyl)amino)-3-{4-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]phenyl}propanoic acid
- 15 (2S)-3-{4-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]phenyl}-2-(((Z)-3-oxo-3-phenyl-1-propyl-1-propenyl)amino)propanoic acid
- 10 (2S)-3-{4-[2-(5-methyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]phenyl}-2-(((Z)-3-oxo-3-phenyl-1-(trifluoromethyl)-1-propenyl)amino)propanoic acid
- 20 (2S)-2-(((Z)-1-ethyl-3-oxo-3-[4-(trifluoromethyl)phenyl]-1-propenyl)amino)-3-(4-{2-[2-(4-fluorophenyl)-5-methyl-1,3-oxazol-4-yl]ethoxy}phenyl)propanoic acid
- 25 (2S)-2-(((Z)-1-ethyl-3-oxo-3-[4-(trifluoromethyl)phenyl]-1-propenyl)amino)-3-(4-{2-[2-(4-methoxyphenyl)-5-methyl-1,3-oxazol-4-yl]ethoxy}phenyl)propanoic acid
- 30 (2S)-3-(4-{2-[2-(4-fluorophenyl)-5-methyl-1,3-oxazol-4-yl]ethoxy}phenyl)-2-(((Z)-3-oxo-3-phenyl-1-(trifluoromethyl)-1-propenyl)amino)propanoic acid
- 20 (2S)-2-(((Z)-1-methyl-3-oxo-3-phenyl-1-propenyl)amino)-3-{4-[(5-methyl-2-phenyl-1,3-oxazol-4-yl)methoxy]phenyl}propanoic acid
- 35 (2S)-3-{4-[2-(5-ethyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]phenyl}-2-(((Z)-1-methyl-3-oxo-3-phenyl-1-propenyl)amino)propanoic acid
- (2S)-2-(((Z)-1-methyl-3-oxo-3-phenyl-1-propenyl)amino)-3-{4-[3-(5-methyl-2-phenyl-1,3-oxazol-4-yl)propoxy]phenyl}propanoic acid
- 25 (2S)-3-{4-[2-(5-ethyl-2-phenyl-1,3-oxazol-4-yl)ethoxy]phenyl}-2-(((Z)-3-oxo-3-phenyl-1-(trifluoromethyl)-1-propenyl)amino)propanoic acid
- 40 (2S)-3-{4-[2-(5-ethyl-2-phenyl-1,3-thiazol-4-yl)ethoxy]phenyl}-2-(((Z)-3-oxo-3-phenyl-1-(trifluoromethyl)-1-propenyl)amino)propanoic acid
- 45 (2S)-3-(4-{2-[5-ethyl-2-(4-fluorophenyl)-1,3-oxazol-4-yl]ethoxy}phenyl)-2-(((Z)-3-oxo-3-phenyl-1-(trifluoromethyl)-1-propenyl)amino)propanoic acid
- 30 (2S)-3-(4-{3-[5-ethyl-2-(4-fluorophenyl)-1,3-oxazol-4-yl]propoxy}phenyl)-2-(((Z)-3-oxo-3-phenyl-1-(trifluoromethyl)-1-propenyl)amino)propanoic acid
- 50 (2S)-3-(4-{3-[5-ethyl-2-(4-fluorophenyl)-1,3-oxazol-4-yl]propoxy}phenyl)-2-(((Z)-3-oxo-3-phenyl-1-(trifluoromethyl)-1-propenyl)amino)propanoic acid

(2S)-3-(4-(2-[5-ethyl-2-(4-fluorophenyl)-1,3-thiazol-4-yl]ethoxy)phenyl)-2-(((Z)-3-oxo-3-phenyl-1-(trifluoromethyl)-1-propenyl)amino)propanoic acid  
(2S)-3-(4-(3-[5-ethyl-2-(4-fluorophenyl)-1,3-oxazol-4-yl]propoxy)phenyl)-2-(((Z)-1-ethyl-3-oxo-3-[4-(trifluoromethyl)phenyl]-1-propenyl)amino)propanoic acid  
(2S)-2-(((Z)-1-ethyl-3-oxo-3-[4-(trifluoromethyl)phenyl]-1-propenyl)amino)-3-(4-[3-(5-ethyl-2-phenyl-1,3-oxazol-4-yl)propoxy]phenyl)propanoic acid  
(2S)-3-(4-(3-[5-ethyl-2-(4-fluorophenyl)-1,3-oxazol-4-yl]propoxy)phenyl)-2-(((Z)-1-methyl-3-oxo-3-[4-(trifluoromethyl)phenyl]-1-propenyl)amino)propanoic acid  
(2S)-3-(4-(3-(5-ethyl-2-phenyl-1,3-oxazol-4-yl)propoxy)phenyl)-2-(((Z)-1-methyl-3-oxo-3-[4-(trifluoromethyl)phenyl]-1-propenyl)amino)propanoic acid  
(2S)-3-(4-[[5-ethyl-2-(4-fluorophenyl)-1,3-oxazol-4-yl]methoxy]phenyl)-2-(((Z)-3-oxo-3-phenyl-1-(trifluoromethyl)-1-propenyl)amino)propanoic acid  
(2S)-3-(4-[[5-ethyl-2-phenyl-1,3-oxazol-4-yl]methoxy]phenyl)-2-(((Z)-1-methyl-3-oxo-3-[4-(trifluoromethyl)phenyl]-1-propenyl)amino)propanoic acid  
(2S)-3-(4-[[5-ethyl-2-(4-fluorophenyl)-1,3-oxazol-4-yl]methoxy]phenyl)-2-(((Z)-1-methyl-3-oxo-3-[4-(trifluoromethyl)phenyl]-1-propenyl)amino)propanoic acid  
(2S)-3-(4-(2-[5-ethyl-2-(4-fluorophenyl)-1,3-thiazol-4-yl]ethoxy)phenyl)-2-(((Z)-1-ethyl-3-oxo-3-[4-(trifluoromethyl)phenyl]-1-propenyl)amino)propanoic acid  
and pharmaceutically acceptable salts and solvates thereof.

13. A compound of any of Claims 1-12 wherein said compound is a dual activator of hPPAR $\gamma$  and hPPAR $\alpha$ .

14. A pharmaceutical composition comprising a therapeutically effective amount of a compound of any of Claims 1-13.

15. A pharmaceutical composition of Claim 14 further comprising a pharmaceutically acceptable diluent or carrier.

16. A compound according to any of claims 1-13 for use in therapy.

5                   17. A method for the treatment or prevention of a hPPAR $\alpha$  and/or hPPAR $\gamma$  mediated disease, risk factor, or condition, comprising administration of a composition of Claim 14 or Claim 15 to a patient in need of such prevention or treatment.

10                   5                   18. Use of a compound of any of Claims 1-13 for the manufacture of a medicament for the treatment of a hPPAR $\gamma$  and/or hPPAR $\alpha$  mediated disease, risk factor, or condition.

15                   10                   19. The method of Claim 17 or the use of Claim 18 wherein said disease, risk factor, or condition is hyperglycemia, dyslipidemia, or type II diabetes mellitus including associated diabetic dyslipidemia.

25

30

35

40

45

50

55

# INTERNATIONAL SEARCH REPORT

International Application No.  
PCT/EP 99/05666

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D263/32 C07D277/24 A61K31/421 A61K31/426 C07D413/12  
C07D417/12 A61K31/422 A61K31/427

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 97 31907 A (GLAXO GROUP LTD ) 4 September 1997 (1997-09-04) cited in the application claims	1-9, 13-19
Y	WO 96 38415 A (SUMITOMO METAL INDUSTRIES LTD) 5 December 1996 (1996-12-05) see abstract and formula 1	1-9, 13-19
A	WO 91 19702 A (PFIZER INC) 26 December 1991 (1991-12-26) claims	1-9, 13-19

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

2 December 1999

Date of mailing of the international search report

10/12/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Henry, J

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/EP 99/05666

## Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 17-19  
because they relate to subject matter not required to be searched by this Authority, namely:  
Remark: Although claims 17 and 19 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this International application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.



# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No  
PCT/EP 99/05666

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9731907 A	04-09-1997	AU 2093597 A	16-09-1997
		BG 102792 A	31-08-1999
		CA 2247443 A	04-09-1997
		CN 1218460 A	02-06-1999
		CZ 9802750 A	13-01-1999
		EP 0888317 A	07-01-1999
		HR 970110 A	30-04-1998
		NO 983940 A	27-10-1998
		PL 328871 A	01-03-1999
		SK 116398 A	13-04-1999
WO 9638415 A	05-12-1996	JP 8325263 A	10-12-1996
		AU 5779196 A	18-12-1996
WO 9119702 A	26-12-1991	US 5089514 A	18-02-1992
		AT 149156 T	15-03-1997
		AU 646052 B	03-02-1994
		AU 7995691 A	07-01-1992
		CA 2084898 A	15-12-1991
		DE 69124798 D	03-04-1997
		DE 69124798 T	12-06-1997
		DK 533781 T	07-07-1997
		EP 0533781 A	31-03-1993
		ES 2098356 T	01-05-1997
		FI 925640 A	11-12-1992
		GR 3022714 T	30-06-1997
		HU 65603 A	28-07-1994
		IL 98447 A	31-12-1995
		JP 2581523 B	12-02-1997
		JP 7149636 A	13-06-1995
		JP 7005513 B	25-01-1995
		NO 924799 A	14-12-1992
		NZ 238528 A	27-07-1993
		PT 97950 A, B	31-03-1992
		US 5438074 A	01-08-1995
		US 5306726 A	26-04-1994